

03.11 – Proteins of Medical Interest

PS-03.11.01 THE CRYSTAL STRUCTURE OF PORCINE PANCREATIC SPASMOLYTIC POLYPEPTIDE; A MEMBER OF A NOVEL FAMILY OF GROWTH FACTORS. By Amitabha De, M. A. Gorman, D. Brown*, M. R. Sanderson*, M. Carr[†], A. N. Lane[‡], and P. S. Freemont, Protein Structure Laboratory, Imperial Cancer Research Fund, *The Randall Institute, King's College, [†]National Institute of Medical Research, London, U.K.

Porcine pancreatic spasmolytic polypeptide (PSP) belongs to a family of novel growth factor-like polypeptides which have been isolated from a variety of sources and includes spasmolysin (frog; Hoffmann, W., *J. Biol. Chem.*, 1988, 263, 7686-7690). APEG (frog; Gmachl, M. et al., *FEBS Lett.*, 1990, 260, 145-148), hSP (man; Tomasetto, C. et al., *EMBO J.*, 1990, 9, 407-414) and pS2 (man; Rio, M. C. et al., *Science*, 1988, 241, 705-708). PSP is produced in large amounts in the pancreas and has been shown to inhibit both intestinal motility and gastric acid secretion suggesting involvement in the exocrine function of the pancreas, Jorgensen, K. D. et al., *Regul. Peptides*, 1982, 3, 231-243). Recent studies of the human analogue hSP suggests a role in the healing of damaged endodermally-derived tissue, such as in gastrointestinal ulcers (Wright, N. A. et al., *Nature* 1990, 343, 82-85). PSP (106 amino acids, 11,700 daltons) is a monomer and comprises seven disulphides and two highly homologous domains, residues 8-47 and 58-96, which are approximately 50% identical (Thim, L. et al., *Biochim. Biophys. Acta*, 1985, 827, 410-418). Each domain has three disulphide bonds which gives rise to the proposed three looped "trefoil-like" motif which is distinct from any other characterised growth factors (Thim, L., *FEBS Lett.*, 1989, 250, 85-90). These repeated domains are also seen in human hSP and a single domain is found in ITF (intestinal trefoil factor) from rat (81 residues) and human pS2 (60 residues). Understanding the role of pS2 is of particular importance since approximately 50% of oestrogen dependent human breast tumours secrete pS2 (Brown, A. M. C. et al., *P.N.A.S.* 1984, 81, 6344-6348; Foekens, J. A. et al., *Cancer Res.*, 1990, 50, 3832-3837) whereas in normal breast tissue there is no significant expression of pS2 (Rio, M. C. et al., *Science*, 1988, 241, 705-708). However, as yet the molecular role of pS2 is still unknown.

In order to understand the molecular function of PSP, hSP and more importantly pS2, we have grown crystals of PSP suitable for high resolution structural analysis (Gorman, M. A. et al., *J. Mol. Biol.*, 1992, 228, 991-994) and have now solved the structure by the single isomorphous replacement method combined with solvent modification. The preliminary model and protein fold was interpreted with the help of the recently determined NMR structure of one domain of PSP (Carr, M. D., *Biochemistry*, 1992, 31, 1998-2004) which proved vital in assigning unambiguous parts of the structure. The structure comprises two domains, as suggested by the sequence, which are related by an approximate two-fold axis. Each domain consists of a 7 residue α -helix followed by a short anti-parallel β -strand which gives rise to three exposed loop regions between secondary structure elements. The topology of PSP is novel and has not been observed in any other protein to date. The crystal structure of PSP will be presented and compared to the NMR solution structure and both structures will be discussed in terms of the function of PSP as a growth factor.

PS-03.11.02 PRELIMINARY CRYSTALLOGRAPHIC STUDIES OF MOUSE LIVER GLUTATHIONE S-TRANSFERASE Yb1. By Q. Zhao*, J. Hayes, R. Wolf and H. Driessen, Imperial Cancer Research Fund, Unit of Structural Molecular Biology, Birkbeck College, London WC1E 7HX, UK

The glutathione S-transferases (GST) are a family of enzymes which play a critical role in cellular detoxification processes of mutagens, carcinogens, and other noxious chemical substances and are implicated in the development of drug resistance to electrophilic anticancer drugs, pesticides and herbicides. The enzymes catalyse the nucleophilic addition of the thiol group (-SH) of reduced glutathione (GSH) to a large number of structurally diverse electrophiles making them more polar. A second function for GST's is that they serve as intracellular carrier proteins, binding with high affinity a number of hydrophobic compounds such as heme, bilirubin and bile salts.

In order to understand the mechanism by which the enzymes bind GSH and to establish structure/function relationships, with an eye on selective inhibition of GSTs in anticancer chemotherapy, we have crystallised mouse liver GST isoenzyme Yb1 (μ class) for three-dimensional structure analysis using single crystal X-ray diffraction methods. The enzyme has been crystallized by the vapour diffusion method in the presence of octyl-glutathione and octyl-glucopyranoside using polyethylene glycol 6000. The crystals are monoclinic, space group P2₁, cell dimensions a=79.52 Å, b=74.81 Å, c=78.86 Å, $\alpha = \gamma = 90.0^\circ$, $\beta = 90.07^\circ$ and diffract to at least 2.5 Å resolution on a rotating anode source. Two dimers in the asymmetric unit ($V_m = 2.27$) interact around 2₁ screw axes in the pseudo-orthorhombic space group P2₁2₁2₁, while the intradimer two-fold axes are at an angle of 12.5° and 77.5° from the b-axis in the b-c plane.

An initial attempt to solve the structure of mouse Yb1 with the technique of molecular replacement using the pig π coordinates (Reinemer, Dirr, Ladenstein, Schäffer, Gally and Huber, *EMBO Journal*, 10, 1997-2005, 1991) has failed. The sequence identity is only 32.9% in this case. The structure of mouse Yb1 has now been solved to 3.0 Å with molecular replacement using the coordinates of one homo-dimer of the rat Yb1 (μ class) structure (Ji, Zhang, Armstrong and Gilliland, *Biochemistry* 31, 10169-10184, 1992). Rat and mouse Yb1 have a sequence identity of 93.1%. Currently the mouse Yb1 structure has been refined to a residual of 21.3% at 2.8 Å. Initial results show that all 4 GST molecules have octyl-GSH bound in the G-site (GSH and its derivatives binding position). The octyl-tails have well-defined electron density and have very similar conformations. The octyl-tail bends away from Tyr (100)S, and from the putative H-site (hydrophobic compounds binding site), and stretches across the cleft to within a distance of 4 Å from side chains on helix 4. The direction of the bending is completely different from that of the hexyl-tail in human.

Detailed analysis has not yet been possible, as the refinement has not been completed. However, there appear to be several areas, where the dimer pairs have different interactions on the surface. These areas do not appear to correlate with the residues where rat and mouse differ and there are clear differences between the positions of the octyl-tail in GSH. Further work is under way to study in more details the substrate binding sites of the enzyme.

PS-03.11.03 THE CRYSTAL STRUCTURE OF CHOLESTEROL OXIDASE COMPLEXED WITH A DEHYDROISOANDROSTERONE SUBSTRATE AT 1.9 Å RESOLUTION by Jiayao Li*, Peter Brick & David Blow Imperial College of Science, Technology and Medicine, London, UK

The crystal structure of Cholesterol Oxidase from *Brevibacterium Sterolicum* complexed with a dehydroisoandrosterone substrate was determined by difference Fourier techniques at 1.9 Å resolution. The K_m measurements and the solubility was no more than 50%. The model refined up to R=14.5% by PROLSQ contains 3877 nonhydrogen protein atom plus a FAD molecule, a steroid molecule and 435 waters. Comparison of this structure with the native enzyme structure (Vrielink, A., et al. 1991, *J. Mol. Biol.* 219, 533-554) shows some interesting features. The substrate binding site which is deeply buried inside the protein molecule and was filled with an ordered lattice of 13 waters is now replaced by density for the