

KEK, Tsukuba, Japan. ^eDepartment of Biology, Tokyo Gakugei University, Tokyo, Japan. E-mail: shio8scp@mbox.nc.kyushu-u.ac.jp

Protocyanin is a complex pigment extracted from flower petals of blue cornflower, *Centaurea cyanus*. The components of protocyanin were recently demonstrated to be anthocyanin (AN), flavone glycoside (FL), Fe³⁺, Mg²⁺ and Ca²⁺ ions^[1]. For X-ray structure determination, protocyanin was reconstructed from the components and crystallized in space group *P2₁2₁2₁* with unit cell dimensions of *a* = 29.7, *b* = 49.2 and *c* = 78.3 Å. Two protocyanin molecules are contained in an asymmetric unit. Data were collected on the beam line 6A at Photon Factory KEK to 1.05 Å resolution.

The refined molecule has pseudo three-fold symmetry and four metals align along the pseudo three-fold axis in order of Ca²⁺, Fe³⁺, Mg²⁺ and Ca²⁺. The four metals are coordinated to six AN and six FL molecules. The inner Fe³⁺ and Mg²⁺ ions are each coordinated to three AN's, respectively, while the outer two Ca²⁺ ions are each coordinated to three FL's. Both AN and FL molecules are self-associated with each other as AN-AN and FL-FL in pair and this hydrophobic association also exists between AN and FL molecules, building copigmentation stacks. Protocyanin is a tetra-metal (Fe³⁺, Mg²⁺, 2Ca²⁺) nuclear complex, a new type of supramolecular pigment.

[1] Takeda K., Osakabe A., Saito S., Furuyama D., Tomita A., Kojima Y., Yamadera M., Sakuta M., *Phytochemistry*, submitted.

Keywords: X-ray structure analysis, pigments, biological molecular complexes

P.05.05.7

Acta Cryst. (2005). A61, C276

Crystal Structures of the Fungal Metabolite Oosporein

Ulrich J. Griesser^a, Ram K. R. Jetti^a, Klaus Wurst^b, Christoph Seger^a, Hermann Stuppner^a, Daniela Erlebach^c, Hermann Strasser^c, ^aInstitute of Pharmacy, University of Innsbruck, Austria. ^bInstitute of Institute of General, Inorganic and Theoretical Chemistry, University of Innsbruck, Austria. ^cInstitute of Microbiology, University of Innsbruck, Austria. E-mail: ulrich.griesser@uibk.ac.at

Oosporein is a symmetrical red coloured 2,5-dihydroxybenzoquinone derivative biosynthesized by a broad variety of soil borne fungi. The compound, being known for almost six decades, is the major secondary metabolite of the entomopathogenic fungi *Beauveria brongniartii* which is successfully applied as a biological control agent against the European cockchafer *Melolontha melolontha*. In the course of isolating and purifying pure oosporein from biological cultures we obtained a dioxane solvate and a non-solvated form which were characterized with different solid state analytical techniques including X-ray diffraction.

The molecular geometry of oosporein is x-shaped with a dihedral angle of 67.8 and 79.9° in the non-solvated form and the dioxane solvate respectively. Surprisingly the two forms crystallize in the same space group (monoclinic, *C₂/c*) showing a similar O-H...O network. The non-solvated form shows two dimensional O-H...O tetrameric layers which are off stacked leading to a densely packed structure. In the dioxane solvate one solvent molecule is involved in the O-H...O hydrogen bond network resembling the overall network of the anhydrous form. This pseudo-tetrameric arrangement results in a large channel along the *c*-axis which is occupied by highly disordered dioxane molecules.

[1] Frank R.L., Clark G.R., Coker J.N., *J. Am. Chem. Soc.*, 1950, 72, 1827.

Keywords: oosporein, natural organic molecules, crystal structure

P.05.06.1

Acta Cryst. (2005). A61, C276

Xanthone Derivatives: Conformational Study and Development of Force Field

Michal Markiewicz^a, Jacek Grochowski^b, Pawel Serda^b, Tadeusz Librowski^c, Henryk Marona^d, Carsten Baecht^e, Michael Knapp^e, Marta Pasenkiewicz-Gierula^f, ^aFaculty of Chemistry, Jagiellonian Univ. (JU), Krakow, Poland. ^bRegional Lab., JU. ^cDept. of Pharmacodynamics, Medical College, JU. ^dDept. of Chemical Technology of Drugs, Medical College, JU. ^eInstitute for Materials

Science, Darmstadt Univ. of Technology, Darmstadt, Germany. ^fDept. of Biophysics, Institute of Molecular Biology and Biotechnology, JU. E-mail: markiewi@chemia.uj.edu.pl

Xanthone derivatives extracted from herbs are important components of homeopathic antibacterial, mycotoxic and cytotoxic medicines. Synthetic substituted xanthones tested against broad spectrum of biological activities revealed: antiinflammatory, cytostatic, antimycotic, and cardiovascular activities. In a series of newly synthesized substituted xanthones, two constitutional isomers, 2-methyl-2-[2-(methyl)-6-xanthonyloxy]-propionic acid, 2-methyl-2-[4-(methyl)-6-xanthonyloxy]-propionic acid, and racemic (RS)-2-[2-(methyl)-6-xanthonyloxy]-propionic acid have shown differentiated antiinflammatory action.

The crystal structures of xanthone derivatives were solved using both single-crystal diffraction and HRPD data recorded with synchrotron radiation. In order to find the native, optimal structures of xanthone derivatives in their natural environment of lipid bilayer, additional force field parameters were obtained using X-ray diffraction data and *ab-initio* calculations.

Keywords: xanthone, *ab-initio* calculations, synchrotron radiation

P.05.07.1

Acta Cryst. (2005). A61, C276

Diffraction and Computational Studies of Hydrogen Bonded Base Paired Systems

Suzanne Harte, Mark Facchini, C.C. Wilson, Department of Chemistry, University of Glasgow, Scotland. E-mail: suzhar@chem.gla.ac.uk

Diffraction methods are often utilised to gain a greater insight hydrogen bonding interactions. In work to date the advantages of single crystal variable temperature x-ray and neutron diffraction, coupled with the imaging capabilities of Fourier difference maps, as complementary techniques for anomalous hydrogen bonding investigations have been highlighted [1]. In addition, recent advances in computational chemistry have enabled calculations to be carried out both on isolated molecules and in the periodic (i.e. crystalline) environment [2].

The main aim of this poster is to discuss the use of complementary methods to promote a better understanding of hydrogen bonding within carboxylic acid dimers and nucleic acid base paired systems. One particular system of interest is 3',5'-di-O-acetylthymidine and various techniques have been utilised to determine the presence of anomalous hydrogen behaviour. A crucial part of this has been a multiple temperature high resolution study carried out on station 9.8 at SRS, Daresbury. Alongside experimental work, recent computational work will highlight how these new methods can augment traditional experimental results. Overall it is hoped that the research presented will again highlight the importance of complementary techniques in crystallographic research.

[1] Parkin A., Harte S. M., Goeta A. E., Wilson C. C., *New J. Chem.*, 2004, 28, 718. [2] Wilson C. C., Morrison C. A., *Chem. Phys. Lett.*, 2002, 362, 85-89.

Keywords: hydrogen bonds, diffraction methods, computation

P.05.07.2

Acta Cryst. (2005). A61, C276-C277

Polymorphism of Crystalline Amino Acids. The Role of Non-covalent Interactions

Elena Boldyreva^{a,b}, Sergei Goryainov^{a,c}, Evgenia Kolesnik^{a,b}, Svetlana Ivashevskaya^{a,d}, Valeri Drebushchak^{a,c}, Tatyana Drebushchak^{a,b}, Hans Ahsbahs^e, Heidrun Sowa^e, ^aREC MDEST Novosibirsk State University, Russia. ^bInstitute of Solid State Chemistry SB RAS, Novosibirsk, Russia. ^cInstitute of Mineralogy SB RAS, Novosibirsk, Russia. ^dInstitute of Geology Karelian Scientific Center RAS, Petrozavodsk, Russia. ^eInstitute of Mineralogy Philipps-University Marburg/Lahn, Germany. E-mail: boldyrev@nsu.ru

The crystals of amino acids are interesting from several points of view – as drugs, as molecular materials (e.g. piezo- and ferroelectrics), but also as biomimetics. Understanding the effects of pressure, temperature, and various chemicals on the crystal structures of these