

Organic and Polymeric Materials, Tokyo Institute of Technology, Japan. E-mail: atsuo.iida@kek.jp

For the direct determination of the microscopic local layer structure in the smectic liquid crystal under the external field, time-resolved synchrotron X-ray μ -diffraction has become a powerful tool [1]. In this presentation, static and dynamic local layer structures and local molecular orientation in the electroclinic effect of the chiral smectic A (SmA) phase were analyzed with μ -diffraction.

An X-ray beam size was about $3 \times 4 \mu\text{m}^2$. The samples were ferroelectric liquid crystals (TK-C101) and measured in the SmA phase. At the high electric field, the compound chevron structure, consisting of the curved vertical chevron and the horizontal chevron, was realized. The molecular orientation measured by the high angle halo-pattern showed the spatially-alternate molecular inclination corresponding to the layer deflection. It was shown that the anchoring effect at the alignment film played an important role for the layer structure. The layer response of the electroclinic effect was similar to that in the high electric field treatment of SmC phase, though the origin of the layer shrinkage was different between them.

[1] a) Takahashi Y., et al, *Jpn. J. Appl. Phys.*, 2001, **40**, 3294; b) *Phys. Rev.* 2003, **E67**, 051706.

Keywords: smectic liquid crystal, time-resolved, X-ray microbeam

P.11.10.1

Acta Cryst. (2005). **A61**, C393

Growth and Properties of KTiOPO₄ Single Crystals doped with Er and Nd

Elena Kharitonova^a, Valentina Voronkova^a, Vladimir Yanovskii^a, Valerii Smirnov^b, ^aPhysics Department, Moscow State University. ^bGeneral Physics Institute, Moscow, Russia. E-mail: harit@polly.phys.msu.ru

KTiOPO₄ (KTP) single crystals are of great interests due to their ferroelectric and nonlinear optical properties. Last time the attention is given to the investigation of KTP-family crystals doped with rare earth and codopant ions as the material for self-doubling [1,2].

In the present work single crystals of KTP codoped with Ln = Er, Nd and Me = Nb, Ta, Ba, Ca, Mg, Bi, Al, Si, KCl were grown by means of flux method and some properties were investigated.

The investigations show that an addition of Nb and Ba leads to decreasing of KTP:Ln:Me optical quality, whereas the quality increases with KCl and Bi and does not markedly depends on Ln.

Maximum luminescence intensity was observed in KTP:Er:Nb crystals. Life time of ⁴F_{3/2}Nd³⁺ and ⁴I_{13/2}Er³⁺ has been found as 250 μ s and 6 ms for KTP:Nd and KTP:Er respectively. In KTP:Ln:Me the life time changes in the area of 1 - 7 ms for ⁴I_{13/2}Er³⁺ and 180 - 300 μ s for ⁴F_{3/2}Nd³⁺ depending on the codopant of ion Me.

The presence of Ln ions in KTP:Ln:Me crystals does not strongly affect on ferroelectric phase transition temperature, but suppresses the dielectric permittivity peak caused by the above-named transition.

[1] Sole R., Nikolov V., Koseva I., et al., *Chem. Mater.*, 1997, **9**, 2745. [2] Carvajal J. J., Nikolov V., Sole R., et al., *Chem. Mater.*, 2002, **14**, 3136.

Keywords: crystal growth, KTiOPO₄, ferroelectrics

P.11.10.2

Acta Cryst. (2005). **A61**, C393

Second-Order Nonlinear Optical Properties of Tetradentate Schiff base Complexes

Alessandra Forni^a, Julieta Gradinaru^b, Vadim Druta^b, Francesca Tessore^c, Sandro Zecchin^d, Silvio Quici^a, Nicolae Gerbeleu^b, ^aISTM-CNR, Milan, Italy. ^bInstitute of Chemistry of the Academy of Sciences of Moldova, Chisinau, R. Moldova. ^cDepartment of Inorganic Chemistry, Milan, Italy. ^dIENI-CNR, Padua, Italy. E-mail: a.forni@istm.cnr.it

Quadratic non-linear optical properties of new unsymmetrical Ni(II), Cu(II), Zn(II) and VO(II) complexes with [N₂O₂] chelate tetradentate ligands are presented. The complexes were synthesized by template condensation of 1-phenylbutane-1,3-dione mono-S-methylisothiosemicarbazone with o-hydroxybenzaldehyde or its 5-

phenylazo-derivative. The crystal structure of some complexes was determined by using synchrotron radiation (XRD1 beamline at ELETTRA, Trieste).

Large values of hyperpolarizability, as obtained for these complexes by the solution-phase dc electric-field-induced second harmonic (EFISH) generation method, together with their high stability constants, make them promising candidates for successful applications in the field of optoelectronic technologies.

The values of theoretical hyperpolarizability, calculated through both Finite Field and Time Dependent DFT methods, are also presented.

Keywords: transition-metal complexes, nonlinear optical properties, theoretical calculations

P.11.10.3

Acta Cryst. (2005). **A61**, C393

Temperature Dependence of Refractive Indices in selected Borates

René Ochrombel, Ladislav Bohatý, Petra Becker, *Institute of Crystallography, University of Cologne, Germany.* E-mail: rene.ochrombel@uni-koeln.de

The knowledge of the temperature dependence of the refractive indices of a nonlinear optical crystal play a key role for its classification as a NLO material. For example, the temperature dependence can be used as a possibility for fine tuning of the phase-matching conditions of a nonlinear optical frequency conversion process. In general it is advantageous to describe the temperature dependence of the refractive indices using the temperature dependence of the polarisation tensor [a_{ij}] (optical dielectric impermeability tensor) $\Delta a_{ij} = b_{ij} \Delta T + c_{ij} \Delta T^2 + \dots$, where [b_{ij}], [c_{ij}], ... describe the linear, quadratic, etc. temperature dependence of the polarisation tensor. In most cases (far from phase transitions) the linear approximation turn out to be an adequate description.

In this work we present a technique for the measurement of temperature dependence of refractive indices based on a Jamin interferometer, which allows to measure optical path length differences as a function of temperature (temperature range: 50...280 K). The knowledge of the refractive indices at a reference temperature T₀ (e.g. room temperature) including the orientation of the indicatrix, and the knowledge of thermal expansion data in the same temperature range are necessary for the evaluation of the experimental data.

As results we present the [b_{ij}] tensors of the non-centrosymmetric borate crystals: Zn₄[O(BO₂)₆] (PG: $\bar{4}3m$), Li₂B₄O₇ (PG: *4mm*), β -BaB₂O₄ (PG: *3m*), PbB₄O₇ (PG: *mm2*), SrB₄O₇ (PG: *mm2*) and BiB₃O₆ (PG: *2*).

Keywords: non-centrosymmetric borates, phase-matching, nonlinear optics

P.11.10.4

Acta Cryst. (2005). **A61**, C393-C394

Nonlinear Optical Properties of Lithium Sulfate Monohydrate, Li₂SO₄ · H₂O

Lkhamsuren Bayarjargal^a, L. Bohatý^a, ^aInstitute of Crystallography, University of Cologne, Zùlpicher Str. 49b, D-50674 Köln, Germany. E-mail: bayarjarl@uni-koeln.de

Lithium sulfate monohydrate crystallizes in the monoclinic polar space group P2₁ [1]. Among the group of non-ferroelectric polar crystals, Li₂SO₄·H₂O possesses the highest pyroelectric coefficient as well as remarkable piezoelectric and electro-optic properties. However, there is only little known about its nonlinear optical properties [2]. Recently, the possibilities of phase-matched second harmonic generation (SHG) in Li₂SO₄·H₂O were analysed in detail [3]. In this work we present the results of our investigation of the SHG. Using the Maker fringe technique and four differently oriented plane slabs of Li₂SO₄·H₂O crystals all eight independent components of the nonlinear optical susceptibility tensor [d_{ijk}] for the fundamental wavelength $\lambda = 1079.5$ nm (Nd:YAP laser) were determined. In comparison to other polar properties of Li₂SO₄·H₂O the coefficients d_{ijk} are surprisingly small; they are one order of magnitude smaller than those of commonly used crystals for frequency conversion of laser light, such as KTiOPO₄, β -BaB₂O₄, LiB₃O₅ or BiB₃O₆. The