

**m05.p01****Advances in charge density studies of ultra high resolution protein structures**

Benoît Guillot<sup>a</sup>, Angélique Lagoutte<sup>a</sup>, Christian Jelsch<sup>a</sup>,  
A. Podjarny<sup>b</sup>, Claude Lecomte<sup>a</sup>

<sup>a</sup>LCM3B, Université H. Poincaré, Nancy, France <sup>b</sup>IGBMC, Strasbourg, France. E-mail: benoit.guillot@lcm3b.uhp-nancy.fr

**Keywords:** charge density studies, high-resolution protein structure, electrostatic properties

The number of proteins and nucleic acids structures solved at atomic and subatomic resolution is increasing continuously. Atomic resolution allows observation of chemical group protonation state and subatomic resolution reveals electron density fine details related to the deformation of the electron cloud due to chemical bonding and intermolecular interactions. As in the field of small molecules crystallography, the use of a non spherical model of atomic electron density allows to take into account these features in the refinement. For this purpose the Hansen & Coppens multipolar model [1] has been implemented in the MoPro software [2], along with methods from small molecules charge density studies and biological crystallography fields. Here we will show how these methods, and using multipolar parameter transferability principle [3], allow biological macromolecules charge density analyses and electrostatic properties computation. We will illustrate our results on several high resolution protein and nucleic acid structures, including Human aldose reductase/inhibitors complexes [4].

[1] Hansen, N.K.; Coppens, P., *Acta. Cryst.* 1978. A34, 909-921.

[2] Jelsch, C.; Guillot, B.; Lagoutte A.; Lecomte, C., *J. Appl. Cryst.* 2005. 38, 38-54.

[3] Jelsch, C.; Pichon-Pesme, V.; Lecomte, C.; Aubry, A., *Acta. Cryst.*, 1998, D54, 1306-1318.

[4] Howard, E. *et. al.*, *Prot. Struct. Funct. & Gen.* 2004. 55, 792-804.

**m06.p01****Structural studies on collagen binding integrin I domains**

Anna-Maria Brandt, J. Santeri Puranen, Yvonne Nymalm,  
Tiina Salminen, Mark S. Johnson

Department of Biochemistry and Pharmacy, Åbo Akademi University, Turku, Finland.

**Keywords:** integrin, I-domain, collagen-binding

Integrins are cell adhesion receptors that mediate cell-cell or cell-extracellular matrix interactions by bidirectional signaling. The  $\alpha\beta$  heterodimeric glycoproteins are composed from 19 different  $\alpha$  subunits and 8 different  $\beta$  subunits. The collagen binding integrin family consists of four collagen receptors that have a common  $\beta 1$  subunit non-covalently bound either to  $\alpha 1$ ,  $\alpha 2$ ,  $\alpha 10$  or  $\alpha 11$  subunit. The  $\alpha$  subunits include an inserted domain of 200 amino acids, also called the I domain, which is the key ligand-binding domain. The I domain alternates between two conformations, which are important for regulating the affinity of the ligand. The  $\alpha I$  domain contains the metal ion-dependent adhesion site (MIDAS), which in the open conformation binds collagen, while in the closed conformation the binding is hindered.

The crystal structure of  $\alpha AI$  and  $\alpha 2I$ , and the complex structure of  $\alpha 2I$  bound to a collagen-like peptide, have been solved [1,2,3]. We have used mutagenesis to lock the structure of  $\alpha 1I$  in the open conformation and showed that the mutant binds collagen with several times higher affinity than the wild-type. Our aim is to crystallize the active mutant and determine the structure of the open conformation. We have also modeled the open conformation of  $\alpha 1I$  in complex with a collagen-like peptide and characterized the binding of the ligand [4].

[1] Salminen TA., Nymalm Y., Kankare J., Käpylä J., Heino J., Johnson MS. (1999) *Acta Crystallogr D Biol Crystallogr.* 55:1365-7.

[2] Emsley J., King SL., Bergelson JM., Liddington RC. (1997) *J Biol Chem.* 272:28512-7.

[3] Emsley J., Knight CG., Farndale RW., Barnes MJ., Liddington RC. (2000) *Cell.* 101:47-56.

[4] Nymalm Y., Puranen JS., Nyholm TK., Käpylä J., Kidron H., Pentikäinen OT., Airene TT., Heino J., Slotte JP., Johnson MS., Salminen TA. (2004) *J Biol Chem.* 279:7962-70.