

[2] I. Goncharenko, P. Cadavez-Peres, O. Makarova, T. Le Bihan, M. Mezouar, *Phys. Rev. B*, 2003, 68, 214418.

[3] N. Ashcroft, *Phys. Rev. Lett.*, 200492, 187002.

[4] I. Goncharenko, *High Press. Res.*, 2007, 27 183

MS22 O4

Single crystal studies of pure elements at high pressure

Lars F. Lundegaard and Malcolm I. McMahon, *SUPA, School of Physics and Centre for Science at Extreme Conditions, The University of Edinburgh, Edinburgh, UK.*
E-mail: lars.lundegaard@ed.ac.uk

Keywords: high-pressure, elements, incommensurate

Although the technique of high-pressure single-crystal diffraction has been used to study pure elements for many years [1], recent developments, such as improved diamond anvil cell design combined with CCD-equipped diffractometers and synchrotron sources, have allowed the structure solution of a number of complex new phases [2,3].

In this talk we will describe the technique of high-pressure single-crystal diffraction, and discuss its advantages and limitations. To illustrate the power of the technique, we will present recent results on the full modulated structure of the incommensurate composite structure of Rb-IV, the structure of Ba-IVb and the structure of epsilon-oxygen [4].

Finally, we will describe our on-going development of techniques that will enable us to do combined high-pressure high-temperature single-crystal diffraction, and single-crystal diffraction at pressures of 50GPa and above.

[1] D. Schiferl, D. T. Cromer & R. L. Mills, *Acta Cryst.* (1981), B37, 1329-1332.

[2] M. I. McMahon, R. J. Nelmes & S. Rekhi, *Phys. Rev. Lett.* (2001), 87, 255502.

[3] R. J. Nelmes, M. I. McMahon, J. S. Loveday & S. Rekhi, *Phys. Rev. Lett.* (2002), 88, 155503.

[4] L. F. Lundegaard, G. Weck, M. I. McMahon, S. Desgreniers & P. Loubeyre, *Nature* (2006), 443, 201-204.

MS22 O5

Electric field induced lattice displacements in BiB₃O₆ crystals,

Ullrich Pietsch^a, Oleg Schmidt^a, Semen Gorfman^a, Petra Becker^b, Ladislav Bohaty^b ^a*Solid State Physics Department of Physics, University of Siegen, Germany.* ^b*Institute of Crystallography, University of Cologne, Germany.* E-mail: pietsch@physik.uni-siegen.de

Keywords: Charge density, Piezoelectric crystals, Electric field induced

Monoclinic BiB₃O₆ is a piezoelectric material with exceptional large piezoelectric coefficients [1]. At this material X-ray charge density analysis is highly affected by the huge difference in electronic numbers of constituting elements and by secondary extinction. However, measurements of intensity variation of selected Bragg reflection caused by an external high electric field provide information about the reorganization of atomic positions within the unit cell and subsequently about a change of chemical bonds [2]. For BiB₃O₆ we have measured several Bragg reflections under influence of an external electric field up to 20 kV/cm using synchrotron radiation. The measured change in angular positions of Bragg reflections is caused by the external piezoelectric effect. Measuring the respective peak shifts at three differently oriented crystal plates we have deduced all eight independent coefficients of the piezoelectric tensor of the material [3]. In addition, the intensity is changed due to the internal piezoelectric effect, i.e. the reorganization of atomic arrangement within the unit cell. Preliminary modelling suggested an effect less than 1%. However, tuning the probing wave length close to the absorption edge of Bi atoms we could measure the linear dependence of intensity variation as a function of the applied field at a [100] oriented crystal plate. Due to the small number of measured reflections we used a rough model for data interpretation, a displacement of rigid lattice of BO₃ units against the fixed Bi atoms. Normalized to E=1kV/mm we deduced a displacement vector $\Delta r = -(14, 2 a_1 + 47, 5 a_3)10^{-5}$, given in relative coordinates of the monoclinic system with $a_{1,2,3}$ as unit vectors. Since the external field was applied in direction $E = 0, 146 a_1 + 0, 043 a_3$ the displacement of the negatively charged BO₃ units are directed by an angle of about 75 degree with respect to the plane of positively charged Bi atoms.

[1] Haussühl, S.; Bohaty, L. ; Becker, P.: Piezoelectric and elastic properties of the nonlinear optical material bismuth triborate, BiB₃O₆. In: *Appl. Phys. A* 82 (2006),S. 495–502

[2] Gorfman, S. V.; Tsirelson, V. G.; Pucher, A.; Morgenroth, W.; Pietsch, U.: X-ray diffraction by a crystal in a permanent external electric field: electric-field-induced structural response in α -GaPO₄. *Acta Cryst.* A62 (2006) 1-10.

[3] S. Gorfman, O. Schmidt, U. Pietsch, P. Becker, L. Bohaty. X-Ray Diffraction Study of the Piezoelectric Properties of BiB₃O₆ Single Crystals (2007) *Z. f. Krist.*, accepted