

There is also a very weak intermolecular Sn...O interaction in complex **4** between Sn2 and the uncoordinated oxygen atom of carbonyl group of a neighboring molecule and link the discrete molecular into a 1D chain with the Sn...Sn separation of 8.629 Å. Although the Sn–O distance [Sn2–O2 = 2.975 Å, (BO = 0.1)] is considerably longer than the normal Sn–O covalent bond, it lies in the range of Sn...O distances of 2.61–3.02 Å that have been reported to suggest Sn–O bonding [7]. However, such an elongation of Sn–O bond from the normal covalent bond distance point to a considerable ionic character associated with Sn–O bond.

Furthermore, the intermolecular NH...Cl hydrogen bonds produce two dimensional supramolecular structures in the lattice.

[1] F. Benetello, C.G. Lobbia, M. Mancini, M. Pellei, C. Santini, *Journal of Organometallic Chemistry* **2005**, 690, 1994–2001. [2] H.P.S. Chauhan, N.M. Shaik, *Journal of Inorganic Biochemistry* **2005**, 99, 538–545. [3] D.C. Menezes, F.T. Vieira, G.M. de Lima, J.L. Wardell, M.E. Cortés, M.P. Ferreira, M.A. Soares, A. Vilas Boas, *Applied. Organometallic Chemistry* **2008**, 22, 221–226. [4] K.E. Gubina, V.A. Ovchinnikov, V.M. Amirkhanov, H. Fischer, R. Stumpf, V.V. Skopenko, *Zeitschrift für Naturforschung* **2000**, 55b, 576–582. [5] K. Gholivand, Z. Shariatinia, *Journal of Organometallic Chemistry* **2006**, 691, 4215–4224. [6] K. Gholivand, S. Farshadian, *Inorgica Chimica Acta* **2011**, 368, 111–123. [7] A. R. Forrester, S. J. Garden, R. A. Howie, J. L. Wardell, *Journal of the Chemical Society, Dalton Transactions* **1992**, 2615–2621.

Keywords: crystal structure, carbacylamidophosphate, hydrogen bond

MS61.P30

Acta Cryst. (2011) A67, C616

2-Chloro-N-[4-(dimethylamino)benzylidene]-N-[4-(3-methyl-3-phenylcyclobutyl)-1,3-thiazol-2-yl]acetohydrazide

Ersin İnkaya, Muharrem Dinçer, *Departament of Physics, Ondokuz Mayıs University, 55139, Samsun, (Turkey)*. E-mail: ersin.inkaya@oposta.omu.edu.tr

3- Substituted cyclobutane carboxylic acids exhibit anti-inflammatory and anti-depressant activity [1], as well as having liquid crystal properties [2]. Also, various thiazole derivatives have been shown to possess herbicidal [3], anti-inflammatory [4], anti-microbial [5], and anti-parasitic properties.

The molecular conformation of the title compound, C₂₅H₂₇ClN₄OS, is stabilized by an intramolecular benzylidene C—H...N_{thiazole} hydrogen bond. The thiazole ring makes dihedral angles of 12.0 (3)° and 20.4 (2)°, respectively, with the phenyl and benzene rings, while the phenyl and benzene rings make a dihedral angle of 22.6 (2)°. The crystal packing involves weak intermolecular thiazole C—H...O_{carbonyl} and methyl C—H...π hydrogen-bonding associations

[1] E. Roger, C.J. Pierre, V. Pualette, G. Gerard, J.P. Chepat, G. Robert. *Eur. J. Med. Chem. Chem. Ther.* **1977**, 12, 501–509. [2] E.V. Dehmlow, S.S. Schmidt. *Liebigs Ann. Chem* **1990**, 411–414. [3] H. Foerster, W. Hofer, V. Mues, L. Eue, R.R. Schmidt FRG Patent 2822155, **1979**. [4] S.N. Sawhney, S.K. Arora, J.V. Singh, O.P. Bansal, S.P. Singh, *J. Indian Chem. Soc. Sect. B*, **1978**, 16, 605–609. [5] N. Suzuki, Y. Tanaka, R. Dohmori, *Chem. Pharm. Bull.* **1979**, 27, 1–11. [6] P.I. Slip, M. Closier, Neville, *J. Med. Chem.* **1974**, 17, 207–209.

Keywords: crystallography, single crystal x-ray

MS61.P31

Acta Cryst. (2011) A67, C616

New Copper(I) Polymers Assembled by Dithioether Ligands

Marek M. Kubicki,^a Yoann Rousselin,^a Michael Knorr,^b Fabrice Guyon,^b ^a*Institut of Molecular Chemistry ICMUB UMR CNRS 5260, Université de Bourgogne, BP47870, F-21078 Dijon Cedex, (France)*. ^b*Institut UTINAM UMR CNRS 6213, Université de Franche-Comté, 16 route de Gray, F-25030 Besançon, (France)*. E-mail: marek.kubicki@u-bourgogne.fr

During the last decade there is a spectacular increase of interest in the field of Cu(I)-based polymers due to their rich structural chemistry and photo-physical properties. The networks resulting from CuX/N,P,S-organic ligand self-assembly processes may adopt linear (1D), layered (2D) or three-dimensional (3D) topologies.[1] We present here some new structural features observed in CuX (X=I, Br, Cl) / RS(C)_nSR limited to n=4 systems. The C₄ spacer may be an n-butyl like (R-S-CH₂-CH₂-CH₂-CH₂-S-R) or 2-butylene (R-S-CH₂-CH=CH-CH₂-S-R) in both the Z- and E- geometries. The most largely studied copper(I) halide is CuI because it easily forms the cubane-like Cu₄I₄ motif as a secondary building unit (SBU) responsible for the intense luminescent properties.[2] However, other Cu_nX_m stoichiometries may also be reached, the most frequent is that of Cu₂X₂ rhomboid.

In C₄ spacer n-butyl-like series with Cu₄I₄ SBU the 2D structures are observed with R=tBu and nBu: (Cu₄I₄S₄, 2:1 CuX: ligand ratio). However, the layers therein are not planar, but undulating over the respective crystal lattice planes. The deviations of respective Cu₄I₄ centroids over these planes reaches ± 1.87 Å for R=tBu (Pbcn) and ± 1.81 Å for R=nBu (P21/c).[3] The 2D structure is obtained for R=tBu series with CuBr (P-1), but the SBU therein is rhomboidal Cu₂Br₂ with the overall ratio CuBr:ligand = 1:1. Going to the salt:ligand ratio of 2:1 a dramatic, but spectacular structure transformation is observed in R=tBu series from centrosymmetric P-1 to tetragonal chiral P₄, bearing the Cu₄Br₄(SC₄Slinker)Cu₄Br₄ helices. An overall 3D structure is reached therein through the strong Cu – S(spacer)S – Cu inter-helice interactions.

In the C₄ 2-butene spacer series the E-(trans) and Z-(cis) geometries of RSC₄SR dithioether influence on the composition and topology. E-(cis) isomers of PhSCH₂CH=CHCH₂SPh lead to 2D planar layers in triclinic (P-1) structures with Cu₂Br₂ and Cu₂I₂ SBU's, whereas the cis (Z) isomer gives with CuI a 2D monoclinic (C2/c) luminescent structure with Cu₄I₄ SBUs.

A steric effect of substitutions on Ph group (Me ortho, tBu meta) of dithioether in saturated R'S(nBu)SR' series has been also checked with CuI. The effect is evidenced by the formation of 1D linear ribbons with the coordination of solvent (NCMe) molecules on the copper atoms.

A particular attention will be paid for discussion of Cu-Cu distances. Variable temperature X-ray studies have been performed in several cases in order to find the structure/thermo-luminescent relationships.

[1] M. Knorr, F. Guyon, *Macromolecules Containing Metal and Metal-like Elements, Volume 10: Photophysics and Photochemistry of Metal-Containing Polymers*, Ed. A.S. Abd-El Aziz, C.E. Carraher Jr., C.U. Pittman Jr., M. Zeldin. **2010** John Wiley & Sons, Inc. [2] M. Knorr, A. Pam, A. Kathyr, C. Strohmman, M. M. Kubicki, Y. Rousselin, S.M. Aly, D. Fortin, P.D. Harvey *Inorg. Chem.*, **2010**, 49, 5834–5844. [3] M. Knorr, F. Guyon, M.M. Kubicki, Y. Rousselin, S.M. Aly, P.D. Harvey, *New J. Chem.*, **2011**, DOI: 10.1039/c0nj00923g.

Keywords: copper(I), dithioethers, topology of the network