

[1] Chiavassa et al 2005 *Astron. & Astroph.* 432,547; Posch Th et al 2007 *Astroph. J.* 668, 993 [2] Ferrarotti and Gail 2005 *Astron. & Astrophys.* 430, 959 [3] Thompson et al. 2012 *Journal of Non-Crystalline Solids* 358 (5), 885 [4] Demyk, et al. *A&A*, 2000,364,170; B.T. Draine. 2003 *Annu. Rev. Astron. Astroph.*, 41,1

Keywords: amorphous silicates, carbonates, powder diffraction, PDF

MS18-P2 Single Crystal High Pressure Diffraction at the Advanced Light Source

Christine M. Beavers^{1,2}, Earl F. O'Bannon², Andrew Doran¹,
Martin Kunz¹, Quentin Williams²

1. Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA USA

2. Department of Earth & Planetary Sciences, University of California Santa Cruz, Santa Cruz, CA USA

email: cmbeavers@lbl.gov

Beamline 12.2.2 at the Advanced Light Source, which was previously optimized for high pressure powder diffraction only, has undergone the substantial addition of a dedicated single crystal diffractometer. Positioned upstream of the existing endstation, the new endstation consists of a Stoe Stadi Vari diffractometer equipped with a RDI CMOS detector. The system was configured with high pressure diffraction in mind: a small sphere of confusion (10 μ m), a sample weight capacity compatible with DACs (1kg), and a detector phosphor optimized for 25keV. System details and commissioning experiment results will be shared.

Keywords: high pressure, single crystal