

*Polder maps: Improving OMIT maps for ligand building and validation*Dorothee Claudia Liebschner¹, Nigel Moriarty¹, Pavel Afonine¹, Billy Poon¹, Oleg Sobolev¹, Paul Adams¹¹Lawrence Berkeley National Laboratory, Berkeley, United States

E-mail: dcliebschner@lbl.gov

In macromolecular crystallography, electron density maps are used to build and validate crystallographic models. In particular, OMIT maps are a common tool to verify the presence of ligands in a crystal structure. The simplest way to compute an OMIT map is to exclude the ligand from the model, update the structure factors and compute a residual map. It is then expected that if the ligand is present in the crystal structure, the electron density for the omitted atoms will appear as positive features in this map. However, this is complicated by the flat bulk-solvent model [1], which postulates constant electron density in the areas of the unit cell that are not occupied by the atomic model. Thus, if atoms are removed from the model, the region where they were modelled will be filled with bulk-solvent, and if the density arising from the omitted atoms is weak, then the bulk-solvent model may obscure it further.


A possible solution to this problem is to prevent bulk-solvent from entering the OMIT regions, which may improve the interpretative power of residual maps. This approach is called polder (OMIT) map [2] and the tool is implemented and available in the software suite Phenix [3]. Polder OMIT maps can be particularly useful for displaying weak densities of ligands. Several examples are presented where polder OMIT maps show clearer features than conventional OMIT maps.

OMIT maps for ligand MES 88 in structure 1ABA. The positive and negative mFobs-DFmodel OMIT difference density is displayed in green and red, respectively. (a) OMIT map contoured at $\pm 3\sigma$. (b) Polder map contoured at $\pm 3\sigma$.

[1] Jiang, J. & Brünger, A. (1994). *J. Mol. Biol.*, 243, 100–115

[2] Liebschner, D. et al. (2017). *Acta Cryst. D*, 73, 148–157

[3] Adams, P. D. et al. (2010). *Acta Cryst. D*, 66, 213–221 (2010).



Keywords: [omit maps](#), [ligand placement](#), [bulk solvent model](#)