

atom, O atom and aromatic ring. This model suggests that higher-affinity ligands for the pirenzepine site could be developed if the relative orientation of the N⁺-containing ring were restricted to the common position found in the four structures: pirenzepine, (+)-biperiden, dexetimide and atropine.

The structural comparisons of chiral ligands for the pirenzepine site show both that structural differences in the two types of binding sites are not evident and that structural similarities in the ligands are not detected by the torsion angles of the consistent conformation for anticholinergics. Since the chiral ligands for both of the pirenzepine binding sites are structurally superimposable, the difference between these sites must arise from factors other than simple shape recognition; these factors may be related to induced conformational changes in either the receptor or the ligand. The ligands compared in this study have a close structural match which belies the wide variation in the values of the torsion angles of the consistent anticholinergic conformation. This discrepancy suggests that more detailed models that compare overall shapes and juxtapose similar heteroatoms are required so that a range of chemical types of ligands for a common receptor can be compared.

This work was supported by the Medical Research Council of Canada (grant MA 8087). The author thanks Dr V. Figala and Dr M. Elitze of Byk Gulden Pharmazeutika for providing the samples and the results of their pharmacological tests and Norma E. Duke for collecting the (+)-biperiden data.

References

- ALBERT, A. & SERJEANT, E. P. (1984). *The Determination of Ionization Constants*, p. 151. London: Chapman and Hall.
- CODDING, P. W. (1983). *Can. J. Chem.* **61**, 529-532.
- CODDING, P. W. & JAMES, M. N. G. (1984). *Acta Cryst.* **B40**, 429-434.
- CROMER, D. T. & MANN, J. B. (1968). *Acta Cryst.* **A24**, 321-324.
- ELTZE, M. & FIGALA, V. (1984). Private communication.
- GERMAIN, G., MAIN, P. & WOOLFSON, M. M. (1971). *Acta Cryst.* **A27**, 368-376.
- GUY, J. J. & HAMOR, T. A. (1975). *J. Chem. Soc. Perkin Trans. 2*, pp. 1074-1077.
- HAMMER, R., BERRIE, C. P., BIRDSALL, N. J. M., BURGEN, A. B. V. & HULME, E. C. (1980). *Nature (London)*, **283**, 90-92.
- JOHNSON, C. K. (1970). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee.
- KLYNE, W. & BUCKINGHAM, J. (1978). *Atlas of Stereochemistry*, Vol. 1, 2nd ed. New York: Oxford Univ. Press.
- KUSSATHER, E. & HAASE, J. (1972). *Acta Cryst.* **B28**, 2896-2899.
- MOTHERWELL, S. (1979). *PLUTO*. Program for plotting molecular and crystal structures. Univ. of Cambridge, England.
- PAULING, P. & DATTA, N. (1980). *Proc. Natl Acad. Sci. USA*, **77**, 708-712.
- RUZIC-TOROS, Z. & KOJIC-PRODIC, B. (1983). *Acta Cryst.* **C39**, 93-95.
- SMITH, G. D. (1980). *PROFIT*. A molecular superposition program. Medical Foundation of Buffalo, Buffalo, New York.
- SPEK, A. L. (1976). *Acta Cryst.* **B32**, 870-877.
- STEWART, J. M., MACHIN, P. A., DICKINSON, C. W., AMMON, H. L., HECK, H. & FLACK, H. (1976). The XRAY76 system. Tech. Rep. TR-446. Computer Science Center, Univ. of Maryland, College Park, Maryland.
- STEWART, R. F., DAVIDSON, E. R. & SIMPSON, W. T. (1965). *J. Chem. Phys.* **42**, 3175-3187.
- TRUMMLITZ, G., SCHMIDT, G., WAGNER, H.-U. & LUGER, P. (1984). *Arzneim. Forsch.* **34**, 849-859.
- YAMAMURA, H. I., WATSON, M. & ROESKE, W. R. (1983). *CNS Receptors—From Molecular Pharmacology to Behavior*, edited by P. MANDEL & F. V. DEFEUDIS, pp. 331-336. New York: Raven Press.

International Union of Crystallography

Acta Cryst. (1986). **B42**, 638

Deposition of Data with Brookhaven Protein Data Bank

Acta Crystallographica policy provides authors of biological-macromolecule papers with the option of requesting that their list of structure factors deposited with the Brookhaven Protein Data Bank be granted privileged status for a period of no longer than four years from the date of publication, see *Notes for Authors* [*Acta Cryst.* (1983), **A39**, 174-186]. In a recent paper by Comarmond, Giegé, Thierry,

Moras & Fischer [*Acta Cryst.* (1986), **B42**, 272-280] the atomic coordinates were wrongly assigned this status also. The atomic coordinates are available for immediate distribution.

Currently acceptable formats for the machine-readable media are: magnetic tape with fixed line length and fixed block size in 9 track, ASCII or EBCDIC, 800, 1600 or 6250 CPI; or punched cards with IBM 026 or IBM 029 codes.

Copies of data in machine-readable form are available from the Protein Data Bank at Brookhaven or one of the affiliated centers at Melbourne or Osaka (but no longer at Cambridge).