

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Chloroquinuclidinium chloride

Isha Azizul, Arifin Zainudin and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 27 February 2008; accepted 21 April 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{N}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.040; wR factor = 0.113; data-to-parameter ratio = 13.0.

The cation of the title compound, $\text{C}_7\text{H}_{13}\text{ClN}^+\cdot\text{Cl}^-$, forms a linear hydrogen bond to the chloride anion. The cation is disordered about a mirror plane.

Related literature

For isomeric 4-chloroquinuclidinium chloride, see: Kurahashi *et al.* (1980), which also reports the parent quinuclidinium chloride.



Experimental

Crystal data

 $\text{C}_7\text{H}_{13}\text{ClN}^+\cdot\text{Cl}^-$
 $M_r = 182.10$

 Orthorhombic, $Pnma$
 $a = 9.379$ (1) Å

 $b = 8.067$ (1) Å

 $c = 11.482$ (2) Å

 $V = 868.7$ (2) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.68$ mm⁻¹
 $T = 100$ (2) K

 $0.15 \times 0.08 \times 0.03$ mm

Data collection

Bruker SMART APEX

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.872$, $T_{\max} = 1.000$

(expected range = 0.855–0.980)

5307 measured reflections

1068 independent reflections

 856 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.113$
 $S = 1.02$

1068 reflections

82 parameters

58 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.57$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{Cl1}$	0.88 (1)	2.13 (1)	3.008 (3)	175 (4)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2008).

We thank the University of Malaya for the purchase of the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2068).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Kurahashi, M., Engel, P. & Nowacki, W. (1980). *Z. Kristallogr.* **152**, 147–156.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2008). *pubCIF*. In preparation.

supporting information

Acta Cryst. (2008). E64, o911 [doi:10.1107/S1600536808011434]

3-Chloroquinuclidinium chloride

Isha Azizul, Arifin Zainudin and Seik Weng Ng

S1. Comment

4-Chloroquinuclidinium chloride features an N–H···Cl hydrogen bond between the cation and anion. The N–C and C–C bonds are somewhat shorter than those in the unsubstituted salt, and this has been attributed to the electron-withdrawing effect of the chlorine substituent (Kurahashi *et al.*, 1980). The present isomeric compound (Scheme I) is expected to show this feature; however, owing to disorder, the effect cannot be unambiguously observed even at low temperature. The cation forms a linear hydrogen bond [N–H···Cl 3.008 (3) Å] to the chloride; the cation is disordered about a mirror plane (Fig. 1).

S2. Experimental

The commercially available compound was a crystalline. A large block was cut into a smaller specimen.

S3. Refinement

The cation is disordered about a mirror plane in the carbon atoms except C1 atom. The N1 and C1 atoms, which lie on this symmetry element, were refined with their normal half occupancies. The other carbon atoms were refined with half occupancies, subject to N–C being restrained to 1.49 ± 0.01 Å and C–C to 1.54 ± 0.01 Å. Additionally, the 1,3-related distances were restrained from 2.43 ± 0.01 Å, to 2.47 ± 0.01 Å as well as 2.52 ± 0.01 Å. The anisotropic temperature factors of the disordered carbon were restrained to be nearly isotropic but the N–H distance was restrained to 0.88 ± 0.01 Å.

Carbon-bound H-atoms were placed in calculated positions (C–H 0.99 to 1.00 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U(\text{C})$. The ammonium H-atom was located in a difference Fourier map, and was refined with an N–H distance restraint of 0.88 ± 0.01 Å; its temperature factor was freely refined.




Figure 1

Thermal ellipsoid plot of the two independent molecules of 2-chloroquinuclidinium chloride at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. The dashed lines denote the hydrogen bond.

3-Chloroquinuclidinium chloride

Crystal data

$C_7H_{13}ClN^+ \cdot Cl^-$

$M_r = 182.10$

Orthorhombic, *Pnma*

Hall symbol: -P 2ac 2n

$a = 9.379 (1) \text{ \AA}$

$b = 8.067 (1) \text{ \AA}$

$c = 11.482 (2) \text{ \AA}$

$V = 868.7 (2) \text{ \AA}^3$

$Z = 4$

$F(000) = 392$

$D_x = 1.408 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 909 reflections

$\theta = 3.1\text{--}22.9^\circ$

$\mu = 0.68 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Block, colorless

$0.15 \times 0.08 \times 0.03 \text{ mm}$

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.872$, $T_{\max} = 1.000$

5307 measured reflections

1068 independent reflections

856 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.8^\circ$

$h = -12 \rightarrow 8$

$k = -10 \rightarrow 10$

$l = -14 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.113$

$S = 1.02$

1068 reflections

82 parameters

58 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0562P)^2 + 0.8408P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.57 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.69230 (8)	0.2500	0.57828 (7)	0.0218 (2)	
C12	0.10483 (11)	0.2500	0.30411 (8)	0.0374 (3)	
N1	0.3718 (3)	0.2500	0.5682 (2)	0.0190 (6)	
H1	0.4648 (12)	0.2500	0.576 (3)	0.035 (12)*	
C1	0.3382 (3)	0.2500	0.4416 (3)	0.0331 (9)	
H1A	0.4092	0.1832	0.3983	0.040*	0.50
H1B	0.3397	0.3645	0.4106	0.040*	0.50
C2	0.1846 (4)	0.1722 (5)	0.4282 (3)	0.0184 (9)	0.50
H2	0.1947	0.0495	0.4194	0.022*	
C3	0.2957 (7)	0.3924 (13)	0.6253 (9)	0.0204 (17)	0.50
H3A	0.3397	0.4988	0.6019	0.024*	0.50
H3B	0.3013	0.3825	0.7111	0.024*	0.50
C4	0.1402 (5)	0.3855 (6)	0.5851 (5)	0.0212 (11)	0.50
H4A	0.1248	0.4653	0.5209	0.025*	0.50
H4B	0.0763	0.4159	0.6503	0.025*	0.50
C5	0.3285 (7)	0.0917 (13)	0.6251 (10)	0.024 (2)	0.50
H5A	0.3698	0.0857	0.7043	0.029*	0.50
H5B	0.3651	-0.0034	0.5795	0.029*	0.50
C6	0.1650 (5)	0.0823 (7)	0.6325 (4)	0.0244 (12)	0.50
H6A	0.1323	0.1111	0.7120	0.029*	0.50
H6B	0.1316	-0.0310	0.6139	0.029*	0.50
C7	0.1058 (4)	0.2079 (5)	0.5430 (4)	0.0205 (12)	0.50
H7	0.0007	0.1931	0.5330	0.025*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0157 (4)	0.0284 (4)	0.0212 (4)	0.000	-0.0006 (3)	0.000
C12	0.0340 (5)	0.0546 (6)	0.0236 (5)	0.000	-0.0111 (4)	0.000
N1	0.0145 (13)	0.0241 (13)	0.0184 (14)	0.000	-0.0020 (10)	0.000
C1	0.0176 (17)	0.064 (3)	0.0182 (19)	0.000	0.0006 (13)	0.000
C2	0.019 (2)	0.0171 (19)	0.019 (2)	0.0018 (17)	-0.0032 (16)	0.0013 (17)
C3	0.019 (3)	0.017 (3)	0.025 (3)	-0.004 (3)	0.008 (3)	0.004 (2)

C4	0.015 (2)	0.022 (3)	0.026 (3)	0.004 (2)	-0.004 (2)	-0.006 (2)
C5	0.017 (3)	0.019 (3)	0.036 (4)	0.000 (3)	0.007 (3)	-0.001 (3)
C6	0.028 (3)	0.026 (3)	0.019 (3)	-0.003 (2)	-0.002 (2)	0.004 (2)
C7	0.0108 (17)	0.030 (4)	0.021 (2)	-0.0037 (16)	0.0014 (15)	-0.0070 (18)

Geometric parameters (Å, °)

C12—C2	1.727 (4)	C3—H3B	0.9900
N1—C5	1.491 (7)	C4—C7	1.546 (6)
N1—C1	1.488 (4)	C4—H4A	0.9900
N1—C3	1.502 (7)	C4—H4B	0.9900
N1—H1	0.88 (1)	C5—C6	1.538 (7)
C1—C2	1.579 (4)	C5—H5A	0.9900
C1—H1A	0.9900	C5—H5B	0.9900
C1—H1B	0.9900	C6—C7	1.546 (5)
C2—C7	1.539 (5)	C6—H6A	0.9900
C2—H2	1.0000	C6—H6B	0.9900
C3—C4	1.531 (7)	C7—H7	1.0000
C3—H3A	0.9900		
C5—N1—C1	111.7 (5)	C3—C4—C7	109.1 (4)
C5—N1—C3	109.5 (3)	C3—C4—H4A	109.9
C1—N1—C3	109.0 (4)	C7—C4—H4A	109.9
C5—N1—H1	103.1 (13)	C3—C4—H4B	109.9
C1—N1—H1	108 (3)	C7—C4—H4B	109.9
C3—N1—H1	115.3 (14)	H4A—C4—H4B	108.3
N1—C1—C2	106.8 (2)	N1—C5—C6	109.8 (5)
N1—C1—H1A	110.4	N1—C5—H5A	109.7
C2—C1—H1A	110.4	C6—C5—H5A	109.7
N1—C1—H1B	110.4	N1—C5—H5B	109.7
C2—C1—H1B	110.4	C6—C5—H5B	109.7
H1A—C1—H1B	108.6	H5A—C5—H5B	108.2
C7—C2—C1	106.3 (3)	C5—C6—C7	106.8 (4)
C7—C2—C12	115.5 (3)	C5—C6—H6A	110.4
C1—C2—C12	109.4 (2)	C7—C6—H6A	110.4
C7—C2—H2	108.5	C5—C6—H6B	110.4
C1—C2—H2	108.5	C7—C6—H6B	110.4
C12—C2—H2	108.5	H6A—C6—H6B	108.6
N1—C3—C4	107.1 (4)	C2—C7—C4	109.9 (3)
N1—C3—H3A	110.3	C2—C7—C6	106.0 (3)
C4—C3—H3A	110.3	C4—C7—C6	108.9 (3)
N1—C3—H3B	110.3	C2—C7—H7	110.6
C4—C3—H3B	110.3	C4—C7—H7	110.6
H3A—C3—H3B	108.6	C6—C7—H7	110.6
C5—N1—C1—C2	42.7 (4)	N1—C5—C6—C7	18.9 (10)
C3—N1—C1—C2	-78.5 (4)	C1—C2—C7—C4	40.7 (4)
N1—C1—C2—C7	27.3 (3)	C12—C2—C7—C4	-80.8 (4)

N1—C1—C2—C12	152.63 (17)	C1—C2—C7—C6	-76.9 (4)
C5—N1—C3—C4	-73.4 (6)	C12—C2—C7—C6	161.6 (3)
C1—N1—C3—C4	49.1 (8)	C3—C4—C7—C2	-70.0 (6)
N1—C3—C4—C7	21.4 (9)	C3—C4—C7—C6	45.7 (6)
C1—N1—C5—C6	-71.2 (9)	C5—C6—C7—C2	49.6 (7)
C3—N1—C5—C6	49.7 (7)	C5—C6—C7—C4	-68.6 (7)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...C11	0.88 (1)	2.13 (1)	3.008 (3)	175 (4)