

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

 Bis[4-(4-methoxyphenyl)-4*H*-1,2,4-triazole- $\kappa$ N<sup>1</sup>]bis(thiocyanato- $\kappa$ N)zinc(II)

Ya Zuo

College of Science, Inner Mongolia Agricultural University, Inner Mongolia 010018, People's Republic of China

Correspondence e-mail: yazuo@mail.nankai.edu.cn


Received 5 March 2008; accepted 27 March 2008

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.009$  Å;  $R$  factor = 0.068;  $wR$  factor = 0.158; data-to-parameter ratio = 16.4.

In the title complex,  $[\text{Zn}(\text{NCS})_2(\text{C}_9\text{H}_9\text{N}_3\text{O})_2]$ , the  $\text{Zn}^{\text{II}}$  ion is coordinated by two N atoms from the  $\text{NCS}^-$  anions and two N atoms from two 4-(4-methoxyphenyl)-4*H*-1,2,4-triazole ligands in a slightly distorted tetrahedral geometry. Three intermolecular weak hydrogen-bonding contacts of the types  $\text{C}-\text{H}\cdots\text{N}$ ,  $\text{C}-\text{H}\cdots\text{S}$  and  $\text{C}-\text{H}\cdots\text{O}$  are observed in the crystal structure.

## Related literature

For related literature, see: Han *et al.* (2006); Ling & Zhang (2007); Thomann *et al.* (1994); Yin *et al.* (2007); Zhao *et al.* (2002); Zhou *et al.* (2007).



## Experimental

## Crystal data

 $[\text{Zn}(\text{NCS})_2(\text{C}_9\text{H}_9\text{N}_3\text{O})_2]$ 
 $M_r = 531.91$ 

 Monoclinic,  $P2_1/n$ 
 $a = 7.812$  (3) Å

 $b = 17.111$  (6) Å

 $c = 18.201$  (6) Å

 $\beta = 99.726$  (6)°

 $V = 2398.0$  (15) Å<sup>3</sup>
 $Z = 4$ 

 Mo  $K\alpha$  radiation

 $\mu = 1.23$  mm<sup>-1</sup>
 $T = 293$  (2) K

 $0.16 \times 0.12 \times 0.10$  mm

## Data collection

Bruker SMART CCD area-detector diffractometer

 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

 $T_{\text{min}} = 0.700$ ,  $T_{\text{max}} = 0.884$ 

13731 measured reflections

4932 independent reflections

 2127 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.121$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.067$ 
 $wR(F^2) = 0.158$ 
 $S = 0.97$ 

4932 reflections

300 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.39$  e Å<sup>-3</sup>
 $\Delta\rho_{\text{min}} = -0.33$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

Zn1—N8	1.923 (6)	Zn1—N1	2.005 (5)
Zn1—N7	1.970 (6)	Zn1—N4	2.009 (5)
N8—Zn1—N7	112.9 (2)	N8—Zn1—N4	116.4 (2)
N8—Zn1—N1	112.1 (2)	N7—Zn1—N4	103.8 (2)
N7—Zn1—N1	108.5 (2)	N1—Zn1—N4	102.2 (2)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1—H1 $\cdots$ N5 <sup>i</sup>	0.93	2.51	3.438 (8)	177
C7—H7 $\cdots$ S1 <sup>ii</sup>	0.93	2.86	3.735 (6)	158
C10—H10 $\cdots$ O1 <sup>iii</sup>	0.93	2.42	3.265 (8)	151

Symmetry codes: (i)  $-x + 1, -y + 2, -z$ ; (ii)  $-x, -y + 2, -z$ ; (iii)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SI2077).

## References

- Bruker (1998). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (1999). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Han, H. Y., Song, Y. L., Hou, H. W., Fan, Y. T. & Zhu, Y. (2006). *Dalton Trans.* pp. 1972–1980.  
 Ling, Y. & Zhang, L. (2007). *Acta Cryst.* **E63**, m4–m6.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Thomann, M., Kahn, O., Guilhem, J. & Varretta, F. (1994). *Inorg. Chem.* **33**, 6029–6037.  
 Yin, G., Zhang, Y. P., Li, B. L. & Zhang, Y. (2007). *J. Mol. Struct.* **837**, 263–268.  
 Zhao, Q. H., Li, H. F., Wang, X. F. & Chen, Z. D. (2002). *New J. Chem.* **26**, 1709–1710.  
 Zhou, J., Yang, J., Qi, L., Shen, X., Zhu, D. R., Xu, Y. & Song, Y. (2007). *Transition Met. Chem.* **32**, 711–715.

## supporting information

*Acta Cryst.* (2008). E64, m609 [doi:10.1107/S1600536808008325]

**Bis[4-(4-methoxyphenyl)-4H-1,2,4-triazole- $\kappa N^1$ ]bis(thiocyanato- $\kappa N$ )zinc(II)**

Ya Zuo

**S1. Comment**

The organic ligands, 1,2,4-triazole and its derivatives continue to attract considerable attention due to the luminescent, magnetic, and electronic properties of their complexes and the potential application in material science (Yin *et al.*, 2007; Zhou, *et al.*, 2007; Han *et al.*, 2006). On the other hand, 1,2,4-triazole and its derivatives combine the coordination geometry of both pyrazole and imidazole with regard to the arrangement and coordination of their three heteroatoms. Compared with the chelating to the metal centers with the nitrogen atom in the 1,2-positions (Thomann *et al.*, 1994), it is familiar that the nitrogen atom serve as the solo coordinated atom (Ling & Zhang, 2007). In order to explore further the coordination chemistry of the 4-(4-methoxyphenyl)-4H-1,2,4-triazole system (hereafter abbreviated as *L*), the title complex was synthesized.

The coordination geometry of the Zn<sup>II</sup> ion is a slightly distorted tetrahedron, in which each Zn<sup>II</sup> ion is coordinated to two nitrogen atoms from the NCS<sup>-</sup> anions and two nitrogen atoms from two ligands *L*. (Fig. 1). The separations of Zn—N range from 1.923 (6) to 2.009 (5) Å (Table 1), which is close to the corresponding distances in the 4-coordinated Zn compound (2.050 (4) Å and 2.023 (4) Å) reported previously (Han *et al.*, 2006), while the average distance of Zn—N is 2.162 (2) Å in the 6-coordinated (Ling & Zhang, 2007) and 2.065 (2) Å in the 5-coordinated (Yin *et al.*, 2007) compounds, respectively. The bond angles around Zn<sup>II</sup> ions vary between 102.2 (2)° and 116.4 (2)°. The central Zn<sup>II</sup> ion, is coordinated with the *L* ligands in the 1-position, which is the common coordination mode (Ling & Zhang, 2007) of 1,2,4-Triazole and its derivatives, differently from that in the 1,2-position (Thomann *et al.*, 1994). As the important structural parameters, the dihedral angles between the triazole and phenyl rings of the two *L* ligands are distinct, that of C1—N3—C3—C8 is equal to 39.2 (9)°, while the value of C10—N6—C12—C17 is 93.8 (8)°. With regard to the two *trans* thiocyanato NCS<sup>-</sup> anions, the group is almost linear with an N7—C19—S1 angle of 178.8 (8)° and N8—C20—S2 angle of 178.1 (7)°. The connection between Zn atoms and NCS groups are bent with a C20—N8—Zn1 angle of 168.1 (6)°, and a C19—N7—Zn1 angle of 160.9 (6)° (Zhao *et al.*, 2002). Three intermolecular hydrogen bonding contacts of the type C—H $\cdots$ N, C—H $\cdots$ S and C—H $\cdots$ O are observed in the title structure (Table 2).


**S2. Experimental**

The compound was synthesized under hydrothermal conditions. A mixture of *L* (*L*=4-(1,2,4-triazol)-1-methoxy-benzene) (0.3 mmol, 0.045 g), ZnSO<sub>4</sub>·7H<sub>2</sub>O (0.1 mmol, 0.029 g), KSCN (0.2 mmol, 0.019 g) and water (10 mL) was placed in a 25 mL acid digestion bomb and heated at 160° for two days, then equably cooled to room temperature for three days. After washed by 5 ml water for twice, colorless block crystals of the title compound were obtained.

**S3. Refinement**

The H atoms (methyl) on the ligands were allowed to ride on their parent atoms with C—H distances of 0.96 Å and  $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}(\text{C})$ , and the rest C—H distances of 0.93 Å and  $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}(\text{C})$ . All of the non-hydrogen atoms were

refined anisotropically.



**Figure 1**

The *ORTEP* plot of the title complex, showing 30% probability displacement ellipsoids and the atom-numbering schemes. H atoms have been omitted for clarity.

**Bis[4-(4-methoxyphenyl)-4H-1,2,4-triazole- $\kappa$ N<sup>1</sup>]bis(thiocyanato- $\kappa$ N)zinc(II)**

*Crystal data*

[Zn(NCS)<sub>2</sub>(C<sub>9</sub>H<sub>9</sub>N<sub>3</sub>O)<sub>2</sub>]

$M_r = 531.91$

Monoclinic,  $P2_1/n$

$a = 7.812$  (3) Å

$b = 17.111$  (6) Å

$c = 18.201$  (6) Å

$\beta = 99.726$  (6)°

$V = 2398.0$  (15) Å<sup>3</sup>

$Z = 4$

$F(000) = 1088$

$D_x = 1.473$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 713 reflections

$\theta = 2.6$ – $21.1$ °

$\mu = 1.23$  mm<sup>-1</sup>

$T = 293$  K

Block, colorless

$0.16 \times 0.12 \times 0.10$  mm

*Data collection*

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.700$ ,  $T_{\max} = 0.884$

13731 measured reflections

4932 independent reflections

2127 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.121$

$\theta_{\max} = 26.5$ °,  $\theta_{\min} = 1.6$ °

$h = -9 \rightarrow 9$

$k = -21 \rightarrow 14$

$l = -22 \rightarrow 18$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.067$   
 $wR(F^2) = 0.158$   
 $S = 0.97$   
 4932 reflections  
 300 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0555P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.39 \text{ e } \text{Å}^{-3}$   
 $\Delta\rho_{\min} = -0.33 \text{ e } \text{Å}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{Å}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.36832 (9)	1.10583 (5)	0.05909 (4)	0.0434 (3)
N1	0.2576 (6)	1.0131 (3)	0.0996 (3)	0.0448 (14)
N2	0.2447 (7)	1.0191 (4)	0.1742 (3)	0.0506 (15)
N3	0.1960 (6)	0.8971 (4)	0.1360 (3)	0.0417 (13)
O1	0.0331 (6)	0.5821 (3)	0.1525 (2)	0.0495 (12)
C1	0.2292 (7)	0.9405 (4)	0.0785 (3)	0.0393 (16)
H1	0.2317	0.9216	0.0307	0.047*
C2	0.2083 (9)	0.9483 (5)	0.1939 (4)	0.055 (2)
H2	0.1927	0.9346	0.2418	0.066*
C3	0.1559 (8)	0.8147 (4)	0.1374 (3)	0.0364 (15)
C4	0.2212 (8)	0.7714 (4)	0.1995 (3)	0.0436 (17)
H4	0.2949	0.7943	0.2391	0.052*
C5	0.1762 (8)	0.6938 (4)	0.2023 (3)	0.0429 (17)
H5	0.2179	0.6646	0.2447	0.051*
C6	0.0701 (8)	0.6587 (4)	0.1433 (3)	0.0383 (16)
C7	0.0088 (8)	0.7023 (4)	0.0807 (3)	0.0462 (18)
H7	-0.0610	0.6791	0.0400	0.055*
C8	0.0506 (8)	0.7795 (4)	0.0784 (3)	0.0467 (18)
H8	0.0073	0.8089	0.0364	0.056*
C9	-0.0628 (8)	0.5413 (4)	0.0903 (4)	0.060 (2)
H9A	-0.1752	0.5649	0.0767	0.090*
H9B	-0.0762	0.4876	0.1035	0.090*
H9C	-0.0014	0.5440	0.0489	0.090*
N4	0.6037 (6)	1.1085 (3)	0.1237 (3)	0.0423 (13)
N5	0.7566 (7)	1.1233 (4)	0.0996 (3)	0.0586 (18)

N6	0.8049 (6)	1.1152 (3)	0.2222 (3)	0.0421 (14)
O2	1.1582 (6)	1.1055 (3)	0.5135 (2)	0.0655 (14)
C10	0.6367 (8)	1.1026 (4)	0.1966 (3)	0.0464 (16)
H10	0.5541	1.0912	0.2264	0.056*
C11	0.8730 (9)	1.1266 (5)	0.1594 (4)	0.065 (2)
H11	0.9902	1.1359	0.1592	0.078*
C12	0.8924 (7)	1.1139 (4)	0.2991 (3)	0.0385 (15)
C13	0.8962 (8)	1.1802 (4)	0.3406 (3)	0.0474 (18)
H13	0.8410	1.2253	0.3202	0.057*
C14	0.9835 (8)	1.1797 (4)	0.4138 (4)	0.0509 (18)
H14	0.9860	1.2246	0.4429	0.061*
C15	1.0660 (7)	1.1130 (4)	0.4433 (3)	0.0436 (16)
C16	1.0579 (8)	1.0461 (4)	0.4010 (4)	0.0505 (18)
H16	1.1114	1.0007	0.4214	0.061*
C17	0.9708 (8)	1.0461 (4)	0.3285 (4)	0.0462 (18)
H17	0.9651	1.0009	0.2999	0.055*
C18	1.1740 (10)	1.1722 (5)	0.5603 (4)	0.080 (3)
H18A	1.2279	1.2138	0.5372	0.120*
H18B	1.2440	1.1595	0.6074	0.120*
H18C	1.0608	1.1884	0.5681	0.120*
S1	0.1894 (3)	1.35631 (13)	0.11268 (10)	0.0765 (7)
N7	0.2543 (7)	1.2013 (4)	0.0874 (3)	0.0527 (16)
C19	0.2283 (9)	1.2663 (5)	0.0986 (3)	0.0445 (18)
S2	0.3020 (3)	1.10216 (14)	-0.19983 (10)	0.0796 (7)
N8	0.3645 (7)	1.0978 (4)	-0.0466 (3)	0.0619 (17)
C20	0.3357 (8)	1.1003 (4)	-0.1112 (4)	0.0500 (17)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.0453 (4)	0.0419 (5)	0.0414 (4)	0.0001 (5)	0.0025 (3)	0.0030 (4)
N1	0.045 (3)	0.045 (4)	0.045 (3)	-0.001 (3)	0.007 (3)	-0.003 (3)
N2	0.070 (4)	0.047 (4)	0.036 (3)	-0.003 (3)	0.012 (3)	-0.007 (3)
N3	0.051 (3)	0.045 (4)	0.029 (3)	-0.001 (3)	0.008 (2)	-0.001 (3)
O1	0.062 (3)	0.040 (3)	0.046 (3)	-0.008 (2)	0.006 (2)	0.001 (2)
C1	0.041 (4)	0.043 (5)	0.033 (4)	0.000 (3)	0.006 (3)	-0.009 (3)
C2	0.073 (5)	0.055 (6)	0.039 (4)	-0.002 (4)	0.013 (4)	-0.005 (4)
C3	0.041 (4)	0.032 (4)	0.037 (4)	0.005 (3)	0.010 (3)	0.000 (3)
C4	0.047 (4)	0.051 (5)	0.031 (4)	-0.007 (4)	0.000 (3)	-0.009 (3)
C5	0.044 (4)	0.048 (5)	0.036 (4)	0.000 (3)	0.004 (3)	0.011 (3)
C6	0.033 (3)	0.043 (5)	0.041 (4)	-0.002 (3)	0.011 (3)	-0.002 (3)
C7	0.052 (4)	0.047 (5)	0.035 (4)	-0.014 (4)	-0.007 (3)	-0.007 (3)
C8	0.059 (4)	0.043 (5)	0.033 (4)	-0.009 (4)	-0.004 (3)	0.012 (3)
C9	0.061 (5)	0.050 (5)	0.068 (5)	-0.010 (4)	0.006 (4)	-0.019 (4)
N4	0.037 (3)	0.052 (4)	0.039 (3)	0.003 (3)	0.007 (2)	0.008 (3)
N5	0.046 (3)	0.087 (5)	0.043 (3)	-0.005 (3)	0.009 (3)	0.009 (3)
N6	0.037 (3)	0.049 (4)	0.039 (3)	-0.003 (3)	0.006 (3)	0.000 (3)
O2	0.071 (3)	0.070 (4)	0.048 (3)	0.003 (3)	-0.012 (2)	-0.011 (3)

C10	0.042 (4)	0.052 (5)	0.045 (4)	-0.002 (4)	0.007 (3)	0.000 (4)
C11	0.045 (4)	0.099 (7)	0.052 (5)	-0.012 (4)	0.012 (4)	0.006 (5)
C12	0.033 (3)	0.046 (5)	0.039 (4)	0.001 (3)	0.009 (3)	-0.003 (4)
C13	0.048 (4)	0.048 (5)	0.046 (4)	0.008 (4)	0.008 (4)	0.008 (4)
C14	0.058 (5)	0.041 (5)	0.053 (4)	0.003 (4)	0.006 (4)	-0.009 (4)
C15	0.039 (3)	0.049 (5)	0.043 (4)	-0.009 (4)	0.006 (3)	-0.006 (4)
C16	0.057 (4)	0.040 (5)	0.050 (4)	0.013 (4)	-0.004 (4)	0.006 (4)
C17	0.045 (4)	0.040 (5)	0.053 (5)	0.009 (4)	0.008 (3)	-0.011 (4)
C18	0.072 (5)	0.100 (8)	0.059 (5)	0.011 (5)	-0.010 (4)	-0.018 (5)
S1	0.128 (2)	0.0453 (13)	0.0492 (12)	0.0171 (13)	-0.0041 (12)	-0.0014 (10)
N7	0.060 (4)	0.050 (4)	0.045 (3)	0.003 (3)	-0.001 (3)	-0.001 (3)
C19	0.053 (4)	0.048 (5)	0.030 (4)	-0.005 (4)	0.000 (3)	0.005 (4)
S2	0.1267 (18)	0.0637 (15)	0.0433 (11)	-0.0087 (14)	-0.0001 (12)	-0.0048 (11)
N8	0.065 (4)	0.070 (5)	0.048 (4)	0.004 (4)	0.003 (3)	0.001 (4)
C20	0.054 (4)	0.038 (4)	0.055 (5)	0.001 (4)	0.001 (4)	-0.001 (4)

*Geometric parameters (Å, °)*

Zn1—N8	1.923 (6)	N4—C10	1.312 (7)
Zn1—N7	1.970 (6)	N4—N5	1.365 (6)
Zn1—N1	2.005 (5)	N5—C11	1.296 (7)
Zn1—N4	2.009 (5)	N6—C10	1.336 (7)
N1—C1	1.309 (8)	N6—C11	1.354 (7)
N1—N2	1.382 (7)	N6—C12	1.450 (7)
N2—C2	1.308 (8)	O2—C15	1.363 (7)
N3—C1	1.344 (7)	O2—C18	1.417 (8)
N3—C2	1.361 (8)	C10—H10	0.9300
N3—C3	1.445 (8)	C11—H11	0.9300
O1—C6	1.359 (7)	C12—C13	1.360 (9)
O1—C9	1.430 (7)	C12—C17	1.377 (8)
C1—H1	0.9300	C13—C14	1.391 (8)
C2—H2	0.9300	C13—H13	0.9300
C3—C8	1.375 (8)	C14—C15	1.374 (9)
C3—C4	1.376 (8)	C14—H14	0.9300
C4—C5	1.378 (9)	C15—C16	1.375 (9)
C4—H4	0.9300	C16—C17	1.379 (8)
C5—C6	1.379 (8)	C16—H16	0.9300
C5—H5	0.9300	C17—H17	0.9300
C6—C7	1.379 (8)	C18—H18A	0.9600
C7—C8	1.364 (9)	C18—H18B	0.9600
C7—H7	0.9300	C18—H18C	0.9600
C8—H8	0.9300	S1—C19	1.598 (8)
C9—H9A	0.9600	N7—C19	1.155 (8)
C9—H9B	0.9600	S2—C20	1.591 (7)
C9—H9C	0.9600	N8—C20	1.160 (7)
N8—Zn1—N7	112.9 (2)	H9B—C9—H9C	109.5
N8—Zn1—N1	112.1 (2)	C10—N4—N5	107.9 (5)

N7—Zn1—N1	108.5 (2)	C10—N4—Zn1	126.4 (4)
N8—Zn1—N4	116.4 (2)	N5—N4—Zn1	125.5 (4)
N7—Zn1—N4	103.8 (2)	C11—N5—N4	105.4 (5)
N1—Zn1—N4	102.2 (2)	C10—N6—C11	103.4 (5)
C1—N1—N2	108.8 (5)	C10—N6—C12	127.6 (5)
C1—N1—Zn1	134.9 (5)	C11—N6—C12	129.0 (5)
N2—N1—Zn1	114.2 (4)	C15—O2—C18	117.8 (6)
C2—N2—N1	104.9 (5)	N4—C10—N6	110.7 (6)
C1—N3—C2	104.5 (6)	N4—C10—H10	124.6
C1—N3—C3	128.5 (5)	N6—C10—H10	124.6
C2—N3—C3	127.0 (6)	N5—C11—N6	112.6 (6)
C6—O1—C9	117.9 (5)	N5—C11—H11	123.7
N1—C1—N3	109.9 (6)	N6—C11—H11	123.7
N1—C1—H1	125.0	C13—C12—C17	121.3 (6)
N3—C1—H1	125.0	C13—C12—N6	119.1 (6)
N2—C2—N3	111.9 (6)	C17—C12—N6	119.7 (6)
N2—C2—H2	124.0	C12—C13—C14	119.3 (6)
N3—C2—H2	124.0	C12—C13—H13	120.4
C8—C3—C4	119.7 (6)	C14—C13—H13	120.4
C8—C3—N3	121.1 (6)	C15—C14—C13	120.0 (6)
C4—C3—N3	119.2 (6)	C15—C14—H14	120.0
C3—C4—C5	119.2 (6)	C13—C14—H14	120.0
C3—C4—H4	120.4	O2—C15—C14	125.7 (7)
C5—C4—H4	120.4	O2—C15—C16	114.3 (7)
C4—C5—C6	121.0 (6)	C14—C15—C16	120.0 (6)
C4—C5—H5	119.5	C15—C16—C17	120.2 (6)
C6—C5—H5	119.5	C15—C16—H16	119.9
O1—C6—C7	125.0 (6)	C17—C16—H16	119.9
O1—C6—C5	115.9 (6)	C12—C17—C16	119.2 (6)
C7—C6—C5	119.2 (6)	C12—C17—H17	120.4
C8—C7—C6	119.9 (6)	C16—C17—H17	120.4
C8—C7—H7	120.1	O2—C18—H18A	109.5
C6—C7—H7	120.1	O2—C18—H18B	109.5
C7—C8—C3	121.0 (6)	H18A—C18—H18B	109.5
C7—C8—H8	119.5	O2—C18—H18C	109.5
C3—C8—H8	119.5	H18A—C18—H18C	109.5
O1—C9—H9A	109.5	H18B—C18—H18C	109.5
O1—C9—H9B	109.5	C19—N7—Zn1	161.0 (6)
H9A—C9—H9B	109.5	N7—C19—S1	178.8 (7)
O1—C9—H9C	109.5	C20—N8—Zn1	168.1 (6)
H9A—C9—H9C	109.5	N8—C20—S2	178.1 (7)
N8—Zn1—N1—C1	24.0 (7)	N7—Zn1—N4—N5	-108.5 (5)
N7—Zn1—N1—C1	149.4 (6)	N1—Zn1—N4—N5	138.7 (5)
N4—Zn1—N1—C1	-101.4 (6)	C10—N4—N5—C11	-0.9 (8)
N8—Zn1—N1—N2	-175.1 (4)	Zn1—N4—N5—C11	174.2 (5)
N7—Zn1—N1—N2	-49.7 (4)	N5—N4—C10—N6	1.8 (8)
N4—Zn1—N1—N2	59.5 (4)	Zn1—N4—C10—N6	-173.2 (4)

C1—N1—N2—C2	-0.1 (7)	C11—N6—C10—N4	-1.9 (8)
Zn1—N1—N2—C2	-165.9 (4)	C12—N6—C10—N4	-180.0 (6)
N2—N1—C1—N3	0.5 (7)	N4—N5—C11—N6	-0.3 (9)
Zn1—N1—C1—N3	162.2 (4)	C10—N6—C11—N5	1.4 (9)
C2—N3—C1—N1	-0.7 (7)	C12—N6—C11—N5	179.4 (6)
C3—N3—C1—N1	179.2 (5)	C10—N6—C12—C13	-86.6 (8)
N1—N2—C2—N3	-0.4 (7)	C11—N6—C12—C13	95.8 (8)
C1—N3—C2—N2	0.7 (7)	C10—N6—C12—C17	93.7 (8)
C3—N3—C2—N2	-179.3 (6)	C11—N6—C12—C17	-83.8 (9)
C1—N3—C3—C8	-39.2 (9)	C17—C12—C13—C14	1.1 (10)
C2—N3—C3—C8	140.7 (6)	N6—C12—C13—C14	-178.5 (5)
C1—N3—C3—C4	142.3 (6)	C12—C13—C14—C15	0.6 (10)
C2—N3—C3—C4	-37.7 (9)	C18—O2—C15—C14	-1.2 (9)
C8—C3—C4—C5	-1.6 (9)	C18—O2—C15—C16	179.4 (6)
N3—C3—C4—C5	176.9 (5)	C13—C14—C15—O2	178.7 (6)
C3—C4—C5—C6	1.4 (9)	C13—C14—C15—C16	-1.9 (10)
C9—O1—C6—C7	6.3 (9)	O2—C15—C16—C17	-179.0 (6)
C9—O1—C6—C5	-174.3 (5)	C14—C15—C16—C17	1.6 (10)
C4—C5—C6—O1	-179.4 (5)	C13—C12—C17—C16	-1.4 (10)
C4—C5—C6—C7	0.1 (9)	N6—C12—C17—C16	178.2 (5)
O1—C6—C7—C8	178.2 (6)	C15—C16—C17—C12	0.1 (10)
C5—C6—C7—C8	-1.3 (9)	N8—Zn1—N7—C19	-71.6 (17)
C6—C7—C8—C3	1.1 (10)	N1—Zn1—N7—C19	163.5 (16)
C4—C3—C8—C7	0.4 (10)	N4—Zn1—N7—C19	55.3 (17)
N3—C3—C8—C7	-178.0 (6)	Zn1—N7—C19—S1	105 (36)
N8—Zn1—N4—C10	-169.6 (6)	N7—Zn1—N8—C20	-29 (3)
N7—Zn1—N4—C10	65.7 (6)	N1—Zn1—N8—C20	93 (3)
N1—Zn1—N4—C10	-47.1 (6)	N4—Zn1—N8—C20	-149 (3)
N8—Zn1—N4—N5	16.2 (6)	Zn1—N8—C20—S2	180 (100)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...N5 <sup>i</sup>	0.93	2.51	3.438 (8)	177
C7—H7...S1 <sup>ii</sup>	0.93	2.86	3.735 (6)	158
C10—H10...O1 <sup>iii</sup>	0.93	2.42	3.265 (8)	151

Symmetry codes: (i)  $-x+1, -y+2, -z$ ; (ii)  $-x, -y+2, -z$ ; (iii)  $-x+1/2, y+1/2, -z+1/2$ .