

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[cyclohexyldiphenyltin(IV)]- μ -hydroxido- κ^2 O:O]

Kong Mun Lo and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 26 March 2008; accepted 20 April 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å;
 R factor = 0.018; wR factor = 0.072; data-to-parameter ratio = 17.6.

The title polymeric mixed-organyl tin hydroxide, $[\text{Sn}(\text{C}_6\text{H}_5)_2(\text{C}_6\text{H}_{11})(\text{OH})]_n$, has a hydroxide-bridged chain structure; the tin center shows *trans*- C_3SnO_2 trigonal bipyramidal coordination. The Sn atom lies on a special position of site symmetry m ; the symmetry element relates one phenyl ring to the other and also relates one half of the cyclohexyl ring to the other half.

Related literature

For background literature on mixed alkyl/diaryltin(IV) compounds, see: Koshy *et al.* (2001). For the synthesis of cyclohexyldiphenyltin hydroxide, see: Teo *et al.* (2007). For the structure of triethyltin hydroxide, see: Deacon *et al.* (1993). For the structure of tribenzyltin hydroxide, see: Chen *et al.* (2005); Reuter (2004). For the structure of triphenyltin hydroxide, see: Fu *et al.* (2003); Glidewell & Liles (1978); Glidewell *et al.* (2002). For the structure of the mixed organyl compound, benzyldimethyltin hydroxide, see: Wannagat *et al.* (1993).



Experimental

Crystal data

$[\text{Sn}(\text{C}_6\text{H}_5)_2(\text{C}_6\text{H}_{11})(\text{OH})]$
 $M_r = 373.05$
Orthorhombic, $Cmc2_1$
 $a = 18.3830$ (2) Å
 $b = 10.2801$ (1) Å
 $c = 8.1762$ (1) Å
 $V = 1545.13$ (3) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.65$ mm⁻¹
 $T = 100$ (2) K
 $0.22 \times 0.09 \times 0.08$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.771$, $T_{\max} = 0.880$
9651 measured reflections
1711 independent reflections
1637 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.071$
 $S = 1.28$
1711 reflections
97 parameters
1 restraint
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.57$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³
Absolute structure: Flack (1983),
650 Friedel pairs
Flack parameter: 0.02 (4)

Table 1

Selected geometric parameters (Å, °).

Sn1—O1	2.201 (4)	Sn1—C5	2.139 (3)
Sn1—C1	2.159 (4)		
C1—Sn1—C5	118.4 (1)	C5—Sn1—O1	90.7 (1)
C1—Sn1—O1	94.1 (2)	C5—Sn1—O1 ⁱ	87.5 (1)
C1—Sn1—O1 ⁱ	89.8 (2)	O1—Sn1—O1 ⁱ	176.1 (1)
C5—Sn1—C5 ⁱⁱ	122.9 (2)	Sn1—O1—Sn1 ⁱⁱⁱ	133.7 (2)

Symmetry codes: (i) $-x + 1, -y + 1, z + \frac{1}{2}$; (ii) $-x + 1, y, z$; (iii) $-x + 1, -y + 1, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

We thank the University of Malaya for funding this study (SF022155/2007 A) and also for the purchase of the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2257).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Chen, Z.-M., Wang, J.-Q., Kuang, D.-Z., Feng, Y.-L. & Zhang, F.-X. (2005). *Chin. J. Inorg. Chem.* **21**, 1186–1190.
Deacon, G. B., Lawrenz, E., Nelson, K. T. & Tiekink, E. R. T. (1993). *Main Group Met. Chem.* **16**, 265–269.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Fu, C.-X., Zhang, J.-H., Ma, C.-L. & Zhang, Z.-T. (2003). *Chin. J. Synth. Chem.* **11**, 189–193.
Glidewell, C. & Liles, D. C. (1978). *Acta Cryst.* **B34**, 129–134.
Glidewell, C., Low, J. N., Bomfim, J. A. S., Filgueiras, C. A. L. & Wardell, J. L. (2002). *Acta Cryst.* **C58**, m199–m201.

- Koshy, J., Ansary, A., Lo, K. M. & Kumar Das, V. G. (2001). *Met.-Based Drugs*, **8**, 107–111.
- Reuter, H. (2004). *Z. Kristallogr. New Cryst. Struct.* **219**, 487–488.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Teo, Y. Y., Lo, K. M. & Ng, S. W. (2007). *Acta Cryst.* **E63**, m1365–m1367.
- Wannagat, U., Dmarath, V., Huch, V., Veith, M. & Harder, U. (1993). *J. Organomet. Chem.* **443**, 153–165.
- Westrip, S. P. (2008). *publCIF*. In preparation.

supporting information

Acta Cryst. (2008). E64, m724–m725 [doi:10.1107/S1600536808011100]

catena-Poly[[cyclohexyldiphenyltin(IV)]- μ -hydroxido- κ^2 O:O]

Kong Mun Lo and Seik Weng Ng

S1. Comment

Mixed alkyl/diaryltin compounds possess much more useful activity against plant pathogens than the symmetrical triorganotin homologs, particularly if one of the alkyl substituent is a cyclic unit (Koshy *et al.*, 2001). The title compound (I) is the starting reactant for the synthesis of mixed organotin carboxylates.

The compound adopts a zigzag chain motif that propagates along the *c*-axis of the orthorhombic unit cell; the tin center shows *trans*-C₃SnO₂ trigonal bipyramidal coordination (Figs 1 and 2 & Table 1).

S2. Experimental

The compound was synthesized as described previously (Teo *et al.*, 2007). Crystals were obtained by recrystallization from ethanol.

S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to $1.2U_{\text{eq}}(\text{C})$. The hydroxo H atom (O—H 0.84 Å) was similarly treated.




Figure 1

70% Probability thermal ellipsoid plot of the asymmetric unit in $\text{Sn}(\text{C}_6\text{H}_{11})(\text{C}_6\text{H}_5)_2(\text{OH})$ (I) extended to show the *trans*- C_3SnO_2 trigonal bipyramidal coordination geometry. Hydrogen atoms are drawn as spheres of arbitrary radius; symmetry-related atoms are not labeled.




Figure 2

Hydroxo-bridged chain motif in (I).

catena-Poly[[cyclohexyldiphenyltin(IV)]- μ -hydroxido- κ^2 O:O]

Crystal data

[Sn(C₆H₅)₂(C₆H₁₁)(OH)]

$M_r = 373.05$

Orthorhombic, *Cmc*2₁

Hall symbol: C 2c -2

$a = 18.3830$ (2) Å

$b = 10.2801$ (1) Å

$c = 8.1762$ (1) Å

$V = 1545.13$ (3) Å³

$Z = 4$

$F(000) = 752$

$D_x = 1.604$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8850 reflections

$\theta = 2.2$ – 28.3°

$\mu = 1.65$ mm⁻¹

$T = 100$ K

Prism, colorless

$0.22 \times 0.09 \times 0.08$ mm

Data collection

Bruker SMART APEXII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.771$, $T_{\max} = 0.880$

9651 measured reflections

1711 independent reflections

1637 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.2^\circ$

$h = -23 \rightarrow 23$
 $k = -13 \rightarrow 13$
 $l = -9 \rightarrow 10$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.071$
 $S = 1.28$
 1711 reflections
 97 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0421P)^2 + 0.0692P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.57 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.31 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 650 Friedel
 pairs
 Absolute structure parameter: 0.02 (4)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.5000	0.494121 (18)	0.50000 (18)	0.01298 (10)
O1	0.5000	0.5850 (3)	0.2563 (4)	0.0165 (6)
H1O	0.5000	0.6666	0.2598	0.025*
C1	0.5000	0.2980 (4)	0.4053 (6)	0.0189 (9)
H1	0.5000	0.3071	0.2836	0.023*
C2	0.43184 (16)	0.2226 (3)	0.4465 (5)	0.0234 (7)
H2A	0.3888	0.2721	0.4084	0.028*
H2B	0.4281	0.2132	0.5667	0.028*
C3	0.4313 (2)	0.0883 (3)	0.3681 (5)	0.0260 (8)
H3A	0.3885	0.0392	0.4078	0.031*
H3B	0.4268	0.0977	0.2480	0.031*
C4	0.5000	0.0125 (4)	0.4074 (8)	0.0248 (13)
H4A	0.5000	-0.0102	0.5251	0.030*
H4B	0.5000	-0.0696	0.3442	0.030*
C5	0.60222 (16)	0.5831 (3)	0.5558 (4)	0.0170 (6)
C6	0.62261 (15)	0.7060 (2)	0.5004 (5)	0.0227 (6)
H6	0.5897	0.7548	0.4352	0.027*
C7	0.69028 (19)	0.7587 (3)	0.5387 (4)	0.0296 (8)
H7	0.7023	0.8441	0.5036	0.036*
C8	0.74004 (18)	0.6867 (4)	0.6280 (5)	0.0308 (8)
H8	0.7867	0.7216	0.6514	0.037*
C9	0.72163 (17)	0.5644 (4)	0.6826 (5)	0.0256 (7)
H9	0.7553	0.5149	0.7449	0.031*
C10	0.6529 (3)	0.5133 (3)	0.6458 (7)	0.0247 (9)
H10	0.6407	0.4287	0.6836	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.01281 (14)	0.01401 (14)	0.01211 (16)	0.000	0.000	0.0003 (2)
O1	0.0227 (14)	0.0168 (15)	0.0101 (15)	0.000	0.000	-0.0008 (11)
C1	0.022 (2)	0.0167 (19)	0.018 (2)	0.000	0.000	0.0005 (17)
C2	0.0200 (15)	0.0215 (14)	0.0288 (19)	-0.0023 (11)	0.0003 (13)	-0.0006 (12)
C3	0.0306 (19)	0.0193 (15)	0.028 (2)	-0.0044 (13)	0.0033 (14)	0.0001 (14)
C4	0.038 (4)	0.019 (2)	0.017 (3)	0.000	0.000	-0.0032 (17)
C5	0.0168 (13)	0.0194 (13)	0.0147 (15)	-0.0006 (12)	0.0029 (11)	-0.0018 (11)
C6	0.0268 (14)	0.0242 (12)	0.0171 (16)	-0.0042 (10)	-0.0025 (17)	0.001 (2)
C7	0.0366 (18)	0.0302 (16)	0.022 (2)	-0.0150 (14)	0.0004 (14)	0.0036 (13)
C8	0.0213 (16)	0.043 (2)	0.028 (2)	-0.0107 (14)	-0.0009 (15)	-0.0088 (16)
C9	0.0174 (15)	0.0343 (19)	0.0250 (18)	0.0007 (13)	-0.0047 (13)	-0.0072 (15)
C10	0.023 (2)	0.0195 (17)	0.032 (3)	-0.0007 (11)	-0.0051 (18)	-0.0014 (14)

Geometric parameters (\AA , $^\circ$)

Sn1—O1	2.201 (4)	C8—C9	1.377 (5)
Sn1—C1	2.159 (4)	C9—C10	1.400 (5)
Sn1—C5	2.139 (3)	O1—H1O	0.8400
Sn1—C5 ⁱ	2.139 (3)	C1—H1	1.0000
Sn1—O1 ⁱⁱ	2.248 (4)	C2—H2A	0.9900
O1—Sn1 ⁱⁱⁱ	2.248 (4)	C2—H2B	0.9900
C1—C2	1.511 (4)	C3—H3A	0.9900
C1—C2 ⁱ	1.511 (4)	C3—H3B	0.9900
C2—C3	1.522 (4)	C4—H4A	0.9900
C3—C4	1.518 (4)	C4—H4B	0.9900
C4—C3 ⁱ	1.518 (4)	C6—H6	0.9500
C5—C10	1.388 (6)	C7—H7	0.9500
C5—C6	1.394 (4)	C8—H8	0.9500
C6—C7	1.393 (4)	C9—H9	0.9500
C7—C8	1.385 (5)	C10—H10	0.9500
C1—Sn1—C5	118.4 (1)	C2—C1—H1	105.6
C1—Sn1—O1	94.1 (2)	Sn1—C1—H1	105.6
C1—Sn1—O1 ⁱⁱ	89.8 (2)	C1—C2—H2A	109.2
C5—Sn1—C5 ⁱ	122.9 (2)	C3—C2—H2A	109.2
C5—Sn1—O1	90.7 (1)	C1—C2—H2B	109.2
C5—Sn1—O1 ⁱⁱ	87.5 (1)	C3—C2—H2B	109.2
C5 ⁱ —Sn1—C1	118.4 (1)	H2A—C2—H2B	107.9
C5 ⁱ —Sn1—O1	90.7 (1)	C4—C3—H3A	109.3
C5 ⁱ —Sn1—O1 ⁱⁱ	87.5 (1)	C2—C3—H3A	109.3
O1—Sn1—O1 ⁱⁱ	176.1 (1)	C4—C3—H3B	109.3
Sn1—O1—Sn1 ⁱⁱⁱ	133.7 (2)	C2—C3—H3B	109.3
C2—C1—C2 ⁱ	112.0 (3)	H3A—C3—H3B	107.9
C2—C1—Sn1	113.5 (2)	C3 ⁱ —C4—H4A	109.1
C2 ⁱ —C1—Sn1	113.5 (2)	C3—C4—H4A	109.1

C1—C2—C3	112.1 (3)	C3 ⁱ —C4—H4B	109.1
C4—C3—C2	111.8 (3)	C3—C4—H4B	109.1
C3 ⁱ —C4—C3	112.5 (4)	H4A—C4—H4B	107.8
C10—C5—C6	117.4 (3)	C7—C6—H6	119.3
C10—C5—Sn1	118.8 (2)	C5—C6—H6	119.3
C6—C5—Sn1	123.7 (2)	C8—C7—H7	120.0
C7—C6—C5	121.3 (3)	C6—C7—H7	120.0
C8—C7—C6	120.0 (3)	C9—C8—H8	120.1
C9—C8—C7	119.8 (3)	C7—C8—H8	120.1
C8—C9—C10	119.6 (4)	C8—C9—H9	120.2
C5—C10—C9	121.7 (3)	C10—C9—H9	120.2
Sn1—O1—H10	113.2	C5—C10—H10	119.1
Sn1 ⁱⁱⁱ —O1—H10	113.2	C9—C10—H10	119.1
C5—Sn1—O1—Sn1 ⁱⁱⁱ	118.53 (8)	C1—Sn1—C5—C10	-47.4 (4)
C5 ⁱ —Sn1—O1—Sn1 ⁱⁱⁱ	-118.53 (8)	O1—Sn1—C5—C10	-142.4 (3)
C1—Sn1—O1—Sn1 ⁱⁱⁱ	0.0	O1 ⁱⁱ —Sn1—C5—C10	41.0 (3)
C5—Sn1—C1—C2	151.7 (2)	C5 ⁱ —Sn1—C5—C6	-56.7 (4)
C5 ⁱ —Sn1—C1—C2	-22.3 (3)	C1—Sn1—C5—C6	129.6 (3)
O1—Sn1—C1—C2	-115.3 (3)	O1—Sn1—C5—C6	34.5 (3)
O1 ⁱⁱ —Sn1—C1—C2	64.7 (3)	O1 ⁱⁱ —Sn1—C5—C6	-142.0 (3)
C5—Sn1—C1—C2 ⁱ	22.3 (3)	C10—C5—C6—C7	-2.2 (5)
C5 ⁱ —Sn1—C1—C2 ⁱ	-151.7 (2)	Sn1—C5—C6—C7	-179.2 (3)
O1—Sn1—C1—C2 ⁱ	115.3 (3)	C5—C6—C7—C8	2.6 (5)
O1 ⁱⁱ —Sn1—C1—C2 ⁱ	-64.7 (3)	C6—C7—C8—C9	-1.8 (6)
C2 ⁱ —C1—C2—C3	-53.8 (5)	C7—C8—C9—C10	0.7 (6)
Sn1—C1—C2—C3	176.0 (3)	C6—C5—C10—C9	1.0 (6)
C1—C2—C3—C4	52.9 (5)	Sn1—C5—C10—C9	178.2 (3)
C2—C3—C4—C3 ⁱ	-52.3 (6)	C8—C9—C10—C5	-0.3 (7)
C5 ⁱ —Sn1—C5—C10	126.3 (3)		

Symmetry codes: (i) $-x+1, y, z$; (ii) $-x+1, -y+1, z+1/2$; (iii) $-x+1, -y+1, z-1/2$.