

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 3-(4-{6-methyl-4-[3-(trifluoromethyl)phenyl]pyridazin-3-yloxy}-phenyl)propanoate

Yu-Hong Xiao, You-Quan Zhu, Xiao-Mao Zou,*
Fang-Zhong Hu and Hua-Zheng Yang

State Key Laboratory and Institute of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, People's Republic of China

Correspondence e-mail: zyzq8165@nankai.edu.cn


Received 14 April 2008; accepted 6 May 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.055; wR factor = 0.147; data-to-parameter ratio = 11.0.

In the title compound, $\text{C}_{22}\text{H}_{19}\text{F}_3\text{N}_2\text{O}_3$, the benzene rings of the trifluoromethylphenyl and benzoylphenyl groups form dihedral angles of 41.89 (10) and 67.44 (10)°, respectively, with the pyridazine ring. The methylpropanoate group is nearly coplanar with the attached benzene ring [dihedral angle = 3.9 (2)°]. The trifluoromethyl group is disordered over two positions; the site-occupancy factors are *ca* 0.64 and 0.36. In the crystal structure, inversion-related molecules are linked through $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the biological activities of pyridazine derivatives, see: Heinisch & Kopelent (1992); Kolar & Tisler (1999).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{19}\text{F}_3\text{N}_2\text{O}_3$
 $M_r = 416.39$
 Triclinic, $P\bar{1}$
 $a = 9.4916$ (19) Å
 $b = 10.232$ (2) Å
 $c = 11.022$ (2) Å
 $\alpha = 81.70$ (3)°
 $\beta = 65.12$ (3)°
 $\gamma = 78.92$ (3)°
 $V = 950.6$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 113$ (2) K
 $0.20 \times 0.16 \times 0.06$ mm

Data collection

Rigaku Saturn CCD area-detector diffractometer
 Absorption correction: multi-scan (*CrystalClear*; Rigaku/MS, 2005)
 $T_{\min} = 0.977$, $T_{\max} = 0.993$
 5529 measured reflections
 3319 independent reflections
 1862 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.147$
 $S = 1.06$
 3319 reflections
 301 parameters
 72 restraints
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.40$ e Å⁻³
 $\Delta\rho_{\min} = -0.39$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C11}-\text{H11B}\cdots\text{O2}^i$	0.96	2.57	3.467 (4)	156
$\text{C22}-\text{H22C}\cdots\text{Cg1}^{ii}$	0.96	2.73	3.486 (4)	136

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x, -y + 1, -z + 1$. Cg1 is the centroid of the C2–C7 ring.

Data collection: *CrystalClear* (Rigaku/MS, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by the National Key Project for Basic Research (No. 20772067).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2581).

References

- Heinisch, G. & Kopelent, H. (1992). *Prog. Med. Chem.* **29**, 141–183.
 Kolar, P. & Tisler, M. (1999). *Adv. Heterocycl. Chem.* **75**, 167–241.
 Rigaku/MS (2005). *CrystalClear*. Rigaku/MS, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2008). E64, o1044 [doi:10.1107/S1600536808013342]

Methyl 3-(4-{6-methyl-4-[3-(trifluoromethyl)phenyl]pyridazin-3-yloxy}phenyl)-propanoate

Yu-Hong Xiao, You-Quan Zhu, Xiao-Mao Zou, Fang-Zhong Hu and Hua-Zheng Yang

S1. Comment

Many pyridazine derivatives have been found to exhibit biological activities such as insecticidal, fungicidal, herbicidal, plant-growth regulatory activity, *etc* (Heinisch & Kopelent, 1992). For example, pyridate, credazine and maleic hydrazide (Kolar & Tisler, 1999) have been commercialized as herbicides. In order to discover new biologically active pyridazine compounds, the title compound was synthesized and its structure is reported here.

In the title molecule (Fig. 1), the C2—C7 and C13—C18 benzene rings form dihedral angles of 41.89 (10)° and 67.44 (10)°, respectively, with the central pyridazine ring (C2—C4/C12/N1/N2). The methylpropanoate group is nearly coplanar with the C13—C18 benzene ring (dihedral angle 3.9 (2)°).


The crystal packing is stabilized by C—H···O hydrogen bonds and C—H··· π interactions (Table 1) involving the C2—C7 ring (centroid Cg1).

S2. Experimental

A mixture of methyl 3-(4-hydroxyphenyl)propanoate (2.3 mmol), 3-chloro-4-(3-(trifluoromethyl)phenyl)-6-methylpyridazine (2.3 mmol) and K₂CO₃ (5 mmol) was stirred at 313 K for 2 h. The crude product was recrystallized from ethanol and single crystals of the title compound suitable for X-ray analysis were grown from ethyl acetate-petroleum ether (3:1 v/v) at room temperature.

S3. Refinement

The trifluoromethyl group is disordered over two orientations (C1/F1/F2/F3 and C1/F1'/F2'/F3') with refined occupancies of 0.636 (12) and 0.364 (12). All C—F distances were restrained to 1.34 (1) Å and the U^{ij} components of disordered F atoms were restrained to be approximately isotropic. All H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

**Figure 1**

The molecular structure of the title compound, showing 30% probability displacement ellipsoids. Only one disorder component is shown.

Methyl 3-(4-(6-methyl-4-[3-(trifluoromethyl)phenyl]pyridazin-3-yloxy)phenyl)propanoate

Crystal data

$C_{22}H_{19}F_3N_2O_3$

$M_r = 416.39$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.4916 (19) \text{ \AA}$

$b = 10.232 (2) \text{ \AA}$

$c = 11.022 (2) \text{ \AA}$

$\alpha = 81.70 (3)^\circ$

$\beta = 65.12 (3)^\circ$

$\gamma = 78.92 (3)^\circ$

$V = 950.6 (4) \text{ \AA}^3$

$Z = 2$

$F(000) = 432$

$D_x = 1.455 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2351 reflections

$\theta = 2.0\text{--}27.9^\circ$

$\mu = 0.12 \text{ mm}^{-1}$

$T = 113 \text{ K}$

Plate, colourless

$0.20 \times 0.16 \times 0.06 \text{ mm}$

Data collection

Rigaku Saturn CCD area-detector
diffractometer

Radiation source: rotating anode

Confocal monochromator

Detector resolution: $7.31 \text{ pixels mm}^{-1}$

ω and φ scans

Absorption correction: multi-scan

(*CrystalClear*; Rigaku/MSC, 2005)

$T_{\min} = 0.977$, $T_{\max} = 0.993$

5529 measured reflections

3319 independent reflections

1862 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.0^\circ$

$h = -10 \rightarrow 11$

$k = -12 \rightarrow 11$

$l = -13 \rightarrow 11$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.147$
 $S = 1.06$
 3319 reflections
 301 parameters
 72 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0675P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.003$
 $\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.39 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
F1	1.0340 (5)	0.3434 (7)	0.0132 (7)	0.0359 (15)	0.636 (12)
F2	0.8747 (11)	0.3445 (7)	-0.0736 (7)	0.072 (2)	0.636 (12)
F3	0.7991 (6)	0.4176 (6)	0.1223 (8)	0.079 (3)	0.636 (12)
F1'	1.0297 (11)	0.3418 (14)	-0.0418 (14)	0.071 (4)	0.364 (12)
F2'	0.7900 (12)	0.3936 (8)	-0.0042 (15)	0.055 (3)	0.364 (12)
F3'	0.8553 (19)	0.4059 (12)	0.1421 (11)	0.076 (4)	0.364 (12)
O1	0.39298 (18)	0.1498 (2)	0.2418 (2)	0.0286 (6)	
O2	-0.5057 (2)	0.5985 (2)	0.6250 (2)	0.0409 (7)	
O3	-0.4466 (2)	0.7531 (2)	0.4543 (2)	0.0330 (6)	
N1	0.3114 (2)	-0.1545 (3)	0.1899 (3)	0.0246 (7)	
N2	0.2934 (2)	-0.0267 (3)	0.2179 (3)	0.0251 (7)	
C1	0.8892 (3)	0.3235 (3)	0.0444 (3)	0.0327 (9)	
C2	0.8618 (3)	0.1868 (3)	0.1025 (3)	0.0228 (8)	
C3	0.9650 (3)	0.1072 (3)	0.1502 (3)	0.0258 (8)	
H3	1.0518	0.1399	0.1464	0.031*	
C4	0.9393 (3)	-0.0202 (3)	0.2033 (3)	0.0266 (8)	
H4	1.0083	-0.0738	0.2361	0.032*	
C5	0.8108 (3)	-0.0688 (3)	0.2079 (3)	0.0241 (8)	
H5	0.7947	-0.1555	0.2430	0.029*	
C6	0.7056 (2)	0.0103 (3)	0.1608 (3)	0.0194 (7)	
C7	0.7313 (3)	0.1389 (3)	0.1080 (3)	0.0205 (7)	
H7	0.6616	0.1931	0.0763	0.025*	
C8	0.5697 (3)	-0.0462 (3)	0.1662 (3)	0.0191 (7)	
C9	0.5840 (3)	-0.1748 (3)	0.1379 (3)	0.0227 (7)	
H9	0.6818	-0.2278	0.1092	0.027*	

C10	0.4511 (3)	-0.2271 (3)	0.1519 (3)	0.0216 (7)
C11	0.4609 (3)	-0.3680 (3)	0.1257 (3)	0.0319 (9)
H11A	0.3575	-0.3868	0.1466	0.048*
H11B	0.5055	-0.4269	0.1806	0.048*
H11C	0.5259	-0.3811	0.0329	0.048*
C12	0.4159 (3)	0.0228 (3)	0.2087 (3)	0.0202 (7)
C13	0.2387 (3)	0.2146 (3)	0.3058 (3)	0.0260 (8)
C14	0.1408 (3)	0.1712 (3)	0.4317 (3)	0.0353 (10)
H14	0.1721	0.0931	0.4743	0.042*
C15	-0.0044 (3)	0.2448 (3)	0.4941 (3)	0.0340 (9)
H15	-0.0710	0.2150	0.5792	0.041*
C16	-0.0543 (3)	0.3619 (3)	0.4340 (3)	0.0215 (7)
C17	0.0473 (3)	0.4015 (3)	0.3070 (3)	0.0311 (9)
H17	0.0168	0.4794	0.2636	0.037*
C18	0.1928 (3)	0.3285 (3)	0.2428 (3)	0.0324 (9)
H18	0.2593	0.3568	0.1570	0.039*
C19	-0.2110 (3)	0.4412 (3)	0.5096 (3)	0.0250 (8)
H19A	-0.2905	0.3833	0.5370	0.030*
H19B	-0.2102	0.4673	0.5905	0.030*
C20	-0.2594 (3)	0.5645 (3)	0.4362 (3)	0.0287 (8)
H20A	-0.1815	0.6240	0.4094	0.034*
H20B	-0.2615	0.5396	0.3554	0.034*
C21	-0.4164 (3)	0.6373 (3)	0.5179 (3)	0.0246 (8)
C22	-0.5929 (3)	0.8355 (3)	0.5257 (4)	0.0392 (9)
H22A	-0.6786	0.7956	0.5298	0.059*
H22B	-0.5942	0.9226	0.4798	0.059*
H22C	-0.6034	0.8431	0.6150	0.059*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0233 (17)	0.036 (2)	0.049 (3)	-0.0158 (15)	-0.0157 (19)	0.011 (3)
F2	0.123 (5)	0.063 (4)	0.071 (4)	-0.058 (3)	-0.075 (4)	0.042 (3)
F3	0.041 (3)	0.028 (3)	0.102 (5)	0.003 (2)	0.027 (3)	0.006 (3)
F1'	0.052 (4)	0.035 (4)	0.069 (7)	-0.004 (3)	0.025 (4)	0.012 (5)
F2'	0.057 (4)	0.035 (4)	0.093 (7)	-0.009 (3)	-0.056 (4)	0.019 (4)
F3'	0.129 (8)	0.040 (5)	0.051 (5)	-0.008 (6)	-0.026 (6)	-0.014 (4)
O1	0.0136 (9)	0.0279 (13)	0.0422 (15)	0.0010 (8)	-0.0076 (9)	-0.0135 (11)
O2	0.0284 (11)	0.0378 (15)	0.0377 (16)	0.0032 (10)	-0.0012 (10)	0.0053 (12)
O3	0.0283 (10)	0.0297 (13)	0.0279 (14)	0.0109 (9)	-0.0045 (9)	-0.0039 (11)
N1	0.0212 (11)	0.0294 (15)	0.0253 (16)	-0.0046 (11)	-0.0108 (10)	-0.0031 (12)
N2	0.0198 (11)	0.0284 (16)	0.0286 (16)	-0.0037 (10)	-0.0104 (11)	-0.0046 (12)
C1	0.0232 (15)	0.038 (2)	0.037 (2)	-0.0089 (15)	-0.0131 (15)	0.0076 (19)
C2	0.0160 (12)	0.0280 (18)	0.0207 (18)	-0.0029 (11)	-0.0039 (11)	-0.0019 (14)
C3	0.0154 (12)	0.0360 (19)	0.0261 (19)	-0.0052 (12)	-0.0083 (12)	-0.0008 (15)
C4	0.0179 (13)	0.0323 (19)	0.030 (2)	-0.0012 (12)	-0.0133 (12)	0.0042 (15)
C5	0.0179 (12)	0.0241 (17)	0.0269 (19)	-0.0018 (11)	-0.0070 (12)	0.0003 (14)
C6	0.0124 (12)	0.0271 (17)	0.0154 (16)	-0.0008 (11)	-0.0022 (11)	-0.0053 (13)

C7	0.0146 (12)	0.0280 (18)	0.0161 (17)	0.0013 (11)	-0.0050 (11)	-0.0031 (13)
C8	0.0159 (12)	0.0283 (18)	0.0124 (17)	-0.0022 (12)	-0.0054 (11)	-0.0016 (14)
C9	0.0180 (12)	0.0278 (18)	0.0191 (18)	0.0007 (12)	-0.0059 (12)	-0.0024 (14)
C10	0.0226 (13)	0.0246 (17)	0.0169 (17)	-0.0042 (12)	-0.0069 (12)	-0.0013 (13)
C11	0.0278 (14)	0.0296 (19)	0.035 (2)	-0.0048 (13)	-0.0091 (14)	-0.0047 (16)
C12	0.0156 (12)	0.0246 (17)	0.0194 (18)	0.0002 (12)	-0.0059 (11)	-0.0059 (14)
C13	0.0140 (12)	0.0303 (18)	0.034 (2)	-0.0019 (12)	-0.0080 (12)	-0.0097 (15)
C14	0.0269 (15)	0.040 (2)	0.033 (2)	0.0083 (14)	-0.0122 (14)	0.0010 (17)
C15	0.0232 (14)	0.041 (2)	0.026 (2)	0.0040 (13)	-0.0049 (13)	0.0035 (16)
C16	0.0151 (12)	0.0253 (17)	0.0235 (18)	-0.0003 (11)	-0.0064 (12)	-0.0079 (14)
C17	0.0251 (14)	0.0235 (18)	0.031 (2)	0.0039 (13)	-0.0021 (13)	0.0000 (15)
C18	0.0244 (14)	0.0294 (19)	0.031 (2)	-0.0042 (13)	0.0008 (13)	0.0002 (16)
C19	0.0210 (13)	0.0270 (18)	0.0226 (18)	0.0019 (12)	-0.0057 (12)	-0.0051 (14)
C20	0.0255 (14)	0.0269 (18)	0.029 (2)	-0.0005 (12)	-0.0073 (13)	-0.0027 (15)
C21	0.0216 (13)	0.0245 (18)	0.027 (2)	-0.0018 (12)	-0.0091 (13)	-0.0046 (15)
C22	0.0351 (16)	0.035 (2)	0.039 (2)	0.0139 (14)	-0.0126 (15)	-0.0098 (17)

Geometric parameters (Å, °)

F1—C1	1.319 (5)	C8—C12	1.405 (3)
F2—C1	1.347 (5)	C9—C10	1.405 (4)
F3—C1	1.304 (7)	C9—H9	0.93
F1'—C1	1.302 (8)	C10—C11	1.489 (4)
F2'—C1	1.320 (6)	C11—H11A	0.96
F3'—C1	1.357 (9)	C11—H11B	0.96
O1—C12	1.353 (4)	C11—H11C	0.96
O1—C13	1.405 (3)	C13—C18	1.363 (4)
O2—C21	1.192 (3)	C13—C14	1.372 (4)
O3—C21	1.333 (3)	C14—C15	1.375 (4)
O3—C22	1.442 (3)	C14—H14	0.93
N1—C10	1.316 (3)	C15—C16	1.384 (4)
N1—N2	1.348 (4)	C15—H15	0.93
N2—C12	1.317 (3)	C16—C17	1.380 (4)
C1—C2	1.475 (4)	C16—C19	1.502 (4)
C2—C3	1.379 (3)	C17—C18	1.379 (4)
C2—C7	1.393 (4)	C17—H17	0.93
C3—C4	1.372 (4)	C18—H18	0.93
C3—H3	0.93	C19—C20	1.501 (4)
C4—C5	1.382 (4)	C19—H19A	0.97
C4—H4	0.93	C19—H19B	0.97
C5—C6	1.388 (3)	C20—C21	1.491 (4)
C5—H5	0.93	C20—H20A	0.97
C6—C7	1.382 (4)	C20—H20B	0.97
C6—C8	1.488 (4)	C22—H22A	0.96
C7—H7	0.93	C22—H22B	0.96
C8—C9	1.364 (4)	C22—H22C	0.96
C12—O1—C13	119.0 (2)	C10—C11—H11A	109.5

C21—O3—C22	116.2 (2)	C10—C11—H11B	109.5
C10—N1—N2	119.7 (2)	H11A—C11—H11B	109.5
C12—N2—N1	119.3 (2)	C10—C11—H11C	109.5
F1'—C1—F3	119.9 (8)	H11A—C11—H11C	109.5
F3—C1—F1	105.4 (5)	H11B—C11—H11C	109.5
F1'—C1—F2'	106.9 (7)	N2—C12—O1	117.9 (2)
F3—C1—F2'	69.2 (6)	N2—C12—C8	124.8 (3)
F1—C1—F2'	126.6 (5)	O1—C12—C8	117.2 (2)
F1'—C1—F2	76.7 (7)	C18—C13—C14	120.7 (3)
F3—C1—F2	109.6 (4)	C18—C13—O1	117.6 (2)
F1—C1—F2	103.9 (4)	C14—C13—O1	121.6 (2)
F1'—C1—F3'	104.6 (9)	C13—C14—C15	119.1 (3)
F1—C1—F3'	82.7 (7)	C13—C14—H14	120.5
F2'—C1—F3'	95.6 (7)	C15—C14—H14	120.5
F2—C1—F3'	132.0 (6)	C14—C15—C16	122.0 (3)
F1'—C1—C2	118.4 (6)	C14—C15—H15	119.0
F3—C1—C2	114.6 (4)	C16—C15—H15	119.0
F1—C1—C2	112.6 (4)	C17—C16—C15	117.1 (3)
F2'—C1—C2	117.7 (4)	C17—C16—C19	123.3 (2)
F2—C1—C2	110.1 (4)	C15—C16—C19	119.5 (2)
F3'—C1—C2	110.7 (6)	C18—C17—C16	121.6 (3)
C3—C2—C7	120.5 (3)	C18—C17—H17	119.2
C3—C2—C1	120.2 (2)	C16—C17—H17	119.2
C7—C2—C1	119.4 (2)	C13—C18—C17	119.6 (3)
C4—C3—C2	119.8 (2)	C13—C18—H18	120.2
C4—C3—H3	120.1	C17—C18—H18	120.2
C2—C3—H3	120.1	C20—C19—C16	116.4 (2)
C3—C4—C5	120.0 (2)	C20—C19—H19A	108.2
C3—C4—H4	120.0	C16—C19—H19A	108.2
C5—C4—H4	120.0	C20—C19—H19B	108.2
C4—C5—C6	120.8 (3)	C16—C19—H19B	108.2
C4—C5—H5	119.6	H19A—C19—H19B	107.3
C6—C5—H5	119.6	C21—C20—C19	113.1 (2)
C7—C6—C5	119.1 (2)	C21—C20—H20A	109.0
C7—C6—C8	121.7 (2)	C19—C20—H20A	109.0
C5—C6—C8	119.2 (2)	C21—C20—H20B	109.0
C6—C7—C2	119.8 (2)	C19—C20—H20B	109.0
C6—C7—H7	120.1	H20A—C20—H20B	107.8
C2—C7—H7	120.1	O2—C21—O3	123.5 (3)
C9—C8—C12	114.3 (2)	O2—C21—C20	125.7 (2)
C9—C8—C6	121.7 (2)	O3—C21—C20	110.9 (2)
C12—C8—C6	123.9 (3)	O3—C22—H22A	109.5
C8—C9—C10	120.0 (2)	O3—C22—H22B	109.5
C8—C9—H9	120.0	H22A—C22—H22B	109.5
C10—C9—H9	120.0	O3—C22—H22C	109.5
N1—C10—C9	121.7 (3)	H22A—C22—H22C	109.5
N1—C10—C11	116.6 (2)	H22B—C22—H22C	109.5
C9—C10—C11	121.7 (2)		

C10—N1—N2—C12	-1.1 (4)	N2—N1—C10—C11	178.8 (3)
F1'—C1—C2—C3	46.6 (11)	C8—C9—C10—N1	1.2 (4)
F3—C1—C2—C3	-104.0 (6)	C8—C9—C10—C11	-178.3 (3)
F1—C1—C2—C3	16.5 (6)	N1—N2—C12—O1	-177.8 (2)
F2'—C1—C2—C3	177.6 (8)	N1—N2—C12—C8	2.6 (4)
F2—C1—C2—C3	131.9 (6)	C13—O1—C12—N2	11.0 (4)
F3'—C1—C2—C3	-74.0 (8)	C13—O1—C12—C8	-169.4 (2)
F1'—C1—C2—C7	-133.2 (10)	C9—C8—C12—N2	-2.1 (4)
F3—C1—C2—C7	76.3 (6)	C6—C8—C12—N2	-178.0 (3)
F1—C1—C2—C7	-163.2 (5)	C9—C8—C12—O1	178.3 (3)
F2'—C1—C2—C7	-2.2 (9)	C6—C8—C12—O1	2.5 (4)
F2—C1—C2—C7	-47.8 (6)	C12—O1—C13—C18	-119.9 (3)
F3'—C1—C2—C7	106.2 (8)	C12—O1—C13—C14	64.9 (4)
C7—C2—C3—C4	0.2 (5)	C18—C13—C14—C15	-0.4 (5)
C1—C2—C3—C4	-179.6 (3)	O1—C13—C14—C15	174.6 (3)
C2—C3—C4—C5	0.5 (5)	C13—C14—C15—C16	-0.4 (5)
C3—C4—C5—C6	-0.8 (5)	C14—C15—C16—C17	0.9 (5)
C4—C5—C6—C7	0.5 (5)	C14—C15—C16—C19	-177.2 (3)
C4—C5—C6—C8	179.4 (3)	C15—C16—C17—C18	-0.5 (5)
C5—C6—C7—C2	0.2 (5)	C19—C16—C17—C18	177.4 (3)
C8—C6—C7—C2	-178.7 (3)	C14—C13—C18—C17	0.7 (5)
C3—C2—C7—C6	-0.5 (5)	O1—C13—C18—C17	-174.5 (3)
C1—C2—C7—C6	179.3 (3)	C16—C17—C18—C13	-0.3 (5)
C7—C6—C8—C9	139.4 (3)	C17—C16—C19—C20	4.1 (5)
C5—C6—C8—C9	-39.5 (4)	C15—C16—C19—C20	-178.0 (3)
C7—C6—C8—C12	-45.0 (4)	C16—C19—C20—C21	179.8 (3)
C5—C6—C8—C12	136.1 (3)	C22—O3—C21—O2	3.2 (5)
C12—C8—C9—C10	0.1 (4)	C22—O3—C21—C20	-177.6 (3)
C6—C8—C9—C10	176.1 (2)	C19—C20—C21—O2	-7.7 (5)
N2—N1—C10—C9	-0.8 (4)	C19—C20—C21—O3	173.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C11—H11 <i>B</i> ...O2 ⁱ	0.96	2.57	3.467 (4)	156
C22—H22 <i>C</i> ...C <i>g</i> 1 ⁱⁱ	0.96	2.73	3.486 (4)	136

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x, -y+1, -z+1$.