

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 5-Bromo-2-methylpyridine N-oxide

Bo-Nian Liu,<sup>a</sup> Shi-Gui Tang,<sup>b</sup> Hao-Yuan Li,<sup>a</sup> Ye-Ming Xu<sup>a</sup>  
and Cheng Guo<sup>a\*</sup><sup>a</sup>College of Science, Nanjing University of Technology, Xinfofan Road No. 5, Nanjing 210009, People's Republic of China, and <sup>b</sup>College of Life Science and Pharmaceutical Engineering, Nanjing University of Technology, Nanjing 210009, People's Republic of China

Correspondence e-mail: guocheng@njut.edu.cn


Received 29 April 2008; accepted 6 May 2008

Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(\text{C}-\text{C}) = 0.013$  Å;  $R$  factor = 0.064;  $wR$  factor = 0.163; data-to-parameter ratio = 15.5.

In the molecule of the title compound,  $\text{C}_6\text{H}_6\text{BrNO}$ , the methyl C and oxide O atoms lie in the pyridine ring plane, while the Br atom is displaced by 0.103 (3) Å. In the crystal structure, intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules into centrosymmetric dimers.

## Related literature

For related literature, see: Ochiai (1953).



## Experimental

## Crystal data

 $\text{C}_6\text{H}_6\text{BrNO}$   
 $M_r = 188.03$ Monoclinic,  $P2_1/n$   
 $a = 7.3060$  (15) Å $b = 11.351$  (2) Å  
 $c = 8.4950$  (17) Å  
 $\beta = 111.01$  (3)°  
 $V = 657.7$  (3) Å<sup>3</sup>  
 $Z = 4$ Mo  $K\alpha$  radiation  
 $\mu = 6.16$  mm<sup>-1</sup>  
 $T = 294$  (2) K  
 $0.10 \times 0.05 \times 0.05$  mm

## Data collection

Enraf-Nonius CAD-4  
diffractometer  
Absorption correction:  $\psi$  scan  
(North *et al.*, 1968)  
 $T_{\min} = 0.578$ ,  $T_{\max} = 0.748$   
1275 measured reflections1180 independent reflections  
747 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.036$   
3 standard reflections  
frequency: 120 min  
intensity decay: none

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.063$   
 $wR(F^2) = 0.162$   
 $S = 1.05$   
1180 reflections76 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.87$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.69$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}-\text{H1A}\cdots\text{O}^i$	0.93	2.41	3.264 (11)	153

Symmetry code: (i)  $-x + 1, -y + 2, -z + 2$ .

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the Center of Testing and Analysis, Nanjing University, for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2459).

## References

- Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.  
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
Ochiai, E. (1953). *J. Org. Chem.* **18**, 534–551.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

## supporting information

*Acta Cryst.* (2008). E64, o1060 [doi:10.1107/S1600536808013391]

## 5-Bromo-2-methylpyridine *N*-oxide

Bo-Nian Liu, Shi-Gui Tang, Hao-Yuan Li, Ye-Ming Xu and Cheng Guo

### S1. Comment

Some derivatives of pyridine are important chemical materials. We report herein the crystal structure of the title compound, (I).

In the molecule of (I), (Fig. 1), ring A (N/C1-C5) is, of course, planar. Br atom is at a distance of 0.103 (3) Å to the plane of ring A, while atoms O and C6 lie in the ring plane.


In the crystal structure, intermolecular C-H...O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers (Fig. 2), in which they may be effective in the stabilization of the structure.

### S2. Experimental

For the preparation of the title compound, 5-bromo-2-methylpyridine (80 g, 462 mmol) was suspended in glacial acetic acid (300 ml), aqueous hydrogen peroxide (35%) was added and the mixture was heated in a water-bath at 343-353 K. After 3 h a further hydrogen peroxide solution (35 ml) was added and the mixture was maintained an additional 9 h at the same temperature. The mixture was concentrated to about 100 ml, diluted with water (100 ml), and then again concentrated in vacuum as far as possible upon cooling to room temperature, a precipitate formed, which was collected by filtration, and then washed with cold ethanol (2 × 50 ml) to afford the ethyl ester as a white solid (yield; 83 g, 95%) (Ochiai, 1953). Crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of a methanol solution.


### S3. Refinement

H atoms were positioned geometrically, with C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms with  $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$ , where  $x = 1.5$  for methyl H, and  $x = 1.2$  for aromatic H atoms.



**Figure 1**

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

A partial packing diagram of (I). Hydrogen bonds are shown as dashed lines.

### 5-Bromo-2-methylpyridine N-oxide

#### *Crystal data*

$C_6H_6BrNO$

$M_r = 188.03$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2_1n$

$a = 7.3060$  (15) Å

$b = 11.351$  (2) Å

$c = 8.4950$  (17) Å

$\beta = 111.01$  (3)°

$V = 657.7$  (3) Å<sup>3</sup>

$Z = 4$

$F(000) = 368$

$D_x = 1.899$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 25 reflections

$\theta = 10\text{--}13^\circ$

$\mu = 6.16$  mm<sup>-1</sup>

$T = 294$  K

Block, colorless

$0.10 \times 0.05 \times 0.05$  mm

*Data collection*

Enraf–Nonius CAD-4 diffractometer	1180 independent reflections 747 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.036$
Graphite monochromator	$\theta_{\text{max}} = 25.2^\circ$ , $\theta_{\text{min}} = 3.1^\circ$
$\omega/2\theta$ scans	$h = -8 \rightarrow 8$
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	$k = 0 \rightarrow 13$
$T_{\text{min}} = 0.578$ , $T_{\text{max}} = 0.748$	$l = 0 \rightarrow 10$
1275 measured reflections	3 standard reflections every 120 min intensity decay: none

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.063$	H-atom parameters constrained
$wR(F^2) = 0.162$	$w = 1/[\sigma^2(F_o^2) + (0.08P)^2 + P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
1180 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
76 parameters	$\Delta\rho_{\text{max}} = 0.87 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.69 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.17025 (14)	0.74086 (7)	0.64566 (12)	0.0448 (4)
N	0.6201 (9)	0.9804 (5)	0.7872 (8)	0.0290 (15)
O	0.6767 (10)	1.0662 (5)	0.8994 (8)	0.0506 (18)
C1	0.4547 (13)	0.9187 (7)	0.7722 (10)	0.038 (2)
H1A	0.3842	0.9381	0.8405	0.046*
C2	0.3901 (11)	0.8283 (6)	0.6579 (10)	0.0289 (18)
C3	0.4939 (15)	0.7989 (8)	0.5576 (12)	0.045 (2)
H3A	0.4523	0.7389	0.4781	0.053*
C4	0.6635 (12)	0.8623 (7)	0.5795 (11)	0.038 (2)
H4A	0.7354	0.8418	0.5127	0.045*
C5	0.7341 (13)	0.9535 (7)	0.6929 (10)	0.036 (2)
C6	0.9107 (13)	1.0232 (8)	0.7239 (12)	0.047
H6A	0.9754	0.9981	0.6493	0.071*
H6B	0.9971	1.0126	0.8387	0.071*
H6C	0.8761	1.1050	0.7043	0.071*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br	0.0512 (6)	0.0387 (6)	0.0354 (6)	-0.0061 (5)	0.0045 (4)	0.0010 (5)
N	0.035 (4)	0.022 (3)	0.022 (4)	0.009 (3)	0.000 (3)	-0.006 (3)
O	0.072 (5)	0.042 (4)	0.032 (4)	-0.021 (3)	0.011 (4)	-0.016 (3)
C1	0.050 (6)	0.030 (4)	0.027 (5)	-0.004 (4)	0.004 (4)	-0.002 (4)
C2	0.032 (4)	0.020 (4)	0.025 (4)	0.006 (3)	-0.002 (4)	0.006 (3)
C3	0.050 (6)	0.035 (5)	0.036 (5)	0.003 (5)	0.001 (5)	-0.014 (4)
C4	0.038 (5)	0.037 (5)	0.034 (5)	0.013 (4)	0.009 (4)	-0.008 (4)
C5	0.051 (5)	0.025 (4)	0.021 (5)	0.011 (4)	0.000 (4)	0.003 (4)
C6	0.047	0.047	0.047	0.000	0.017	0.000

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Br—C2	1.859 (8)	C3—H3A	0.9300
N—O	1.322 (8)	C4—C5	1.382 (12)
N—C1	1.362 (10)	C4—H4A	0.9300
N—C5	1.381 (10)	C5—C6	1.455 (12)
C1—C2	1.375 (11)	C6—H6A	0.9600
C1—H1A	0.9300	C6—H6B	0.9600
C2—C3	1.369 (12)	C6—H6C	0.9600
C3—C4	1.387 (12)		
O—N—C1	118.9 (7)	C5—C4—C3	125.0 (8)
O—N—C5	118.9 (7)	C5—C4—H4A	117.5
C1—N—C5	122.1 (7)	C3—C4—H4A	117.5
N—C1—C2	121.2 (8)	N—C5—C4	114.7 (8)
N—C1—H1A	119.4	N—C5—C6	117.1 (7)
C2—C1—H1A	119.4	C4—C5—C6	128.2 (8)
C3—C2—C1	119.7 (8)	C5—C6—H6A	109.5
C3—C2—Br	119.7 (6)	C5—C6—H6B	109.5
C1—C2—Br	120.6 (6)	H6A—C6—H6B	109.5
C2—C3—C4	117.3 (8)	C5—C6—H6C	109.5
C2—C3—H3A	121.4	H6A—C6—H6C	109.5
C4—C3—H3A	121.4	H6B—C6—H6C	109.5
O—N—C1—C2	-179.7 (7)	O—N—C5—C4	179.7 (7)
C5—N—C1—C2	-1.9 (12)	C1—N—C5—C4	2.0 (11)
N—C1—C2—C3	0.3 (12)	O—N—C5—C6	-0.3 (11)
N—C1—C2—Br	177.3 (6)	C1—N—C5—C6	-178.1 (7)
C1—C2—C3—C4	1.0 (13)	C3—C4—C5—N	-0.5 (13)
Br—C2—C3—C4	-176.0 (6)	C3—C4—C5—C6	179.5 (9)
C2—C3—C4—C5	-0.9 (14)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C1—H1A $\cdots$ O <sup>i</sup>	0.93	2.41	3.264 (11)	153

Symmetry code: (i)  $-x+1, -y+2, -z+2$ .