

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-Amino-2,4,6-triiodoisophthalic acid monohydrate

Tobias Beck* and George M. Sheldrick

Department of Structural Chemistry, Georg-August Universität, Tammannstrasse 4, 37077 Göttingen, Germany

Correspondence e-mail: tbeck@shelx.uni-ac.gwdg.de


Received 22 May 2008; accepted 11 June 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.026; wR factor = 0.062; data-to-parameter ratio = 15.7.

The title compound, $\text{C}_8\text{H}_4\text{I}_3\text{NO}_4 \cdot \text{H}_2\text{O}$, shows an extensive hydrogen-bond network; in the crystal structure, molecules are linked by $\text{O}-\text{H} \cdots \text{O}$, $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds involving all possible donors and also the water molecule.

Related literature

For the synthetic procedure, see Larsen *et al.* (1956). For related crystal structure determinations: 1,3,5-triiodobenzene, see: Margraf & Bats (2006); sodium diatrizoate, see: Tonnessen *et al.* (1996). For the 1,3,5-triiodobenzene core as the basis of contrast agents, see: Yu & Watson (1999).



Experimental

Crystal data

 $\text{C}_8\text{H}_4\text{I}_3\text{NO}_4 \cdot \text{H}_2\text{O}$ $M_r = 576.84$ Orthorhombic, *Pbca* $a = 9.214$ (1) Å $b = 15.735$ (2) Å $c = 18.816$ (2) Å $V = 2728.0$ (5) Å³ $Z = 8$ Cu $K\alpha$ radiation $\mu = 54.11$ mm⁻¹ $T = 100$ (2) K $0.08 \times 0.05 \times 0.03$ mm

Data collection

Bruker SMART 6000 diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.106$, $T_{\max} = 0.345$
(expected range = 0.061–0.197)

49139 measured reflections
2716 independent reflections
2545 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.026$ $wR(F^2) = 0.061$ $S = 1.03$

2716 reflections

173 parameters

14 restraints

Only H-atom coordinates refined

 $\Delta\rho_{\text{max}} = 0.71$ e Å⁻³ $\Delta\rho_{\text{min}} = -1.71$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O11–H11 ⁱ ··O14	0.79 (5)	1.75 (5)	2.540 (5)	173 (7)
O8–H8··O12 ⁱ	0.80 (5)	1.90 (5)	2.662 (5)	161 (7)
O14–H14A··O9 ⁱⁱ	0.81 (4)	1.95 (4)	2.751 (5)	170 (6)
O14–H14B··N13 ⁱⁱⁱ	0.81 (4)	2.05 (4)	2.841 (5)	166 (6)
N13–H13A··O14 ^{iv}	0.88 (4)	2.30 (5)	3.067 (6)	147 (5)
N13–H13B··O12 ^{iv}	0.88 (4)	2.68 (5)	3.478 (5)	152 (5)

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{3}{2}, -z$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, z$; (iii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y + \frac{1}{2}, z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* and *SHELXTL*.

Financial support of the ICDD (Ludo Frevel Scholarship Award 2008 for TB) and DFG (IRTG 1422) is gratefully acknowledged. The authors thank Regine Herbst-Irmer and Stephan Rühl for advice regarding the refinement.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2102).

References

- Bruker (2007). *APEX* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Larsen, A. A., Moore, C., Sprague, J., Cloke, B., Moss, J. & Hoppe, J. O. (1956). *J. Am. Chem. Soc.* **78**, 3210–3216.
- Margraf, D. & Bats, J. W. (2006). *Acta Cryst.* **E62**, o502–o504.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tonnessen, L. E., Pedersen, B. F. & Klaveness, J. (1996). *Acta Chem. Scand.* **50**, 603–608.
- Yu, S. B. & Watson, A. D. (1999). *Chem. Rev.* **99**, 2353–2378.

supporting information

Acta Cryst. (2008). E64, o1286 [doi:10.1107/S1600536808017741]

5-Amino-2,4,6-triiodoisophthalic acid monohydrate

Tobias Beck and George M. Sheldrick

S1. Comment

Iodine-based compounds have always been in the focus of contrast agents for X-ray imaging. The 1,3,5-triiodobenzene core has been the basis of many contrast agents (Yu & Watson 1999). The ionic monomer diatrizoate was one of the first compounds used (Tonnessen *et al.* 1996).

The title compound, 5-Amino-2,4,6-triiodoisophthalic acid (hereafter I3C), crystallizes as a monohydrate, due to water impurities in the crystallization solution. It forms hydrogen bonds with all potential donors as well as the lattice water being involved (Fig. 2, Table 1). However, the interaction between N13 and O12 is slightly weaker. In the crystal, the molecules are positioned perpendicular to each other, showing no π - π interactions of the phenyl rings (Fig. 3).


The three functional groups for hydrogen bonding, along with the three iodine atoms, render I3C a suitable agent for experimental phasing of macromolecules (Beck *et al.*, unpublished results). The iodine atoms give rise to a large anomalous signal, even at in-house sources. Additionally, they form an equilateral triangle (I—I 6.0 Å) which is easy to recognize in the heavy atom substructure when this compound is used as a heavy atom derivative for macromolecular phasing.

S2. Experimental


The title compound was prepared according to the reported procedure (Larsen *et al.* 1956). It was recrystallized from a methanol-acetonitrile solution by slowly evaporating the solvents to obtain crystals suitable for X-ray single-crystal diffraction.

S3. Refinement


Hydrogen atoms were located *via* the difference Fourier map and their geometrical positions were refined with restraints. The U values were set to 1.5 U_{eq} of their parent atom. Bond lengths for hydrogen atoms were restrained to be equal (SADI in SHELXL-97). Phenyl ring and carboxylate groups were restrained to planarity.

**Figure 1**


A view of I3C. Displacement ellipsoids are drawn at the 50% probability level. The hydrogen bond within the asymmetric unit is shown as a dashed line.

**Figure 2**

Hydrogen bonding of I3C. Symmetry equivalents are depicted in orange.

**Figure 3**

Packing of I3C, viewed along b. Hydrogen atoms are omitted for clarity. In alternating layers molecules are positioned perpendicular to each other. Hydrogen bonds are shown as dashed lines.

**Figure 4**

Synthetic scheme of I3C.

5-Amino-2,4,6-triiodoisophthalic acid monohydrate

Crystal data

C₈H₄I₃NO₄·H₂O $M_r = 576.84$ Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

 $a = 9.214 (1) \text{ \AA}$ $b = 15.735 (2) \text{ \AA}$ $c = 18.816 (2) \text{ \AA}$ $V = 2728.0 (5) \text{ \AA}^3$ $Z = 8$ $F(000) = 2080$ $D_x = 2.809 \text{ Mg m}^{-3}$ Cu $K\alpha$ radiation, $\lambda = 1.54178 \text{ \AA}$

Cell parameters from 9945 reflections

 $\theta = 4.7\text{--}60.8^\circ$ $\mu = 54.11 \text{ mm}^{-1}$ $T = 100 \text{ K}$

Block, yellow

 $0.08 \times 0.05 \times 0.03 \text{ mm}$

Data collection

Bruker SMART 6000

diffractometer

Radiation source: rotating anode

INCOATEC multilayer optics monochromator

Detector resolution: $5.602 \text{ pixels mm}^{-1}$ ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.106$, $T_{\max} = 0.345$

49139 measured reflections

2716 independent reflections

2545 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.043$ $\theta_{\max} = 74.3^\circ$, $\theta_{\min} = 4.7^\circ$ $h = -10 \rightarrow 10$ $k = -17 \rightarrow 19$ $l = -23 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.026$ $wR(F^2) = 0.061$ $S = 1.03$

2716 reflections

173 parameters

14 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: difference Fourier map

Only H-atom coordinates refined

 $w = 1/[\sigma^2(F_o^2) + (0.0277P)^2 + 17.3624P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.002$ $\Delta\rho_{\max} = 0.72 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -1.71 \text{ e \AA}^{-3}$ Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.000058 (8)

Special details

Experimental. Intensities were measured with a Bruker SMART 6000 area-detector**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.**Refinement.** Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. Hydrogen atoms were located via the difference Fourier map and their geometrical positions were refined with restraints. The U values were set to $1.5 U_{\text{eq}}$ of their parent atom.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	1.18424 (3)	0.630616 (18)	0.012748 (16)	0.02419 (10)
I2	0.71499 (3)	0.674286 (18)	0.234093 (15)	0.02375 (10)
I3	1.04238 (3)	0.979824 (17)	0.124476 (17)	0.02693 (10)
C1	1.0827 (5)	0.7970 (3)	0.0781 (2)	0.0161 (8)
C2	1.0606 (5)	0.7094 (3)	0.0789 (2)	0.0161 (9)
C3	0.9544 (5)	0.6745 (3)	0.1231 (2)	0.0153 (9)
C4	0.8655 (5)	0.7276 (3)	0.1631 (2)	0.0163 (9)
C5	0.8803 (5)	0.8172 (3)	0.1604 (2)	0.0165 (9)
C6	0.9945 (5)	0.8492 (3)	0.1199 (2)	0.0180 (9)
C7	1.1981 (5)	0.8362 (3)	0.0320 (3)	0.0202 (10)
O8	1.1458 (4)	0.8648 (2)	−0.02803 (18)	0.0256 (7)
H8	1.205 (6)	0.883 (4)	−0.055 (3)	0.038*
O9	1.3242 (4)	0.8414 (2)	0.05017 (19)	0.0290 (8)
C10	0.9364 (5)	0.5794 (3)	0.1285 (2)	0.0173 (9)
O11	1.0376 (4)	0.5445 (2)	0.16663 (17)	0.0224 (7)
H11	1.031 (7)	0.494 (3)	0.170 (3)	0.034*
O12	0.8358 (4)	0.54161 (19)	0.10032 (17)	0.0213 (7)
N13	0.7889 (5)	0.8702 (2)	0.1997 (2)	0.0210 (8)
H13A	0.698 (5)	0.855 (4)	0.204 (3)	0.031*
H13B	0.785 (6)	0.922 (3)	0.183 (3)	0.031*
O14	1.0395 (4)	0.3834 (2)	0.17540 (18)	0.0235 (7)
H14A	1.074 (7)	0.366 (4)	0.139 (3)	0.035*
H14B	1.099 (6)	0.376 (4)	0.206 (3)	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.02850 (18)	0.02100 (16)	0.02307 (16)	0.00785 (12)	0.00724 (12)	0.00027 (11)
I2	0.02603 (18)	0.02156 (16)	0.02364 (16)	−0.00544 (11)	0.00910 (12)	−0.00016 (11)
I3	0.02852 (19)	0.01260 (15)	0.03966 (19)	−0.00146 (11)	0.00662 (13)	0.00096 (11)
C1	0.018 (2)	0.0143 (19)	0.0162 (19)	0.0012 (17)	0.0007 (17)	0.0000 (16)
C2	0.019 (2)	0.014 (2)	0.0150 (19)	0.0033 (17)	−0.0002 (17)	−0.0002 (16)
C3	0.020 (2)	0.013 (2)	0.0135 (19)	0.0006 (17)	−0.0052 (17)	0.0019 (15)
C4	0.018 (2)	0.016 (2)	0.0150 (19)	−0.0072 (16)	0.0005 (17)	0.0017 (16)
C5	0.018 (2)	0.015 (2)	0.016 (2)	0.0003 (17)	−0.0023 (17)	−0.0006 (16)
C6	0.023 (2)	0.013 (2)	0.017 (2)	−0.0022 (18)	−0.0013 (18)	0.0009 (16)
C7	0.023 (3)	0.014 (2)	0.023 (2)	−0.0007 (17)	0.0010 (19)	−0.0016 (17)
O8	0.0251 (19)	0.0276 (18)	0.0241 (17)	−0.0019 (14)	0.0036 (15)	0.0093 (14)
O9	0.0216 (19)	0.038 (2)	0.0276 (18)	−0.0045 (15)	0.0035 (15)	0.0013 (15)
C10	0.023 (2)	0.014 (2)	0.015 (2)	−0.0007 (18)	0.0048 (18)	0.0001 (16)
O11	0.0274 (18)	0.0132 (15)	0.0265 (17)	−0.0007 (13)	−0.0074 (14)	0.0043 (13)
O12	0.0279 (18)	0.0121 (14)	0.0239 (16)	−0.0039 (13)	−0.0064 (14)	0.0014 (12)
N13	0.021 (2)	0.0148 (18)	0.027 (2)	0.0019 (15)	0.0035 (17)	−0.0026 (15)
O14	0.0255 (19)	0.0217 (16)	0.0233 (17)	0.0028 (14)	−0.0003 (14)	0.0027 (14)

Geometric parameters (Å, °)

I1—C2	2.094 (4)	C5—N13	1.396 (6)
I2—C4	2.100 (4)	C7—O9	1.214 (6)
I3—C6	2.103 (4)	C7—O8	1.308 (6)
C1—C2	1.394 (6)	O8—H8	0.80 (5)
C1—C6	1.398 (6)	C10—O12	1.223 (6)
C1—C7	1.504 (6)	C10—O11	1.299 (6)
C2—C3	1.396 (6)	O11—H11	0.79 (5)
C3—C4	1.391 (6)	N13—H13A	0.88 (4)
C3—C10	1.509 (6)	N13—H13B	0.88 (4)
C4—C5	1.418 (6)	O14—H14A	0.81 (4)
C5—C6	1.393 (6)	O14—H14B	0.81 (4)
C2—C1—C6	119.4 (4)	C5—C6—C1	122.3 (4)
C2—C1—C7	121.0 (4)	C5—C6—I3	119.3 (3)
C6—C1—C7	119.6 (4)	C1—C6—I3	118.4 (3)
C1—C2—C3	119.8 (4)	O9—C7—O8	125.0 (4)
C1—C2—I1	120.0 (3)	O9—C7—C1	122.7 (4)
C3—C2—I1	120.2 (3)	O8—C7—C1	112.3 (4)
C4—C3—C2	120.0 (4)	C7—O8—H8	115 (5)
C4—C3—C10	119.6 (4)	O12—C10—O11	125.3 (4)
C2—C3—C10	120.4 (4)	O12—C10—C3	122.5 (4)
C3—C4—C5	121.3 (4)	O11—C10—C3	112.3 (4)
C3—C4—I2	119.6 (3)	C10—O11—H11	114 (5)
C5—C4—I2	119.0 (3)	C5—N13—H13A	117 (4)
C6—C5—N13	122.0 (4)	C5—N13—H13B	113 (4)
C6—C5—C4	116.9 (4)	H13A—N13—H13B	105 (6)
N13—C5—C4	121.0 (4)	H14A—O14—H14B	108 (6)
C6—C1—C2—C3	−2.2 (6)	N13—C5—C6—C1	−177.4 (4)
C7—C1—C2—C3	179.0 (4)	C4—C5—C6—C1	6.0 (6)
C6—C1—C2—I1	176.7 (3)	N13—C5—C6—I3	5.2 (5)
C7—C1—C2—I1	−2.0 (5)	C4—C5—C6—I3	−171.4 (3)
C1—C2—C3—C4	3.5 (6)	C2—C1—C6—C5	−2.8 (6)
I1—C2—C3—C4	−175.4 (3)	C7—C1—C6—C5	176.0 (4)
C1—C2—C3—C10	−175.6 (4)	C2—C1—C6—I3	174.7 (3)
I1—C2—C3—C10	5.5 (5)	C7—C1—C6—I3	−6.5 (5)
C2—C3—C4—C5	−0.1 (6)	C2—C1—C7—O9	−84.0 (6)
C10—C3—C4—C5	179.1 (4)	C6—C1—C7—O9	97.2 (5)
C2—C3—C4—I2	−176.2 (3)	C2—C1—C7—O8	97.3 (5)
C10—C3—C4—I2	3.0 (5)	C6—C1—C7—O8	−81.5 (5)
C3—C4—C5—C6	−4.6 (6)	C4—C3—C10—O12	75.4 (5)
I2—C4—C5—C6	171.5 (3)	C2—C3—C10—O12	−105.5 (5)
C3—C4—C5—N13	178.8 (4)	C4—C3—C10—O11	−103.6 (4)
I2—C4—C5—N13	−5.1 (5)	C2—C3—C10—O11	75.5 (5)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O11—H11 \cdots O14	0.79 (5)	1.75 (5)	2.540 (5)	173 (7)
O8—H8 \cdots O12 ⁱ	0.80 (5)	1.90 (5)	2.662 (5)	161 (7)
O14—H14 <i>A</i> \cdots O9 ⁱⁱ	0.81 (4)	1.95 (4)	2.751 (5)	170 (6)
O14—H14 <i>B</i> \cdots N13 ⁱⁱⁱ	0.81 (4)	2.05 (4)	2.841 (5)	166 (6)
N13—H13 <i>A</i> \cdots O14 ^{iv}	0.88 (4)	2.30 (5)	3.067 (6)	147 (5)
N13—H13 <i>B</i> \cdots O12 ^{iv}	0.88 (4)	2.68 (5)	3.478 (5)	152 (5)

Symmetry codes: (i) $x+1/2, -y+3/2, -z$; (ii) $-x+5/2, y-1/2, z$; (iii) $-x+2, y-1/2, -z+1/2$; (iv) $-x+3/2, y+1/2, z$.