

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2R)-Ethyl 2-(5-bromo-2,3-dioxoindolin-1-yl)propanoate

Alexander V. Kurkin,* Anna A. Bernovskaya, Marina A. Yurovskaya and Victor B. Rybakov

Department of Chemistry, Moscow State University, 119991 Moscow, Russian Federation

Correspondence e-mail: kurkin@direction.chem.msu.ru


Received 10 June 2008; accepted 3 July 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.052; wR factor = 0.140; data-to-parameter ratio = 15.8.

The title compound, $\text{C}_{13}\text{H}_{12}\text{BrNO}_4$, was obtained from an optically active aniline derivative. The structure was characterized by ^1H NMR, ^{13}C NMR, MS and X-ray diffraction techniques. 86% of the atoms of the two independent molecules in the asymmetric unit show non-crystallographic inversion symmetry.

Related literature

For related structures, see: Akkurt *et al.* (2006); Miehe *et al.* (1991); Robeyns *et al.* (2007). For general background, see: Sandmeyer (1919); Silva *et al.* (2001); Spek (2003).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{12}\text{BrNO}_4$
 $M_r = 326.14$
 Monoclinic, $P2_1$

$a = 9.7390$ (13) Å
 $b = 14.355$ (2) Å
 $c = 9.8361$ (10) Å

$\beta = 95.779$ (9)°
 $V = 1368.1$ (3) Å³
 $Z = 4$
 Cu $K\alpha$ radiation

$\mu = 4.20$ mm⁻¹
 $T = 293$ (2) K
 $0.20 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.385$, $T_{\max} = 0.432$
 6047 measured reflections

5502 independent reflections
 3935 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$
 1 standard reflection
 frequency: 60 min
 intensity decay: 2%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.140$
 $S = 1.02$
 5502 reflections
 348 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.45$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.38$ e Å⁻³
 Absolute structure: Flack (1983),
 2566 Friedel pairs
 Flack parameter: -0.06 (3)

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are indebted to the Russian Foundation for Basic Research for covering the licence fee for use of the Cambridge Structural Database.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2723).

References

- Akkurt, M., Türktekin, S., Jarrahpour, A. A., Khalili, D. & Büyükgüngör, O. (2006). *Acta Cryst.* **E62**, o1575–o1577.
 Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
 Miehe, G., Susse, P., Kupcik, V., Egert, E., Nieger, M., Kunz, G., Gerke, R., Knieriem, B., Niemeyer, M. & Luttkie, W. (1991). *Angew. Chem. Int. Ed. Engl.* **30**, 964–967.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Robeyns, K., Rohand, T., Bouhfid, R., Essassi, E. L. M. & Van Meervelt, L. (2007). *Acta Cryst.* **E63**, o1747–o1748.
 Sandmeyer, T. (1919). *Helv. Chim. Acta*, **2**, 234–242.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Silva, J. F., Garden, S. J. & Pinto, A. C. (2001). *J. Braz. Chem. Soc.* **12**, 273–324.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2008). E64, o1448 [doi:10.1107/S1600536808020588]

(2R)-Ethyl 2-(5-bromo-2,3-dioxindolin-1-yl)propanoate

Alexander V. Kurkin, Anna A. Bernovskaya, Marina A. Yurovskaya and Victor B. Rybakov

S1. Comment

Nowadays, much attention has been focused on isatin derivatives for their broad-spectrum biological and pharmacological activities, such as antibacterial, antiprotozoal, antifungal, antiviral, anti-HIV, anticonvulsant, antihelminthic activities, influence *CNS*, participate in metabolism and stimulate growth of plants (Silva *et al.*, 2001). There is a modern tendency to use pure enantiomers of heterocyclic compounds instead of their racemic mixtures, for example, as starting materials in preparation of pharmaceuticals. It is true for the derivatives of isatin. In this paper, we report the synthesis and crystal structure of the ethyl (2*R*)-2-(5-bromisatin-1-yl)propanoate.

The asymmetric unit of the title compound has two independent molecules (hereafter called A and B), which depicted in Fig. 1. The ADDSYM test by *PLATON* (Spek, 2003), shown a noncrystallographic inversion.

In the principle, the geometric parameters of heterobicycle are closely agree with ones in molecular structures of ethyl 2-(2,3-dioxindolin-1-yl)acetate (Robeyns *et al.*, 2007), *N*-benzylindole-2,3-dion (*N*-benzylisatin) (Akkurt *et al.*, 2006) and 1-methyl-1*H*-indole-2,3-dione (Miehe *et al.*, 1991).

The short interatomic contacts O12*a*...C3*b* = 3.01 Å and O12*b*...C3*a* = 2.98 Å were found in the crystal structure.

S2. Experimental

We have synthesized ethyl (2*R*)-2-(5-bromisatin-1-yl)propanoate from optically active aniline by Sandmeyer method (Sandmeyer, 1919) (Fig. 2). A mixture of solutions of chloralhydrate (0.003 mol) in water (5.1 ml), a solution of ethyl (*R*)-*N*-(4-bromophenylamino)propanoate (0.0018 mol) in water (1.23 ml) with concentrated hydrochloric acid (0.26 g), a solution of hydroxylamine hydrochloride (0.0061 mol) in water (1.03 ml) and Na₂SO₄ (0.42 g), was stirred at reflux for 1–2 min. In addition we have used ethanol as a solvent to increase aniline solubility. The reaction mixture was cooled to r.t., extracted with CH₂Cl₂ and dried over anhydrous Na₂SO₄. The solvent was evaporated under reduced pressure to afford isonitroso-substance as brown oil (83%). The isonitroso-substance (0.0015 mol) was added to concentrate sulfuric acid (1.46 g) at 323 K so that the temperature of the reaction mixture did not exceed 343 K. The reaction mixture was stirred at 353 K for 10–15 min. The resulting mixture was cooled to r. t., diluted with cold water, extracted with CH₂Cl₂ and dried over anhydrous Na₂SO₄. The solvent was evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel (eluent - ethylacetate/petroleum ether: 10/1) to afford ethyl (2*R*)-2-(5-bromisatin-1-yl)propanoate (60%) as a red solid, its enantiomeric purity determinates by HPLC with chiral stationary phase achieved 97% *ee*. *M.p.* 404–405 K.

The ¹H NMR (CDCl₃), δ, p.p.m., J (Hz): 1.15 (t, J=7.0, 3H, —CH₂—CH₃), 1.53 (d, J=7.1, 3H, —CH—CH₃), 4.08–4.21 (m, 2H, —CH₂—CH₃), 5.15 (q, J=7.1, 1H, —CH—CH₃), 7.12 (d, J=8.5, 1H, 7-H), 7.77 (s, 1H, 4-H), 7.85 (d, J=8.1, 1H, 6-H). ¹³C NMR (DMSO-*d*₆), δ, p.p.m.: 14.19 (CH₃), 14.42 (CH₃), 49.55 (CH), 61.89 (CH₂), 113.94 (CH), 115.71 (C), 119.92 (C), 127.52 (CH), 140.41 (CH), 148.95 (C), 157.70 (C=O), 169.73 (C=O), 181.87 (C=O). Mass-spectr., *m/z* (I, %): 224 [*M*⁺—CH₃CHCO₂Et], (10), 145 (2), 117 (8), 91 (36), 41 (39).

S3. Refinement

In the compound **I** hydrogen atoms bonded to C-atoms were included in calculated positions and refined as riding atoms. Calculated C—H bond lengths are in the range of 0.93–0.97 Å. For methyl H-atoms U_{iso} values were set equal to $1.5U_{\text{eq}}$ of the carrier atoms, for other H-atoms U_{iso} values were set to $1.2U_{\text{eq}}$ of the carrier atoms.




Figure 1

ORTEP-3 (Farrugia, 1997) plot of the molecules (A and B) of compound **I** with the numbering scheme. Thermal displacement ellipsoids are shown at the 30% probability level. H atoms are drawn as a small spheres of arbitrary radius.




Figure 2

Synthesis of ethyl (2*R*)-2-(5-bromisatin-1-yl)propanoate, I.

(2*R*)-Ethyl 2-(5-bromo-2,3-dioxindolin-1-yl)propanoate

Crystal data

$C_{13}H_{12}BrNO_4$

$M_r = 326.14$

Monoclinic, $P2_1$

Hall symbol: $P\ 2_1yb$

$a = 9.7390$ (13) Å

$b = 14.355$ (2) Å

$c = 9.8361$ (10) Å

$\beta = 95.779$ (9)°

$V = 1368.1$ (3) Å³

$Z = 4$

$F(000) = 656$

$D_x = 1.583$ Mg m⁻³

Melting point: 404.5 K

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 25 reflections

$\theta = 32.2$ – 34.4 °

$\mu = 4.20$ mm⁻¹

$T = 293$ K

Prism, red

$0.20 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: Fine-focus sealed tube

Graphite monochromator

Non-profiled ω scans

Absorption correction: ψ scan

(North *et al.*, 1968)

$T_{\min} = 0.385$, $T_{\max} = 0.432$

6047 measured reflections

5502 independent reflections

3935 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.020$

$\theta_{\max} = 74.9$ °, $\theta_{\min} = 4.5$ °

$h = -12 \rightarrow 12$

$k = -17 \rightarrow 17$

$l = -12 \rightarrow 12$

1 standard reflections every 60 min

intensity decay: 2%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.051$

$wR(F^2) = 0.140$

$S = 1.02$

5502 reflections

348 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0642P)^2 + 0.5869P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.45$ e Å⁻³

$\Delta\rho_{\min} = -0.38$ e Å⁻³

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0039 (3)

Absolute structure: Flack (1983)

Absolute structure parameter: -0.06 (3)

Special details

Experimental. Number of ψ -scan sets used was 8. The θ correction was applied. Averaged transmission function was used. No Fourier smoothing was applied.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1a	0.0320 (4)	0.0742 (3)	0.5522 (4)	0.0519 (9)
C2a	0.0599 (5)	-0.0181 (3)	0.5310 (6)	0.0558 (11)
O21a	0.1299 (4)	-0.0683 (3)	0.6073 (5)	0.0748 (11)
C3a	-0.0156 (6)	-0.0421 (3)	0.3892 (6)	0.0599 (13)
O31a	-0.0148 (5)	-0.1195 (2)	0.3406 (5)	0.0838 (13)
C4a	-0.0793 (5)	0.0440 (3)	0.3390 (5)	0.0536 (12)
C5a	-0.1571 (6)	0.0641 (4)	0.2187 (6)	0.0655 (14)
H5a	-0.1750	0.0188	0.1517	0.079*
C6a	-0.2075 (6)	0.1520 (4)	0.1997 (6)	0.0683 (15)
Br6a	-0.31241 (9)	0.18369 (7)	0.03306 (8)	0.1113 (3)
C7a	-0.1789 (6)	0.2207 (4)	0.2980 (6)	0.0603 (13)
H7a	-0.2141	0.2804	0.2822	0.072*
C8a	-0.1000 (5)	0.2019 (3)	0.4175 (5)	0.0542 (12)
H8a	-0.0799	0.2482	0.4827	0.065*
C9a	-0.0506 (5)	0.1117 (3)	0.4392 (5)	0.0468 (10)
C10a	0.0842 (5)	0.1204 (3)	0.6760 (5)	0.0524 (11)
H10a	0.1315	0.0724	0.7341	0.063*
C11a	-0.0284 (6)	0.1595 (4)	0.7571 (5)	0.0663 (15)
H11a	-0.0726	0.2112	0.7085	0.099*
H11b	0.0119	0.1801	0.8450	0.099*
H11c	-0.0954	0.1119	0.7687	0.099*
C12a	0.1957 (5)	0.1904 (4)	0.6456 (5)	0.0539 (11)
O12a	0.2353 (4)	0.2019 (3)	0.5372 (4)	0.0787 (12)
O13a	0.2439 (4)	0.2352 (3)	0.7588 (4)	0.0654 (10)
C14a	0.3501 (6)	0.3044 (5)	0.7423 (6)	0.0764 (17)
H14a	0.4308	0.2749	0.7111	0.092*
H14b	0.3160	0.3506	0.6753	0.092*
C15A	0.3859 (7)	0.3487 (5)	0.8757 (7)	0.089 (2)
H15A	0.3052	0.3772	0.9060	0.133*
H15B	0.4552	0.3954	0.8675	0.133*
H15C	0.4208	0.3026	0.9408	0.133*
N1b	0.5010 (4)	0.5226 (3)	0.4543 (4)	0.0580 (10)
C2b	0.4556 (6)	0.6130 (4)	0.4568 (7)	0.0662 (15)

O21b	0.3847 (5)	0.6536 (3)	0.3690 (6)	0.0947 (15)
C3b	0.5115 (6)	0.6509 (4)	0.5989 (7)	0.0671 (15)
O31b	0.4928 (5)	0.7288 (3)	0.6363 (6)	0.0896 (15)
C4b	0.5872 (5)	0.5728 (4)	0.6681 (5)	0.0567 (12)
C5b	0.6581 (6)	0.5656 (5)	0.7945 (6)	0.0692 (15)
H5b	0.6639	0.6155	0.8551	0.083*
C6b	0.7207 (5)	0.4817 (5)	0.8295 (5)	0.0689 (14)
Br6b	0.82477 (10)	0.46814 (8)	1.00249 (8)	0.1231 (4)
C7b	0.7096 (6)	0.4070 (4)	0.7425 (6)	0.0671 (14)
H7b	0.7517	0.3510	0.7701	0.080*
C8b	0.6371 (5)	0.4131 (4)	0.6151 (5)	0.0553 (12)
H8b	0.6295	0.3623	0.5560	0.066*
C9b	0.5761 (5)	0.4980 (3)	0.5786 (5)	0.0459 (10)
C10b	0.4649 (5)	0.4574 (4)	0.3427 (5)	0.0596 (12)
H10b	0.5158	0.3999	0.3665	0.072*
C11b	0.5102 (7)	0.4896 (7)	0.2087 (7)	0.106 (3)
H11d	0.4538	0.5411	0.1747	0.159*
H11e	0.5008	0.4394	0.1440	0.159*
H11f	0.6050	0.5090	0.2219	0.159*
C12b	0.3120 (5)	0.4334 (4)	0.3396 (5)	0.0559 (12)
O12b	0.2423 (4)	0.4511 (3)	0.4281 (4)	0.0780 (11)
O13b	0.2720 (4)	0.3850 (3)	0.2285 (4)	0.0750 (11)
C14b	0.1338 (7)	0.3446 (7)	0.2229 (9)	0.105 (3)
H14c	0.1250	0.3089	0.3054	0.126*
H14d	0.0654	0.3940	0.2178	0.126*
C15b	0.1098 (8)	0.2856 (6)	0.1066 (9)	0.114 (3)
H15d	0.1309	0.3188	0.0265	0.171*
H15e	0.0147	0.2668	0.0959	0.171*
H15f	0.1676	0.2315	0.1189	0.171*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1a	0.059 (2)	0.0364 (19)	0.059 (2)	0.0009 (17)	-0.0007 (18)	-0.0023 (17)
C2a	0.052 (3)	0.038 (2)	0.079 (3)	-0.001 (2)	0.015 (2)	0.002 (2)
O21a	0.074 (2)	0.045 (2)	0.104 (3)	0.0080 (18)	0.006 (2)	0.008 (2)
C3a	0.065 (3)	0.041 (3)	0.075 (3)	-0.006 (2)	0.016 (3)	-0.012 (3)
O31a	0.094 (3)	0.0413 (19)	0.116 (4)	-0.0057 (19)	0.011 (3)	-0.027 (2)
C4a	0.062 (3)	0.040 (3)	0.059 (3)	-0.008 (2)	0.008 (2)	-0.008 (2)
C5a	0.076 (4)	0.056 (3)	0.064 (3)	-0.021 (3)	0.003 (3)	-0.009 (3)
C6a	0.065 (3)	0.073 (4)	0.065 (3)	-0.015 (3)	-0.003 (3)	0.000 (3)
Br6A	0.1135 (6)	0.1265 (7)	0.0846 (5)	-0.0145 (5)	-0.0355 (4)	0.0150 (5)
C7a	0.059 (3)	0.048 (3)	0.074 (3)	-0.002 (2)	0.005 (3)	0.008 (2)
C8a	0.059 (3)	0.035 (2)	0.067 (3)	-0.004 (2)	-0.002 (2)	-0.004 (2)
C9a	0.047 (2)	0.040 (2)	0.053 (2)	-0.0066 (19)	0.005 (2)	-0.005 (2)
C10a	0.059 (3)	0.042 (2)	0.054 (3)	-0.003 (2)	0.000 (2)	0.006 (2)
C11a	0.069 (3)	0.075 (4)	0.057 (3)	-0.014 (3)	0.017 (3)	-0.006 (3)
C12a	0.048 (2)	0.053 (3)	0.062 (3)	-0.002 (2)	0.009 (2)	-0.003 (3)

O12a	0.074 (2)	0.100 (3)	0.066 (2)	-0.025 (2)	0.0229 (19)	-0.011 (2)
O13a	0.062 (2)	0.073 (2)	0.061 (2)	-0.0174 (18)	0.0030 (17)	-0.0111 (18)
C14a	0.061 (3)	0.091 (4)	0.078 (4)	-0.031 (3)	0.013 (3)	-0.021 (3)
C15a	0.071 (4)	0.102 (5)	0.092 (5)	-0.023 (4)	0.006 (3)	-0.029 (4)
N1b	0.058 (2)	0.051 (2)	0.063 (3)	0.000 (2)	-0.003 (2)	-0.0029 (19)
C2b	0.055 (3)	0.050 (3)	0.093 (4)	0.003 (2)	0.004 (3)	0.010 (3)
O21b	0.077 (3)	0.078 (3)	0.125 (4)	0.009 (2)	-0.011 (3)	0.035 (3)
C3b	0.057 (3)	0.048 (3)	0.099 (4)	-0.006 (2)	0.018 (3)	-0.002 (3)
O31b	0.085 (3)	0.044 (2)	0.142 (4)	-0.002 (2)	0.022 (3)	-0.016 (2)
C4b	0.054 (3)	0.051 (3)	0.066 (3)	-0.007 (2)	0.009 (2)	-0.012 (2)
C5b	0.060 (3)	0.076 (4)	0.071 (3)	-0.011 (3)	0.003 (3)	-0.027 (3)
C6b	0.056 (3)	0.093 (4)	0.057 (3)	0.004 (3)	-0.001 (2)	-0.006 (3)
Br6b	0.1132 (6)	0.1801 (10)	0.0686 (4)	0.0217 (6)	-0.0264 (4)	-0.0147 (6)
C7b	0.058 (3)	0.077 (4)	0.066 (3)	0.010 (3)	0.005 (3)	0.003 (3)
C8b	0.056 (3)	0.055 (3)	0.055 (3)	0.000 (2)	0.005 (2)	-0.003 (2)
C9b	0.042 (2)	0.045 (2)	0.051 (2)	-0.0056 (19)	0.0017 (18)	-0.0036 (19)
C10b	0.052 (3)	0.071 (3)	0.054 (2)	-0.002 (3)	-0.003 (2)	-0.006 (3)
C11b	0.088 (5)	0.164 (8)	0.069 (4)	-0.043 (5)	0.022 (3)	-0.027 (5)
C12b	0.058 (3)	0.059 (3)	0.051 (3)	0.000 (2)	0.003 (2)	-0.001 (2)
O12b	0.072 (2)	0.084 (3)	0.082 (3)	-0.013 (2)	0.027 (2)	-0.018 (2)
O13b	0.054 (2)	0.099 (3)	0.072 (2)	-0.013 (2)	0.0042 (18)	-0.018 (2)
C14b	0.061 (4)	0.143 (7)	0.112 (6)	-0.028 (4)	0.016 (4)	-0.037 (5)
C15b	0.086 (5)	0.117 (6)	0.140 (7)	-0.035 (5)	0.017 (5)	-0.049 (6)

Geometric parameters (Å, °)

N1a—C2a	1.372 (6)	N1b—C2b	1.373 (7)
N1a—C9a	1.412 (6)	N1b—C9b	1.405 (6)
N1a—C10a	1.434 (6)	N1b—C10b	1.457 (6)
C2a—O21a	1.202 (6)	C2b—O21b	1.201 (7)
C2a—C3a	1.548 (7)	C2b—C3b	1.547 (9)
C3a—O31a	1.209 (6)	C3b—O31b	1.197 (6)
C3a—C4a	1.447 (7)	C3b—C4b	1.471 (8)
C4a—C5a	1.370 (7)	C4b—C5b	1.364 (8)
C4a—C9a	1.392 (6)	C4b—C9b	1.386 (7)
C5a—C6a	1.361 (8)	C5b—C6b	1.378 (9)
C5a—H5a	0.9300	C5b—H5b	0.9300
C6a—C7a	1.390 (8)	C6b—C7b	1.369 (8)
C6a—Br6a	1.897 (6)	C6b—Br6b	1.901 (5)
C7a—C8a	1.365 (7)	C7b—C8b	1.378 (7)
C7a—H7a	0.9300	C7b—H7b	0.9300
C8a—C9a	1.391 (7)	C8b—C9b	1.388 (7)
C8a—H8a	0.9300	C8b—H8b	0.9300
C10a—C11a	1.526 (7)	C10b—C11b	1.504 (8)
C10a—C12a	1.531 (7)	C10b—C12b	1.526 (7)
C10a—H10a	0.9800	C10b—H10b	0.9800
C11a—H11a	0.9600	C11b—H11d	0.9600
C11a—H11b	0.9600	C11b—H11e	0.9600

C11a—H11c	0.9600	C11b—H11f	0.9600
C12a—O12a	1.182 (6)	C12b—O12b	1.185 (6)
C12a—O13a	1.329 (6)	C12b—O13b	1.321 (6)
O13a—C14a	1.455 (6)	O13b—C14b	1.462 (7)
C14a—C15a	1.469 (8)	C14b—C15b	1.424 (10)
C14a—H14a	0.9700	C14b—H14c	0.9700
C14a—H14b	0.9700	C14b—H14d	0.9700
C15a—H15a	0.9600	C15b—H15d	0.9600
C15a—H15b	0.9600	C15b—H15e	0.9600
C15a—H15c	0.9600	C15b—H15f	0.9600
C2a—N1a—C9a	110.7 (4)	C2b—N1b—C9b	111.2 (4)
C2a—N1a—C10a	121.2 (4)	C2b—N1b—C10b	124.5 (5)
C9a—N1a—C10a	128.1 (4)	C9b—N1b—C10b	124.0 (4)
O21a—C2a—N1a	126.3 (5)	O21b—C2b—N1b	127.5 (6)
O21a—C2a—C3a	128.1 (5)	O21b—C2b—C3b	127.1 (6)
N1a—C2a—C3a	105.6 (4)	N1b—C2b—C3b	105.3 (5)
O31a—C3a—C4a	132.0 (5)	O31b—C3b—C4b	130.9 (7)
O31a—C3a—C2a	122.6 (5)	O31b—C3b—C2b	123.8 (6)
C4a—C3a—C2a	105.4 (4)	C4b—C3b—C2b	105.3 (5)
C5a—C4a—C9a	121.2 (5)	C5b—C4b—C9b	121.4 (5)
C5a—C4a—C3a	131.0 (5)	C5b—C4b—C3b	131.6 (5)
C9a—C4a—C3a	107.8 (4)	C9b—C4b—C3b	107.0 (5)
C4a—C5a—C6a	118.4 (5)	C4b—C5b—C6b	117.7 (5)
C4a—C5a—H5a	120.8	C4b—C5b—H5b	121.2
C6a—C5a—H5a	120.8	C6b—C5b—H5b	121.2
C5a—C6a—C7a	121.2 (5)	C5b—C6b—C7b	121.6 (5)
C5a—C6a—Br6a	119.8 (4)	C5b—C6b—Br6b	119.7 (5)
C7a—C6a—Br6a	118.9 (4)	C7b—C6b—Br6b	118.7 (5)
C8a—C7a—C6a	121.0 (5)	C8b—C7b—C6b	121.3 (6)
C8a—C7a—H7a	119.5	C8b—C7b—H7b	119.3
C6a—C7a—H7a	119.5	C6b—C7b—H7b	119.3
C7a—C8a—C9a	118.2 (4)	C7b—C8b—C9b	117.2 (5)
C7a—C8a—H8a	120.9	C7b—C8b—H8b	121.4
C9a—C8a—H8a	120.9	C9b—C8b—H8b	121.4
C8a—C9a—C4a	120.0 (4)	C8b—C9b—C4b	120.8 (4)
C8a—C9a—N1a	129.6 (4)	C8b—C9b—N1b	128.1 (4)
C4a—C9a—N1a	110.4 (4)	C4b—C9b—N1b	111.1 (4)
N1a—C10a—C11a	113.7 (4)	N1b—C10b—C11b	113.1 (5)
N1a—C10a—C12a	109.6 (4)	N1b—C10b—C12b	108.7 (4)
C11a—C10a—C12a	115.0 (4)	C11b—C10b—C12b	115.1 (5)
N1a—C10a—H10a	105.9	N1b—C10b—H10b	106.4
C11a—C10a—H10a	105.9	C11b—C10b—H10b	106.4
C12a—C10a—H10a	105.9	C12b—C10b—H10b	106.4
C10a—C11a—H11a	109.5	C10b—C11b—H11d	109.5
C10a—C11a—H11b	109.5	C10b—C11b—H11e	109.5
H11a—C11a—H11b	109.5	H11d—C11b—H11e	109.5
C10a—C11a—H11c	109.5	C10b—C11b—H11f	109.5

H11a—C11a—H11c	109.5	H11d—C11b—H11f	109.5
H11b—C11a—H11c	109.5	H11e—C11b—H11f	109.5
O12a—C12a—O13a	124.6 (5)	O12b—C12b—O13b	125.3 (5)
O12a—C12a—C10a	124.8 (5)	O12b—C12b—C10b	124.6 (5)
O13a—C12a—C10a	110.6 (4)	O13b—C12b—C10b	110.0 (5)
C12a—O13a—C14a	115.6 (4)	C12b—O13b—C14b	115.7 (5)
C15a—C14a—O13a	107.7 (5)	C15b—C14b—O13b	110.0 (6)
C15a—C14a—H14a	110.2	C15b—C14b—H14c	109.7
O13a—C14a—H14a	110.2	O13b—C14b—H14c	109.7
C15a—C14a—H14b	110.2	C15b—C14b—H14d	109.7
O13a—C14a—H14b	110.2	O13b—C14b—H14d	109.7
H14a—C14a—H14b	108.5	H14c—C14b—H14d	108.2
C14a—C15a—H15a	109.5	C14b—C15b—H15d	109.5
C14a—C15a—H15b	109.5	C14b—C15b—H15e	109.5
H15a—C15a—H15b	109.5	H15d—C15b—H15e	109.5
C14a—C15a—H15c	109.5	C14b—C15b—H15f	109.5
H15a—C15a—H15c	109.5	H15d—C15b—H15f	109.5
H15b—C15a—H15c	109.5	H15e—C15b—H15f	109.5
C9a—N1a—C2a—O21a	178.3 (5)	C9b—N1b—C2b—O21b	-176.9 (6)
C10a—N1a—C2a—O21a	-1.7 (8)	C10b—N1b—C2b—O21b	-2.6 (9)
C9a—N1a—C2a—C3a	-0.9 (5)	C9b—N1b—C2b—C3b	1.4 (6)
C10a—N1a—C2a—C3a	179.0 (4)	C10b—N1b—C2b—C3b	175.7 (4)
O21a—C2a—C3a—O31a	2.6 (9)	O21b—C2b—C3b—O31b	-2.5 (10)
N1a—C2a—C3a—O31a	-178.2 (5)	N1b—C2b—C3b—O31b	179.2 (6)
O21a—C2a—C3a—C4a	-177.6 (5)	O21b—C2b—C3b—C4b	177.8 (6)
N1a—C2a—C3a—C4a	1.7 (5)	N1b—C2b—C3b—C4b	-0.5 (6)
O31a—C3a—C4a—C5a	-1.2 (11)	O31b—C3b—C4b—C5b	0.0 (11)
C2a—C3a—C4a—C5a	179.0 (6)	C2b—C3b—C4b—C5b	179.7 (6)
O31a—C3a—C4a—C9a	178.0 (6)	O31b—C3b—C4b—C9b	179.8 (6)
C2a—C3a—C4a—C9a	-1.8 (6)	C2b—C3b—C4b—C9b	-0.5 (6)
C9a—C4a—C5a—C6a	-0.7 (8)	C9b—C4b—C5b—C6b	1.3 (9)
C3a—C4a—C5a—C6a	178.4 (6)	C3b—C4b—C5b—C6b	-179.0 (6)
C4a—C5a—C6a—C7a	1.3 (9)	C4b—C5b—C6b—C7b	-1.9 (9)
C4a—C5a—C6a—Br6a	178.7 (4)	C4b—C5b—C6b—Br6b	178.5 (4)
C5a—C6a—C7a—C8a	-0.5 (9)	C5b—C6b—C7b—C8b	1.3 (9)
Br6a—C6a—C7a—C8a	-177.8 (4)	Br6b—C6b—C7b—C8b	-179.1 (4)
C6a—C7a—C8a—C9a	-1.0 (8)	C6b—C7b—C8b—C9b	0.0 (8)
C7a—C8a—C9a—C4a	1.5 (7)	C7b—C8b—C9b—C4b	-0.6 (8)
C7a—C8a—C9a—N1a	180.0 (5)	C7b—C8b—C9b—N1b	178.0 (5)
C5a—C4a—C9a—C8a	-0.7 (8)	C5b—C4b—C9b—C8b	-0.1 (8)
C3a—C4a—C9a—C8a	-180.0 (5)	C3b—C4b—C9b—C8b	-179.8 (5)
C5a—C4a—C9a—N1a	-179.4 (5)	C5b—C4b—C9b—N1b	-178.8 (5)
C3a—C4a—C9a—N1a	1.3 (6)	C3b—C4b—C9b—N1b	1.4 (6)
C2a—N1a—C9a—C8a	-178.8 (5)	C2b—N1b—C9b—C8b	179.5 (5)
C10a—N1a—C9a—C8a	1.3 (8)	C10b—N1b—C9b—C8b	5.1 (8)
C2a—N1a—C9a—C4a	-0.2 (6)	C2b—N1b—C9b—C4b	-1.8 (6)
C10a—N1a—C9a—C4a	179.9 (5)	C10b—N1b—C9b—C4b	-176.2 (5)

C2a—N1a—C10a—C11a	-119.8 (5)	C2b—N1b—C10b—C11b	60.4 (7)
C9a—N1a—C10a—C11a	60.1 (6)	C9b—N1b—C10b—C11b	-125.9 (6)
C2a—N1a—C10a—C12a	109.9 (5)	C2b—N1b—C10b—C12b	-68.7 (6)
C9a—N1a—C10a—C12a	-70.1 (6)	C9b—N1b—C10b—C12b	104.9 (5)
N1a—C10a—C12a—O12a	-2.3 (7)	N1b—C10b—C12b—O12b	-12.8 (8)
C11a—C10a—C12a—O12a	-131.8 (6)	C11b—C10b—C12b—O12b	-140.8 (7)
N1a—C10a—C12a—O13a	179.0 (4)	N1b—C10b—C12b—O13b	171.3 (4)
C11A—C10a—C12a—O13a	49.5 (6)	C11b—C10b—C12b—O13b	43.3 (7)
O12A—C12a—O13a—C14a	1.8 (8)	O12b—C12b—O13b—C14b	-5.3 (9)
C10A—C12a—O13a—C14a	-179.5 (5)	C10b—C12b—O13b—C14b	170.6 (6)
C12A—O13a—C14a—C15a	177.7 (5)	C12b—O13b—C14b—C15b	-173.2 (7)
