

catena-Poly[[*(nitrate-κ²O,O')*silver(I)]-*μ*-1,2-bis(diphenylphosphino)ethane-κ²P:P']

Xing-Cong Wang, Yan-Li Wu and Xiu-Li You*

 Jiangxi Key Laboratory of Organic Chemistry, Jiangxi Science and Technology Normal University, Nanchang 330013, People's Republic of China
 Correspondence e-mail: jxstnu116@yahoo.cn


Received 25 June 2008; accepted 26 June 2008

 Key indicators: single-crystal X-ray study; $T = 223$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.036; wR factor = 0.088; data-to-parameter ratio = 15.4.

In the title chain compound, $[\text{Ag}(\text{NO}_3)(\text{C}_{26}\text{H}_{24}\text{P}_2)]_n$, the bis(diphenylphosphino)ethane (dppe) units link the Ag^+ ions into chains along [001]. A nitrate anion is coordinated to the Ag atom. There is a centre of symmetry at the mid-point of the ethane C—C bond and a twofold rotation axis passes through the Ag, N and terminal O atoms. Each Ag atom is four-coordinated in a distorted tetrahedral geometry by two O atoms of the nitrate anion and two P atoms of dppe ligands. The two aromatic rings are oriented at a dihedral angle of $73.77(3)^\circ$.

Related literature

For related literature, see: Harker & Tiekink (1990); Huang *et al.* (1991); Menezes Vicenti & Burrow (2007); Yang *et al.* (1992).



Experimental

Crystal data

$[\text{Ag}(\text{NO}_3)(\text{C}_{26}\text{H}_{24}\text{P}_2)]$	$V = 2542.0(9) \text{ \AA}^3$
$M_r = 568.27$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 17.123(3) \text{ \AA}$	$\mu = 0.95 \text{ mm}^{-1}$
$b = 14.064(3) \text{ \AA}$	$T = 223.2 \text{ K}$
$c = 11.120(2) \text{ \AA}$	$0.30 \times 0.26 \times 0.20 \text{ mm}$
$\beta = 108.33(3)^\circ$	

Data collection

Rigaku Mercury diffractometer	12170 measured reflections
Absorption correction: multi-scan (Jacobson, 1998)	2327 independent reflections
$T_{\min} = 0.704$, $T_{\max} = 0.833$	2161 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	151 parameters
$wR(F^2) = 0.087$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.97 \text{ e \AA}^{-3}$
2327 reflections	$\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$

Table 1

Selected geometric parameters (Å, °).

Ag1—P1	2.4066 (9)	Ag1—O1	2.508 (2)
P1—Ag1—P1 ⁱ	137.49 (4)	P1—Ag1—O1	102.63 (6)
P1—Ag1—O1 ⁱ	115.92 (7)	O1 ⁱ —Ag1—O1	50.52 (11)

 Symmetry code: (i) $-x, y, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku/MSC, 2001); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku/MSC, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97*; software used to prepare material for publication: *SHELXL97*.

The authors acknowledge Jiangxi Science and Technology Normal University for funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2480).

References

- Harker, C. S. W. & Tiekink, E. R. T. (1990). *J. Coord. Chem.* **21**, 287–293.
 Huang, M. S., Zhang, P., Zhang, Y., Yang, H. H. & Zheng, L. S. (1991). *Acta Phys. Chim. Sinica*, **7**, 694–698.
 Jacobson, R. (1998). Private communication to the Rigaku Corporation, Tokyo, Japan.
 Menezes Vicenti, J. R. de & Burrow, R. A. (2007). *Acta Cryst.* **C63**, m88–m90.
 Rigaku/MSC (2001). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
 Rigaku/MSC (2004). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yang, H. H., Zheng, L. S., Xu, Y. J. & Zhang, Q. E. (1992). *Chin. J. Inorg. Chem.* **8**, 65–67.

supporting information

Acta Cryst. (2008). E64, m981 [doi:10.1107/S1600536808019442]

catena-Poly[[nitrate- κ^2O,O' silver(I)]- μ -1,2-bis(diphenylphosphino)ethane- $\kappa^2P:P'$]

Xing-Cong Wang, Yan-Li Wu and Xiu-Li You

S1. Comment

The complexes obtained by the reaction of AgNO_3 with bis(diphenylphosphino)- ethane), dppe, include mono-nuclear complex $\text{Ag}(\text{dppe})(\text{NO}_3)$, (II) (Harker & Tiekink, 1990), binuclear complex $[\text{Ag}(\text{dppe})]_2(\text{NO}_3)_2 \cdot 2\text{MeOH}$, (III) (Yang *et al.*, 1992), and one-dimensional polymers: $[\text{Ag}_4(\text{dppe})_3(\text{NO}_3)_4]_n$, (III) (Huang *et al.*, 1991) and $[\text{Ag}(\text{dppe})(\text{NO}_3)(\text{DMF})]_n$, (IV) (Menezes Vicenti & Burrow, 2007). We report herein the formation of a one-dimensional coordination polymer, (I), using AgNO_3 as the metal source and dppe as bidentate bridging ligand, and its crystal structure.

The structure of the title compound, (I), is polymeric with dppe bridging ligands between Ag centres to form a chain (Fig. 1). There is one dppe ligand in the asymmetric unit. The remaining parts are generated by crystallographic centres of inversion at the mid-points of the C-C bond of the ethane group. The polymeric chains are elongated along [001] direction (Fig. 2). A nitrate anion is coordinated to the Ag atom, in which a twofold rotation axis passes through the N1-O2 bond. Each Ag atom is four-coordinated in a distorted tetrahedral geometry (Table 1) by two O atoms of the nitrate anion and two P atoms of dppe ligands. The two aromatic rings are oriented at a dihedral angle of $73.77(3)^\circ$.

S2. Experimental

For the preparation of the title compound, dppe (20 mg, 0.05 mmol) was dissolved in CH_2Cl_2 (5 ml) and was poured into the tube, then MeOH (3 ml) was layered on it. Finally, MeOH solution (5 ml) containing AgNO_3 (8.5 mg, 0.05 mmol) was layered on the top of the tube. The crystals of the title compound, (I), were obtained for about 3 d.

S3. Refinement

H atoms were positioned geometrically, with C-H = 0.93 and 0.97 Å for aromatic and methylene H and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level [symmetry codes: (i) $-x, 1 - y, -z$; (ii) $-x, y, 1/2 - z$].




Figure 2

The coordination polymer of (I) [symmetry code: (A) $-x, y, 1/2 - z$] along the *c* axis.

catena-Poly[[*(nitrate-κ²O,O')*silver(I)- *μ*-1,2-bis(diphenylphosphino)ethane-κ²P:P']

Crystal data

[Ag(NO₃)(C₂₆H₂₄P₂)]

M_r = 568.27

Monoclinic, *C2/c*

Hall symbol: $-C 2yc$

a = 17.123 (3) Å

b = 14.064 (3) Å

$c = 11.120 (2) \text{ \AA}$
 $\beta = 108.33 (3)^\circ$
 $V = 2542.0 (9) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 1152$
 $D_x = 1.485 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5049 reflections
 $\theta = 3.5\text{--}25.3^\circ$
 $\mu = 0.95 \text{ mm}^{-1}$
 $T = 223 \text{ K}$
 Block, colorless
 $0.30 \times 0.26 \times 0.20 \text{ mm}$

Data collection

Rigaku Mercury
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: $14.6306 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (Jacobson, 1998)
 $T_{\min} = 0.704$, $T_{\max} = 0.833$

12170 measured reflections
 2327 independent reflections
 2161 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 3.5^\circ$
 $h = -20 \rightarrow 19$
 $k = -16 \rightarrow 15$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.087$
 $S = 1.07$
 2327 reflections
 151 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0457P)^2 + 4.2828P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.97 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.45 \text{ e \AA}^{-3}$

Special details

Experimental. no

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag1	0.0000	0.33644 (2)	0.2500	0.03718 (14)
P1	0.08413 (5)	0.39847 (5)	0.13087 (7)	0.02910 (19)
O1	0.06120 (15)	0.17517 (16)	0.3151 (2)	0.0493 (6)
O2	0.0000	0.0421 (3)	0.2500	0.0783 (13)
N1	0.0000	0.1292 (3)	0.2500	0.0409 (9)
C1	0.18506 (17)	0.4287 (2)	0.2384 (3)	0.0342 (7)
C2	0.2341 (2)	0.5014 (3)	0.2188 (3)	0.0486 (8)
H2A	0.2160	0.5381	0.1457	0.058*

C3	0.3100 (2)	0.5200 (3)	0.3075 (4)	0.0635 (11)
H3A	0.3426	0.5693	0.2943	0.076*
C4	0.3370 (2)	0.4648 (3)	0.4153 (4)	0.0633 (11)
H4A	0.3878	0.4771	0.4750	0.076*
C5	0.2902 (2)	0.3933 (3)	0.4345 (4)	0.0612 (11)
H5A	0.3094	0.3560	0.5069	0.073*
C6	0.2142 (2)	0.3747 (3)	0.3484 (3)	0.0472 (8)
H6A	0.1822	0.3257	0.3639	0.057*
C7	0.10284 (19)	0.3224 (2)	0.0110 (3)	0.0351 (7)
C8	0.1568 (2)	0.3472 (3)	-0.0540 (4)	0.0529 (9)
H8A	0.1849	0.4048	-0.0367	0.064*
C9	0.1693 (3)	0.2867 (3)	-0.1445 (4)	0.0690 (12)
H9A	0.2050	0.3040	-0.1888	0.083*
C10	0.1292 (3)	0.2022 (3)	-0.1684 (4)	0.0702 (13)
H10A	0.1385	0.1613	-0.2282	0.084*
C11	0.0753 (3)	0.1761 (3)	-0.1058 (4)	0.0677 (13)
H11A	0.0477	0.1183	-0.1237	0.081*
C12	0.0620 (2)	0.2364 (2)	-0.0154 (3)	0.0490 (9)
H12A	0.0255	0.2189	0.0274	0.059*
C13	0.04348 (17)	0.5080 (2)	0.0442 (3)	0.0339 (6)
H13A	0.0432	0.5582	0.1039	0.041*
H13B	0.0787	0.5278	-0.0046	0.041*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ag1	0.0364 (2)	0.0378 (2)	0.0391 (2)	0.000	0.01439 (15)	0.000
P1	0.0287 (4)	0.0281 (4)	0.0297 (4)	0.0031 (3)	0.0081 (3)	0.0020 (3)
O1	0.0397 (13)	0.0446 (14)	0.0527 (14)	0.0038 (10)	-0.0010 (11)	0.0004 (11)
O2	0.077 (3)	0.038 (2)	0.118 (4)	0.000	0.029 (3)	0.000
N1	0.042 (2)	0.033 (2)	0.049 (2)	0.000	0.0154 (19)	0.000
C1	0.0290 (15)	0.0384 (16)	0.0348 (15)	0.0051 (12)	0.0096 (13)	-0.0052 (13)
C2	0.0403 (19)	0.049 (2)	0.054 (2)	-0.0047 (15)	0.0113 (16)	-0.0005 (16)
C3	0.0357 (19)	0.063 (3)	0.088 (3)	-0.0127 (17)	0.014 (2)	-0.018 (2)
C4	0.0361 (19)	0.082 (3)	0.059 (2)	0.006 (2)	-0.0046 (18)	-0.028 (2)
C5	0.047 (2)	0.083 (3)	0.043 (2)	0.015 (2)	-0.0008 (18)	-0.0020 (19)
C6	0.0372 (18)	0.062 (2)	0.0404 (18)	0.0085 (16)	0.0087 (15)	0.0075 (16)
C7	0.0380 (17)	0.0331 (16)	0.0326 (15)	0.0098 (12)	0.0088 (13)	-0.0003 (12)
C8	0.054 (2)	0.052 (2)	0.060 (2)	-0.0011 (17)	0.0288 (19)	-0.0087 (17)
C9	0.077 (3)	0.081 (3)	0.062 (3)	0.013 (2)	0.040 (2)	-0.015 (2)
C10	0.097 (3)	0.063 (3)	0.049 (2)	0.027 (2)	0.021 (2)	-0.014 (2)
C11	0.107 (4)	0.040 (2)	0.047 (2)	0.000 (2)	0.011 (2)	-0.0088 (17)
C12	0.071 (2)	0.0376 (18)	0.0369 (17)	-0.0022 (16)	0.0145 (17)	0.0004 (14)
C13	0.0335 (16)	0.0294 (15)	0.0366 (15)	0.0031 (12)	0.0080 (13)	0.0042 (12)

Geometric parameters (Å, °)

Ag1—P1	2.4066 (9)	C5—C6	1.376 (5)
Ag1—P1 ⁱ	2.4066 (9)	C5—H5A	0.9300
Ag1—O1 ⁱ	2.508 (2)	C6—H6A	0.9300
Ag1—O1	2.508 (2)	C7—C12	1.381 (5)
P1—C7	1.815 (3)	C7—C8	1.386 (5)
P1—C1	1.816 (3)	C8—C9	1.385 (5)
P1—C13	1.834 (3)	C8—H8A	0.9300
O1—N1	1.250 (3)	C9—C10	1.357 (7)
O2—N1	1.225 (5)	C9—H9A	0.9300
N1—O1 ⁱ	1.250 (3)	C10—C11	1.370 (7)
C1—C2	1.382 (5)	C10—H10A	0.9300
C1—C6	1.393 (4)	C11—C12	1.387 (5)
C2—C3	1.388 (5)	C11—H11A	0.9300
C2—H2A	0.9300	C12—H12A	0.9300
C3—C4	1.381 (6)	C13—C13 ⁱⁱ	1.521 (6)
C3—H3A	0.9300	C13—H13A	0.9700
C4—C5	1.343 (6)	C13—H13B	0.9700
C4—H4A	0.9300		
P1—Ag1—P1 ⁱ	137.49 (4)	C4—C5—H5A	119.6
P1—Ag1—O1 ⁱ	115.92 (7)	C6—C5—H5A	119.6
P1 ⁱ —Ag1—O1 ⁱ	102.63 (7)	C5—C6—C1	120.4 (4)
P1—Ag1—O1	102.63 (6)	C5—C6—H6A	119.8
P1 ⁱ —Ag1—O1	115.92 (7)	C1—C6—H6A	119.8
O1 ⁱ —Ag1—O1	50.52 (11)	C12—C7—C8	119.1 (3)
C7—P1—C1	105.68 (14)	C12—C7—P1	118.6 (3)
C7—P1—C13	103.59 (14)	C8—C7—P1	122.4 (3)
C1—P1—C13	105.93 (14)	C9—C8—C7	120.3 (4)
C7—P1—Ag1	117.71 (11)	C9—C8—H8A	119.8
C1—P1—Ag1	109.46 (10)	C7—C8—H8A	119.8
C13—P1—Ag1	113.56 (10)	C10—C9—C8	119.8 (4)
N1—O1—Ag1	95.88 (19)	C10—C9—H9A	120.1
O2—N1—O1	121.14 (18)	C8—C9—H9A	120.1
O2—N1—O1 ⁱ	121.14 (18)	C9—C10—C11	121.0 (4)
O1—N1—O1 ⁱ	117.7 (4)	C9—C10—H10A	119.5
C2—C1—C6	118.3 (3)	C11—C10—H10A	119.5
C2—C1—P1	124.7 (2)	C10—C11—C12	119.7 (4)
C6—C1—P1	117.0 (3)	C10—C11—H11A	120.2
C1—C2—C3	120.5 (4)	C12—C11—H11A	120.2
C1—C2—H2A	119.8	C7—C12—C11	120.1 (4)
C3—C2—H2A	119.8	C7—C12—H12A	119.9
C4—C3—C2	119.6 (4)	C11—C12—H12A	119.9
C4—C3—H3A	120.2	C13 ⁱⁱ —C13—P1	110.4 (3)
C2—C3—H3A	120.2	C13 ⁱⁱ —C13—H13A	109.6
C5—C4—C3	120.4 (3)	P1—C13—H13A	109.6
C5—C4—H4A	119.8	C13 ⁱⁱ —C13—H13B	109.6

C3—C4—H4A	119.8	P1—C13—H13B	109.6
C4—C5—C6	120.9 (4)	H13A—C13—H13B	108.1

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $-x, -y+1, -z$.