



# 2,2'-Dimethyl-7,7'-(methylenediiimino)di-1,8-naphthyridin-1-ium bis(perchlorate). Erratum

**IUCr Editorial Office**


5 Abbey Square, Chester CH1 2HU, England

This article reports the correction of a paper by Mo *et al.* [*Acta Cryst.* (2008), **E64**, o1702].

After thorough investigation, an article by Mo *et al.* (2008) has been shown to have problems with its data set. The article is therefore withdrawn from the published literature.

**References**

- Mo, J., Liu, J.-H., Pan, Y.-S., Zhang, S.-M. & Du, X.-D. (2008). *Acta Cryst.* **E64**, o1702.



OPEN ACCESS

## 2,2'-Dimethyl-7,7'-(methylenediamino)di-1,8-naphthyridin-1-i um bis(perchlorate)

Juan Mo,\* Jian-Hua Liu, Yu-Shan Pan, Su-Mei Zhang and Xiang-Dang Du

College of Animal Husbandry and Veterinary Studies, Henan Agricultural University, Zhengzhou, Henan Province 450002, People's Republic of China

Correspondence e-mail: mojuan52@126.com


Received 14 June 2008; accepted 31 July 2008

Key indicators: single-crystal X-ray study;  $T = 113\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ; disorder in solvent or counterion;  $R$  factor = 0.060;  $wR$  factor = 0.134; data-to-parameter ratio = 12.6.

In the title salt,  $\text{C}_{19}\text{H}_{20}\text{N}_6^{2+} \cdot 2\text{ClO}_4^-$ , the two planar 1,8-naphthyridine systems are linked by a methylenediamine group with a dihedral angle of  $60.6(1)^\circ$  between the two systems. The crystal structure involves extensive  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonding.

### Related literature

For related literature, see: Baker & Norman (2004); Gavrilova & Bosnich (2004); Nakatani *et al.* (2000, 2001); Stadie *et al.* (2007); Ferrarini *et al.* (1997).



### Experimental

#### Crystal data

$\text{C}_{19}\text{H}_{20}\text{N}_6^{2+} \cdot 2\text{ClO}_4^-$   
 $M_r = 531.31$   
 Orthorhombic,  $Pbca$   
 $a = 8.191(1)\text{ \AA}$   
 $b = 19.325(2)\text{ \AA}$   
 $c = 27.885(2)\text{ \AA}$

#### Data collection

Bruker SMART CCD area-detector diffractometer

$V = 4413.9(5)\text{ \AA}^3$   
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.36\text{ mm}^{-1}$   
 $T = 113(2)\text{ K}$   
 $0.34 \times 0.16 \times 0.14\text{ mm}$

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.908$ ,  $T_{\max} = 0.952$

31220 measured reflections  
3882 independent reflections

3598 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.061$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$   
 $wR(F^2) = 0.134$   
 $S = 1.16$   
 3882 reflections  
 308 parameters  
 16 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.35\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.41\text{ e \AA}^{-3}$

**Table 1**  
 Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 $\cdots$ O6 <sup>i</sup>	0.88 (4)	1.92	2.794 (4)	168
N3—H3A $\cdots$ O3	0.80 (3)	2.23	2.990 (6)	159
N4—H4A $\cdots$ O5 <sup>i</sup>	0.71 (4)	2.56	3.233 (5)	160
N4—H4A $\cdots$ O7 <sup>i</sup>	0.71 (4)	2.52	3.133 (5)	145
N6—H6 $\cdots$ O1	0.84 (3)	2.00	2.838 (8)	178
C1—H1A $\cdots$ O7 <sup>ii</sup>	0.98	2.54	3.501 (3)	167
C1—H1B $\cdots$ O4 <sup>iii</sup>	0.98	2.33	3.078 (4)	132
C4—H4 $\cdots$ O8 <sup>iv</sup>	0.95	2.52	3.392 (7)	152
C10—H10B $\cdots$ O4 <sup>iv</sup>	0.99	2.35	3.082 (5)	130
C13—H13 $\cdots$ O2 <sup>i</sup>	0.95	2.41	3.259 (3)	149
C19—H19B $\cdots$ O5	0.98	2.57	3.351 (6)	136
C19—H19C $\cdots$ O7 <sup>v</sup>	0.98	2.58	3.207 (7)	122
C19—H19C $\cdots$ O8 <sup>v</sup>	0.98	2.57	3.548 (4)	172

Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $-x + \frac{3}{2}, -y, z - \frac{1}{2}$ ; (iii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iv)  $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ ; (v)  $x - 1, y, z$ .

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *XP* in *SHELXTL*.

We thank Henan Agricultural University for the generous support of this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2415).

### References

- Baker, R. S. & Norman, R. E. (2004). *Acta Cryst. E60*, m1761–m1763.
- Bruker (1997). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ferrarini, P. L., Mori, C., Badawneh, M., Manera, C., Martinelli, A., Miceli, M., Ramagnoli, F. & Saccomanni, G. (1997). *J. Heterocycl. Chem.* **34**, 1501–1504.
- Gavrilova, E. L. & Bosnich, B. (2004). *Chem. Rev.* **104**, 349–383.
- Nakatani, K., Sando, S. & Saito, I. (2000). *J. Am. Chem. Soc.* **122**, 2172–2178.
- Nakatani, K., Sando, S. & Saito, I. (2001). *Nat. Biotechnol.* **19**, 51–55.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stadie, N. P., Sanchez-Smith, R. & Groy, T. L. (2007). *Acta Cryst. E63*, m2153–m2154.

# supporting information

*Acta Cryst.* (2008). E64, o1702 [doi:10.1107/S1600536808024616]

## 2,2'-Dimethyl-7,7'-(methylenediimino)di-1,8-naphthyridin-1-i um bis-(perchlorate)

Juan Mo, Jian-Hua Liu, Yu-Shan Pan, Su-Mei Zhang and Xiang-Dang Du

### S1. Comment

1,8-Naphthyridine and its derivatives are used for binding of mismatched guanine or used as versatile ligands which are able to form metal aggregates with monodentates fashion or chelating bidentate fashion(Nakatani *et al.*, 2000; Nakatani *et al.*, 2001; Ferrarini *et al.*, 1997; Gavrilova & Bosnich, 2004; Baker & Norman, 2004; Stadie *et al.*, 2007). We report here a new 1,8-Naphthyridine compound (Fig. 1).


The title compound reveals 1,8-naphthyridine rings are linked by methenediamine with a dihedral angle between two 1,8-naphthyridine rings of 60.6 (1) $^{\circ}$ . Each 1,8-naphthyridine ring is an almost planar in which the ten atoms forming the 1,8-naphthyridine ring have mean deviation of 0.03 $\text{\AA}$  from the least-squares plane calculated using the ten atoms. To balance hydrogen ion charge of two 1,8-naphthyridine rings, there are two perchlorate groups in crystal cell. From the packing diagram (Fig. 2), it seems that the intramolecular N—H $\cdots$ O and C—H $\cdots$ O and hydrogen bonds are effective in the stabilization of the crystal structure.

### S2. Experimental


To the solution of 2-amino-7-methyl-1,8-naphthyridine (3.18 g, 0.02 mol) in mixed solvent of water (28 mL) and ethanol (2 mL), 37% formadehyde solution (0.86 mL, 0.01 mol) was added dropwise at 0°C and the reaction mixture was stirred at room temperature for 24h. The white precipitate formed was filtered, washed several times with water and then with diethyl ether and dried. Yield: 55% (1.81 g). FTIR (KBr)cm $^{-1}$ :  $\nu_{\text{NH}}$  3389, 3266;  $\nu_{\text{CH}}$  3026. Anal. Calc. For C<sub>19</sub>H<sub>18</sub>N<sub>6</sub>: C, 69.07; H, 5.49; N, 25.44. Found: C, 68.86; H, 5.56; N, 25.37. Single crystals of (I) suitable for an X-ray study were obtained by slow evaporation of an aqueous ethanol solution (30% v/v) under the conditions in the presence of perchloric acid at 293 K over a period of one month.

### S3. Refinement

Hydrogen atoms of NH (naphthyridine and amine) were located in a Fourier map and refined freely. All the other hydrogen atoms were generated geometrically (C—H bond lengths of methyl group fixed at 0.98 $\text{\AA}$ , C—H bond lengths of naphthyridine fixed at 0.95  $\text{\AA}$ ) assigned appropriated isotropic thermal parameters,  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . Each perchlorate anion is disordered over two different orientations. The Cl—O distances were restrained to 1.43 (4) $\text{\AA}$ .

**Figure 1**


Molecular structure of the cation of the title compound showing the atom-numbering scheme and displacement ellipsoids drawn at the 40% probability level.

**Figure 2**

Unit-cell packing diagram as viewed down the *c*-direction. Hydrogen bonds are shown as dashed lines.

### **2,2'-Dimethyl-7,7'-(methylenedimino)di-1,8-naphthyridin-1-ium bis(perchlorate)**

#### *Crystal data*



$$M_r = 531.31$$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$$a = 8.1910 (5) \text{ \AA}$$

$$b = 19.3250 (12) \text{ \AA}$$

$c = 27.8850$  (19) Å  
 $V = 4413.9$  (5) Å<sup>3</sup>  
 $Z = 8$   
 $F(000) = 2192$   
 $D_x = 1.599$  Mg m<sup>-3</sup>  
Mo  $K\alpha$  radiation,  $\lambda = 0.71070$  Å

Cell parameters from 7263 reflections  
 $\theta = 2.1\text{--}28.0^\circ$   
 $\mu = 0.36$  mm<sup>-1</sup>  
 $T = 113$  K  
Prism, colorless  
 $0.34 \times 0.16 \times 0.14$  mm

#### Data collection

Bruker SMART CCD area-detector  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 7.31 pixels mm<sup>-1</sup>  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.909$ ,  $T_{\max} = 0.952$

31220 measured reflections  
3882 independent reflections  
3598 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.061$   
 $\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 2.1^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -22 \rightarrow 22$   
 $l = -33 \rightarrow 32$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.060$   
 $wR(F^2) = 0.134$   
 $S = 1.16$   
3882 reflections  
308 parameters  
16 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0498P)^2 + 5.3196P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.003$   
 $\Delta\rho_{\max} = 0.35$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.41$  e Å<sup>-3</sup>

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cl1	0.24246 (10)	0.22574 (4)	0.56855 (3)	0.0360 (2)	
Cl2	1.00163 (9)	0.01682 (3)	0.69459 (3)	0.0323 (2)	
O1	0.1693 (6)	0.1730 (2)	0.59780 (16)	0.0534 (14)	0.747 (5)
O2	0.3894 (5)	0.2014 (2)	0.5483 (2)	0.0758 (17)	0.747 (5)
O3	0.1302 (6)	0.2438 (2)	0.53084 (12)	0.0738 (15)	0.747 (5)
O4	0.2651 (6)	0.28664 (19)	0.59698 (15)	0.0700 (14)	0.747 (5)
O5	0.8509 (4)	0.02894 (17)	0.67091 (14)	0.0664 (12)	0.870 (6)
O6	0.9850 (5)	-0.03196 (14)	0.73264 (10)	0.0657 (13)	0.870 (6)
O7	1.1129 (3)	-0.01123 (15)	0.65972 (10)	0.0490 (10)	0.870 (6)

O8	1.0661 (6)	0.0802 (2)	0.7134 (2)	0.0444 (13)	0.870 (6)
O1'	0.1358 (14)	0.1676 (5)	0.5766 (4)	0.039 (3)	0.253 (5)
O2'	0.3710 (11)	0.2233 (6)	0.6037 (3)	0.062 (4)	0.253 (5)
O3'	0.3135 (14)	0.2178 (6)	0.5220 (3)	0.062 (4)	0.253 (5)
O4'	0.1558 (13)	0.2875 (4)	0.5708 (4)	0.052 (3)	0.253 (5)
O5'	0.8301 (12)	0.0031 (10)	0.7058 (7)	0.053 (6)	0.130 (6)
O6'	1.0870 (19)	-0.0453 (6)	0.7069 (7)	0.048 (6)	0.130 (6)
O7'	1.012 (3)	0.0307 (10)	0.6446 (4)	0.062 (7)	0.130 (6)
O8'	1.056 (3)	0.0738 (10)	0.7228 (8)	0.039 (10)	0.130 (6)
N1	0.1466 (3)	0.16498 (13)	0.27420 (9)	0.0328 (6)	
N2	0.1460 (3)	0.16453 (12)	0.35633 (8)	0.0293 (6)	
N3	0.1396 (3)	0.16601 (15)	0.43819 (10)	0.0353 (6)	
N6	0.3295 (3)	0.04505 (14)	0.58097 (9)	0.0336 (6)	
N5	0.2041 (3)	0.04253 (13)	0.50731 (9)	0.0327 (6)	
N4	0.0922 (4)	0.04231 (16)	0.43264 (11)	0.0417 (7)	
C2	0.1739 (4)	0.19127 (16)	0.23034 (10)	0.0377 (8)	
C3	0.2463 (5)	0.25570 (18)	0.22665 (11)	0.0435 (9)	
H3	0.2634	0.2759	0.1960	0.052*	
C4	0.2935 (4)	0.29061 (16)	0.26746 (11)	0.0390 (8)	
H4	0.3453	0.3344	0.2647	0.047*	
C5	0.2662 (4)	0.26239 (14)	0.31277 (10)	0.0285 (7)	
C6	0.1858 (4)	0.19774 (14)	0.31595 (10)	0.0268 (6)	
C7	0.1866 (4)	0.19636 (15)	0.39716 (10)	0.0288 (7)	
C8	0.2786 (4)	0.25928 (15)	0.39855 (10)	0.0314 (7)	
H8	0.3132	0.2779	0.4284	0.038*	
C9	0.3159 (4)	0.29193 (15)	0.35722 (10)	0.0324 (7)	
H9	0.3748	0.3343	0.3578	0.039*	
C1	0.1243 (6)	0.14783 (18)	0.18868 (12)	0.0518 (10)	
H1A	0.2123	0.1154	0.1808	0.078*	
H1B	0.1026	0.1776	0.1610	0.078*	
H1C	0.0253	0.1219	0.1968	0.078*	
C16	0.4320 (4)	0.02351 (16)	0.61578 (11)	0.0360 (8)	
C17	0.5112 (4)	-0.03943 (17)	0.60916 (12)	0.0402 (8)	
H17	0.5862	-0.0556	0.6326	0.048*	
C18	0.4812 (4)	-0.07835 (17)	0.56878 (12)	0.0398 (8)	
H18	0.5322	-0.1222	0.5653	0.048*	
C14	0.3773 (4)	-0.05443 (15)	0.53293 (12)	0.0357 (7)	
C15	0.3015 (4)	0.01043 (16)	0.53928 (11)	0.0316 (7)	
C11	0.1802 (4)	0.00934 (17)	0.46592 (11)	0.0368 (8)	
C12	0.2463 (4)	-0.05808 (16)	0.45608 (12)	0.0424 (8)	
H12	0.2221	-0.0808	0.4267	0.051*	
C13	0.3428 (4)	-0.08878 (16)	0.48880 (12)	0.0411 (8)	
H13	0.3878	-0.1332	0.4825	0.049*	
C19	0.4554 (5)	0.06921 (19)	0.65809 (11)	0.0446 (8)	
H19A	0.4863	0.1156	0.6473	0.067*	
H19B	0.5418	0.0503	0.6785	0.067*	
H19C	0.3533	0.0718	0.6763	0.067*	
C10	0.0224 (4)	0.10976 (18)	0.43970 (12)	0.0431 (8)	

H10A	-0.0336	0.1106	0.4712	0.052*
H10B	-0.0612	0.1178	0.4147	0.052*
H3A	0.163 (4)	0.1865 (17)	0.4619 (12)	0.035 (10)*
H1	0.101 (5)	0.124 (2)	0.2764 (13)	0.055 (11)*
H4A	0.081 (5)	0.0269 (19)	0.4097 (13)	0.040 (12)*
H6	0.281 (4)	0.0828 (18)	0.5853 (12)	0.035 (9)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0433 (5)	0.0317 (4)	0.0331 (4)	-0.0003 (3)	0.0046 (4)	-0.0008 (3)
Cl2	0.0360 (5)	0.0256 (4)	0.0353 (4)	-0.0007 (3)	0.0018 (3)	-0.0012 (3)
O1	0.064 (3)	0.046 (2)	0.050 (3)	0.016 (2)	0.026 (2)	0.022 (2)
O2	0.055 (3)	0.049 (2)	0.124 (4)	-0.002 (2)	0.053 (3)	-0.011 (3)
O3	0.118 (4)	0.059 (2)	0.045 (2)	0.031 (2)	-0.025 (2)	0.0033 (18)
O4	0.089 (3)	0.052 (2)	0.070 (3)	-0.001 (2)	0.002 (3)	-0.027 (2)
O5	0.0421 (19)	0.063 (2)	0.095 (3)	0.0139 (15)	-0.0291 (19)	-0.019 (2)
O6	0.126 (4)	0.0396 (16)	0.0313 (15)	-0.0408 (19)	-0.0042 (19)	0.0050 (12)
O7	0.0410 (17)	0.0542 (18)	0.0519 (18)	-0.0075 (14)	0.0127 (14)	-0.0213 (14)
O8	0.055 (3)	0.0232 (18)	0.055 (2)	-0.0038 (16)	-0.002 (2)	-0.0076 (17)
O1'	0.036 (6)	0.030 (5)	0.051 (7)	-0.003 (4)	0.014 (5)	0.002 (5)
O2'	0.049 (6)	0.086 (7)	0.052 (6)	-0.022 (5)	-0.013 (5)	0.031 (5)
O3'	0.056 (7)	0.087 (7)	0.041 (6)	-0.016 (6)	0.023 (5)	-0.020 (5)
O4'	0.063 (6)	0.027 (4)	0.065 (6)	0.017 (4)	0.012 (5)	0.007 (4)
O5'	0.030 (8)	0.067 (9)	0.061 (10)	0.008 (7)	0.004 (7)	0.003 (8)
O6'	0.046 (9)	0.033 (8)	0.064 (10)	0.009 (7)	0.000 (8)	0.003 (7)
O7'	0.085 (11)	0.060 (10)	0.041 (9)	-0.012 (8)	0.014 (8)	-0.003 (7)
O8'	0.053 (14)	0.032 (13)	0.032 (12)	0.002 (8)	-0.006 (8)	-0.010 (8)
N1	0.0448 (17)	0.0233 (13)	0.0302 (14)	-0.0046 (12)	-0.0070 (12)	0.0000 (10)
N2	0.0321 (14)	0.0275 (12)	0.0283 (13)	-0.0027 (11)	-0.0009 (11)	0.0035 (10)
N3	0.0381 (16)	0.0419 (15)	0.0260 (14)	0.0023 (13)	-0.0004 (12)	0.0032 (12)
N6	0.0353 (15)	0.0344 (15)	0.0310 (14)	0.0059 (13)	0.0054 (12)	0.0103 (12)
N5	0.0314 (14)	0.0360 (14)	0.0306 (13)	-0.0023 (11)	0.0053 (11)	0.0101 (11)
N4	0.0479 (19)	0.0467 (18)	0.0306 (16)	-0.0128 (14)	-0.0024 (15)	0.0084 (14)
C2	0.053 (2)	0.0331 (16)	0.0266 (16)	0.0011 (15)	-0.0066 (15)	-0.0008 (13)
C3	0.064 (2)	0.0381 (18)	0.0287 (16)	-0.0036 (17)	-0.0008 (16)	0.0063 (14)
C4	0.050 (2)	0.0317 (16)	0.0349 (17)	-0.0098 (15)	0.0017 (15)	0.0043 (13)
C5	0.0341 (17)	0.0212 (14)	0.0302 (15)	-0.0031 (12)	-0.0020 (13)	-0.0004 (12)
C6	0.0325 (16)	0.0223 (13)	0.0257 (14)	-0.0012 (12)	-0.0021 (13)	-0.0003 (11)
C7	0.0299 (16)	0.0301 (15)	0.0264 (15)	0.0067 (13)	0.0012 (13)	0.0022 (12)
C8	0.0359 (18)	0.0293 (15)	0.0290 (15)	0.0050 (13)	-0.0035 (13)	-0.0066 (12)
C9	0.0374 (18)	0.0262 (15)	0.0335 (16)	-0.0018 (13)	-0.0029 (14)	-0.0030 (12)
C1	0.083 (3)	0.0408 (19)	0.0316 (18)	-0.0031 (19)	-0.0128 (19)	-0.0052 (15)
C16	0.0290 (17)	0.0429 (18)	0.0362 (17)	-0.0019 (14)	0.0067 (14)	0.0172 (14)
C17	0.0313 (18)	0.0449 (19)	0.0444 (19)	0.0015 (15)	0.0056 (15)	0.0222 (16)
C18	0.0337 (18)	0.0324 (17)	0.053 (2)	0.0026 (14)	0.0174 (16)	0.0202 (15)
C14	0.0355 (18)	0.0285 (15)	0.0430 (18)	-0.0041 (14)	0.0150 (15)	0.0115 (14)
C15	0.0287 (17)	0.0350 (16)	0.0309 (16)	-0.0028 (13)	0.0097 (13)	0.0104 (13)

C11	0.0381 (19)	0.0409 (18)	0.0314 (16)	-0.0138 (15)	0.0053 (15)	0.0099 (14)
C12	0.052 (2)	0.0331 (17)	0.0417 (19)	-0.0172 (16)	0.0116 (17)	0.0006 (15)
C13	0.048 (2)	0.0273 (16)	0.048 (2)	-0.0080 (15)	0.0165 (17)	0.0061 (14)
C19	0.042 (2)	0.058 (2)	0.0341 (17)	-0.0008 (17)	0.0011 (16)	0.0121 (16)
C10	0.0343 (19)	0.061 (2)	0.0343 (18)	-0.0038 (17)	0.0030 (15)	0.0176 (16)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C11—O4'	1.390 (7)	C2—C3	1.383 (5)
C11—O2	1.410 (4)	C2—C1	1.490 (4)
C11—O3'	1.430 (7)	C3—C4	1.378 (4)
C11—O4	1.431 (3)	C3—H3	0.9500
C11—O1	1.436 (4)	C4—C5	1.394 (4)
C11—O2'	1.439 (7)	C4—H4	0.9500
C11—O3	1.440 (3)	C5—C6	1.415 (4)
C11—O1'	1.441 (8)	C5—C9	1.424 (4)
C12—O5	1.420 (3)	C7—C8	1.431 (4)
C12—O7'	1.423 (9)	C8—C9	1.349 (4)
C12—O8'	1.424 (10)	C8—H8	0.9500
C12—O6	1.426 (3)	C9—H9	0.9500
C12—O6'	1.432 (9)	C1—H1A	0.9800
C12—O8	1.433 (3)	C1—H1B	0.9800
C12—O7	1.439 (3)	C1—H1C	0.9800
C12—O5'	1.464 (9)	C16—C17	1.390 (5)
N1—C2	1.343 (4)	C16—C19	1.486 (5)
N1—C6	1.364 (4)	C17—C18	1.376 (5)
N1—H1	0.88 (4)	C17—H17	0.9500
N2—C7	1.336 (4)	C18—C14	1.392 (5)
N2—C6	1.336 (4)	C18—H18	0.9500
N3—C7	1.342 (4)	C14—C15	1.410 (4)
N3—C10	1.451 (4)	C14—C13	1.427 (5)
N3—H3A	0.80 (3)	C11—C12	1.437 (5)
N6—C16	1.350 (4)	C12—C13	1.345 (5)
N6—C15	1.360 (4)	C12—H12	0.9500
N6—H6	0.84 (3)	C13—H13	0.9500
N5—C11	1.335 (4)	C19—H19A	0.9800
N5—C15	1.348 (4)	C19—H19B	0.9800
N4—C11	1.337 (4)	C19—H19C	0.9800
N4—C10	1.437 (5)	C10—H10A	0.9900
N4—H4A	0.71 (4)	C10—H10B	0.9900
O4'—Cl1—O2	137.8 (4)	N1—C2—C3	118.7 (3)
O4'—Cl1—O3'	109.9 (6)	N1—C2—C1	116.8 (3)
O4'—Cl1—O4	48.3 (4)	C3—C2—C1	124.5 (3)
O2—Cl1—O4	112.7 (3)	C4—C3—C2	120.0 (3)
O3'—Cl1—O4	122.6 (5)	C4—C3—H3	120.0
O4'—Cl1—O1	111.8 (5)	C2—C3—H3	120.0
O2—Cl1—O1	110.3 (3)	C3—C4—C5	120.8 (3)

O3'—Cl1—O1	127.5 (5)	C3—C4—H4	119.6
O4—Cl1—O1	108.9 (3)	C5—C4—H4	119.6
O4'—Cl1—O2'	111.7 (6)	C4—C5—C6	118.5 (3)
O2—Cl1—O2'	68.8 (5)	C4—C5—C9	125.9 (3)
O3'—Cl1—O2'	108.5 (6)	C6—C5—C9	115.6 (3)
O4—Cl1—O2'	63.6 (5)	N2—C6—N1	116.0 (2)
O1—Cl1—O2'	84.0 (4)	N2—C6—C5	126.2 (3)
O4'—Cl1—O3	59.9 (5)	N1—C6—C5	117.8 (3)
O2—Cl1—O3	109.5 (3)	N2—C7—N3	117.0 (3)
O3'—Cl1—O3	67.9 (5)	N2—C7—C8	123.0 (3)
O4—Cl1—O3	106.7 (2)	N3—C7—C8	120.0 (3)
O1—Cl1—O3	108.7 (3)	C9—C8—C7	119.6 (3)
O2'—Cl1—O3	166.5 (4)	C9—C8—H8	120.2
O4'—Cl1—O1'	110.7 (6)	C7—C8—H8	120.2
O2—Cl1—O1'	108.6 (6)	C8—C9—C5	119.4 (3)
O3'—Cl1—O1'	107.7 (6)	C8—C9—H9	120.3
O4—Cl1—O1'	129.3 (5)	C5—C9—H9	120.3
O2'—Cl1—O1'	108.2 (6)	C2—C1—H1A	109.5
O3—Cl1—O1'	85.2 (5)	C2—C1—H1B	109.5
O5—Cl2—O7'	64.3 (8)	H1A—C1—H1B	109.5
O5—Cl2—O8'	113.6 (11)	C2—C1—H1C	109.5
O7'—Cl2—O8'	112.1 (10)	H1A—C1—H1C	109.5
O5—Cl2—O6	111.8 (2)	H1B—C1—H1C	109.5
O7'—Cl2—O6	149.3 (8)	N6—C16—C17	117.7 (3)
O8'—Cl2—O6	97.5 (11)	N6—C16—C19	117.9 (3)
O5—Cl2—O6'	132.4 (7)	C17—C16—C19	124.4 (3)
O7'—Cl2—O6'	111.3 (8)	C18—C17—C16	120.2 (3)
O8'—Cl2—O6'	111.4 (10)	C18—C17—H17	119.9
O6—Cl2—O6'	46.6 (7)	C16—C17—H17	119.9
O5—Cl2—O8	110.5 (2)	C17—C18—C14	121.0 (3)
O7'—Cl2—O8	100.1 (8)	C17—C18—H18	119.5
O6—Cl2—O8	109.1 (3)	C14—C18—H18	119.5
O6'—Cl2—O8	116.7 (8)	C18—C14—C15	118.3 (3)
O5—Cl2—O7	107.38 (19)	C18—C14—C13	125.9 (3)
O7'—Cl2—O7	50.9 (8)	C15—C14—C13	115.8 (3)
O8'—Cl2—O7	117.9 (11)	N5—C15—N6	116.0 (3)
O6—Cl2—O7	108.3 (2)	N5—C15—C14	125.9 (3)
O6'—Cl2—O7	62.4 (8)	N6—C15—C14	118.1 (3)
O8—Cl2—O7	109.6 (3)	N5—C11—N4	116.8 (3)
O5—Cl2—O5'	45.2 (7)	N5—C11—C12	123.1 (3)
O7'—Cl2—O5'	107.7 (8)	N4—C11—C12	120.2 (3)
O8'—Cl2—O5'	108.7 (10)	C13—C12—C11	119.5 (3)
O6—Cl2—O5'	68.2 (8)	C13—C12—H12	120.3
O6'—Cl2—O5'	105.3 (8)	C11—C12—H12	120.3
O8—Cl2—O5'	115.5 (8)	C12—C13—C14	119.8 (3)
O7—Cl2—O5'	133.2 (8)	C12—C13—H13	120.1
C2—N1—C6	124.2 (3)	C14—C13—H13	120.1
C2—N1—H1	118 (2)	C16—C19—H19A	109.5

C6—N1—H1	117 (2)	C16—C19—H19B	109.5
C7—N2—C6	115.9 (2)	H19A—C19—H19B	109.5
C7—N3—C10	122.8 (3)	C16—C19—H19C	109.5
C7—N3—H3A	115 (2)	H19A—C19—H19C	109.5
C10—N3—H3A	121 (2)	H19B—C19—H19C	109.5
C16—N6—C15	124.6 (3)	N4—C10—N3	114.4 (3)
C16—N6—H6	117 (2)	N4—C10—H10A	108.7
C15—N6—H6	118 (2)	N3—C10—H10A	108.7
C11—N5—C15	116.0 (3)	N4—C10—H10B	108.7
C11—N4—C10	123.5 (3)	N3—C10—H10B	108.7
C11—N4—H4A	120 (3)	H10A—C10—H10B	107.6
C10—N4—H4A	117 (3)		
C6—N1—C2—C3	-0.1 (5)	C15—N6—C16—C19	176.9 (3)
C6—N1—C2—C1	179.5 (3)	N6—C16—C17—C18	-1.5 (4)
N1—C2—C3—C4	2.1 (5)	C19—C16—C17—C18	179.8 (3)
C1—C2—C3—C4	-177.4 (4)	C16—C17—C18—C14	2.8 (5)
C2—C3—C4—C5	-1.4 (6)	C17—C18—C14—C15	-0.8 (4)
C3—C4—C5—C6	-1.3 (5)	C17—C18—C14—C13	177.1 (3)
C3—C4—C5—C9	177.0 (3)	C11—N5—C15—N6	178.3 (3)
C7—N2—C6—N1	179.2 (3)	C11—N5—C15—C14	-0.7 (4)
C7—N2—C6—C5	0.2 (4)	C16—N6—C15—N5	-175.3 (3)
C2—N1—C6—N2	178.4 (3)	C16—N6—C15—C14	3.8 (4)
C2—N1—C6—C5	-2.5 (5)	C18—C14—C15—N5	176.7 (3)
C4—C5—C6—N2	-177.9 (3)	C13—C14—C15—N5	-1.5 (4)
C9—C5—C6—N2	3.7 (5)	C18—C14—C15—N6	-2.3 (4)
C4—C5—C6—N1	3.1 (4)	C13—C14—C15—N6	179.5 (3)
C9—C5—C6—N1	-175.3 (3)	C15—N5—C11—N4	-176.0 (3)
C6—N2—C7—N3	176.2 (3)	C15—N5—C11—C12	3.0 (4)
C6—N2—C7—C8	-4.9 (4)	C10—N4—C11—N5	-0.9 (5)
C10—N3—C7—N2	-12.6 (4)	C10—N4—C11—C12	-179.9 (3)
C10—N3—C7—C8	168.5 (3)	N5—C11—C12—C13	-2.9 (5)
N2—C7—C8—C9	5.7 (5)	N4—C11—C12—C13	176.0 (3)
N3—C7—C8—C9	-175.5 (3)	C11—C12—C13—C14	0.5 (5)
C7—C8—C9—C5	-1.4 (5)	C18—C14—C13—C12	-176.5 (3)
C4—C5—C9—C8	178.9 (3)	C15—C14—C13—C12	1.5 (4)
C6—C5—C9—C8	-2.8 (4)	C11—N4—C10—N3	74.2 (4)
C15—N6—C16—C17	-1.9 (4)	C7—N3—C10—N4	85.6 (4)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D\cdots H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N1—H1 $\cdots$ O6 <sup>i</sup>	0.88 (4)	1.92	2.794 (4)	168
N3—H3A $\cdots$ O3	0.80 (3)	2.23	2.990 (6)	159
N4—H4A $\cdots$ O5 <sup>i</sup>	0.71 (4)	2.56	3.233 (5)	160
N4—H4A $\cdots$ O7 <sup>i</sup>	0.71 (4)	2.52	3.133 (5)	145
N6—H6 $\cdots$ O1	0.84 (3)	2.00	2.838 (8)	178
C1—H1A $\cdots$ O7 <sup>ii</sup>	0.98	2.54	3.501 (3)	167

---

C1—H1 <i>B</i> ···O4 <sup>iii</sup>	0.98	2.33	3.078 (4)	132
C4—H4···O8 <sup>iv</sup>	0.95	2.52	3.392 (7)	152
C10—H10 <i>B</i> ···O4 <sup>iv</sup>	0.99	2.35	3.082 (5)	130
C13—H13···O2 <sup>i</sup>	0.95	2.41	3.259 (3)	149
C19—H19 <i>B</i> ···O5	0.98	2.57	3.351 (6)	136
C19—H19 <i>C</i> ···O7 <sup>v</sup>	0.98	2.58	3.207 (7)	122
C19—H19 <i>C</i> ···O8 <sup>v</sup>	0.98	2.57	3.548 (4)	172

---

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $-x+3/2, -y, z-1/2$ ; (iii)  $x, -y+1/2, z-1/2$ ; (iv)  $x-1/2, -y+1/2, -z+1$ ; (v)  $x-1, y, z$ .