

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

2-Amino-5-(1*H*-tetrazol-5-yl)pyridinium chloride

Jing Dai and Xiao-Chun Wen\*

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China  
Correspondence e-mail: fudavid88@yahoo.com.cn


Received 11 September 2008; accepted 26 September 2008

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  
 $R$  factor = 0.048;  $wR$  factor = 0.125; data-to-parameter ratio = 16.4.

In the title salt,  $\text{C}_6\text{H}_7\text{N}_6^+\cdot\text{Cl}^-$ , there are two organic cations with similar conformations and two chloride anions in the asymmetric unit. The pyridine and tetrazole rings are essentially coplanar in each cation, with dihedral angles of 4.94 (15) and 5.41 (14)°. The pyridine N atoms are protonated. The crystal packing is stabilized by  $\text{N}-\text{H}\cdots\text{N}$  and  $\text{N}-\text{H}\cdots\text{Cl}$  hydrogen bonds, forming an infinite sheets parallel to the (101).

## Related literature

For uses of tetrazole derivatives, see: Dai & Fu (2008); Wang *et al.* (2005); Wen (2008); Xiong *et al.* (2002).



## Experimental

## Crystal data

 $\text{C}_6\text{H}_7\text{N}_6^+\cdot\text{Cl}^-$  $M_r = 198.63$ Monoclinic,  $P2_1/c$  $a = 14.043$  (3) Å $b = 13.238$  (3) Å $c = 9.5766$  (19) Å $\beta = 109.35$  (3)° $V = 1679.7$  (6) Å<sup>3</sup> $Z = 8$ Mo  $K\alpha$  radiation $\mu = 0.41$  mm<sup>-1</sup> $T = 298$  (2) K

0.45 × 0.25 × 0.20 mm

## Data collection

Rigaku Mercury2 diffractometer

Absorption correction: multi-scan

(CrystalClear; Rigaku, 2005)

 $T_{\min} = 0.859$ ,  $T_{\max} = 0.920$ 

16820 measured reflections

3855 independent reflections

3123 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.034$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$  $wR(F^2) = 0.125$  $S = 1.11$ 

3855 reflections

235 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.28$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.23$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{N5}^{\text{i}}$	0.86	2.59	3.317 (2)	143
$\text{N1}-\text{H1}\cdots\text{N6}^{\text{i}}$	0.86	2.04	2.894 (2)	171
$\text{N2}-\text{H2A}\cdots\text{N5}^{\text{i}}$	0.86	2.15	2.972 (3)	160
$\text{N7}-\text{H7}\cdots\text{N11}^{\text{ii}}$	0.86	2.57	3.304 (2)	144
$\text{N7}-\text{H7}\cdots\text{N12}^{\text{ii}}$	0.86	2.06	2.913 (2)	169
$\text{N8}-\text{H8A}\cdots\text{N11}^{\text{ii}}$	0.86	2.20	3.024 (3)	160
$\text{N2}-\text{H2B}\cdots\text{Cl1}^{\text{iii}}$	0.86	2.36	3.2187 (18)	177
$\text{N3}-\text{H3A}\cdots\text{Cl2}^{\text{iv}}$	0.86	2.17	3.0047 (19)	165
$\text{N8}-\text{H8B}\cdots\text{Cl2}^{\text{iv}}$	0.86	2.39	3.2432 (18)	172
$\text{N9}-\text{H9A}\cdots\text{Cl1}^{\text{v}}$	0.86	2.18	3.0031 (18)	160

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 2, y + \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (iv)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (v)  $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a start-up grant from Southeast University to Professor Ren-Gen Xiong.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2196).

## References

- Dai, W. & Fu, D.-W. (2008). *Acta Cryst.* **E64**, o1444.  
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Wang, X.-S., Tang, Y.-Z., Huang, X.-F., Qu, Z.-R., Che, C.-M., Chan, P. W. H. & Xiong, R.-G. (2005). *Inorg. Chem.* **44**, 5278–5285.  
Wen, X.-C. (2008). *Acta Cryst.* **E64**, m1033.  
Xiong, R.-G., Xue, X., Zhao, H., You, X.-Z., Abrahams, B. F. & Xue, Z. (2002). *Angew. Chem. Int. Ed.* **41**, 3800–3803.

## supporting information

*Acta Cryst.* (2008). E64, o2232 [doi:10.1107/S1600536808031164]

**2-Amino-5-(1*H*-tetrazol-5-yl)pyridinium chloride**

Jing Dai and Xiao-Chun Wen

**S1. Comment**

Tetrazole derivatives have found a wide range of applications in coordination chemistry because of their multiple coordination modes as ligands to metal ions, and for the construction of novel metal-organic frameworks (Wang *et al.*, 2005; Xiong *et al.*, 2002; Wen, 2008; Dai & Fu, 2008). We report here the crystal structure of the title salt (Fig. 1).


The title compound contains two organic cations with similar conformation and two Cl<sup>-</sup> ions in the asymmetric unit. The pyridine N atoms are protonated. The pyridine rings and tetrazole rings are nearly coplanar and are twisted from each other by a dihedral angle of only 4.94 (15) and 5.41 (14)°. The packing of ions is stabilized by N—H···N and N—H···Cl hydrogen bonds, to form an infinite two-dimensional layers parallel to the (1 0 1) plane in the crystal (Table 1 and Fig. 2).

**S2. Experimental**

2-Amino-5-cyanopyridine (30 mmol), NaN<sub>3</sub> (45 mmol), NH<sub>4</sub>Cl (33 mmol) and DMF (50 ml) were mixed in a flask under nitrogen atmosphere, and the mixture stirred at 383 K for 20 h. The resulting solution was then poured into ice-water (100 ml), and a white solid was obtained after adding hydrochloric acid (6 M) until pH=6. The precipitate was filtered and washed with distilled water. Colourless block-shaped crystals suitable for X-ray analysis were obtained from the crude product by slow evaporation of an ethanol/hydrochloric acid (50:1 v/v) solution.


**S3. Refinement**

All H atoms were fixed geometrically and treated as riding to their carrier atoms, with C—H = 0.93 Å (aromatic) and N—H = 0.86 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier atom})$ .



**Figure 1**

A view of the title compound with the atomic numbering scheme. Displacement ellipsoids were drawn at the 30% probability level.



**Figure 2**

The crystal packing of the title compound viewed along the *c* axis, showing the two-dimensional hydrogen bonds network. H atoms not involved in hydrogen bonding (dashed lines) have been omitted for clarity.

### 2-Amino-5-(1*H*-tetrazol-5-yl)pyridinium chloride

#### Crystal data

$C_6H_7N_6^+ \cdot Cl^-$   
 $M_r = 198.63$   
 Monoclinic,  $P2_1/c$   
 Hall symbol:  $-P\ 2_1/c$   
 $a = 14.043\ (3)\ \text{\AA}$   
 $b = 13.238\ (3)\ \text{\AA}$   
 $c = 9.5766\ (19)\ \text{\AA}$   
 $\beta = 109.35\ (3)^\circ$   
 $V = 1679.7\ (6)\ \text{\AA}^3$   
 $Z = 8$

$F(000) = 816$   
 $D_x = 1.571\ \text{Mg m}^{-3}$   
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$   
 Cell parameters from 3690 reflections  
 $\theta = 2.3\text{--}27.5^\circ$   
 $\mu = 0.41\ \text{mm}^{-1}$   
 $T = 298\ \text{K}$   
 Block, colorless  
 $0.45 \times 0.25 \times 0.20\ \text{mm}$

#### Data collection

Rigaku Mercury2  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 Detector resolution:  $13.6612\ \text{pixels mm}^{-1}$   
 $\omega$  scans

Absorption correction: multi-scan  
 (*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.859$ ,  $T_{\max} = 0.920$   
 16820 measured reflections  
 3855 independent reflections  
 3123 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$   
 $\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 2.7^\circ$   
 $h = -18 \rightarrow 18$

$k = -17 \rightarrow 17$   
 $l = -12 \rightarrow 12$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.125$   
 $S = 1.12$   
 3855 reflections  
 235 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0556P)^2 + 0.4684P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.28 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.23 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.56207 (12)	0.65987 (12)	0.17350 (18)	0.0344 (4)
H1	0.5639	0.7245	0.1662	0.041*
N3	0.36346 (13)	0.52444 (13)	0.37792 (19)	0.0368 (4)
H3A	0.3600	0.5892	0.3818	0.044*
C1	0.62505 (14)	0.60404 (14)	0.1236 (2)	0.0313 (4)
N2	0.68458 (13)	0.64965 (13)	0.0611 (2)	0.0413 (4)
H2A	0.6831	0.7143	0.0523	0.050*
H2B	0.7247	0.6146	0.0292	0.050*
N6	0.40734 (13)	0.37577 (12)	0.33322 (19)	0.0382 (4)
N4	0.31050 (14)	0.45963 (13)	0.4305 (2)	0.0428 (4)
C6	0.42266 (14)	0.47298 (14)	0.3184 (2)	0.0310 (4)
N5	0.33724 (14)	0.37053 (13)	0.4030 (2)	0.0430 (4)
C4	0.49237 (14)	0.51714 (13)	0.2516 (2)	0.0304 (4)
C2	0.62316 (14)	0.49827 (14)	0.1406 (2)	0.0333 (4)
H2	0.6657	0.4574	0.1086	0.040*
C3	0.55896 (15)	0.45611 (14)	0.2039 (2)	0.0336 (4)
H3	0.5587	0.3864	0.2160	0.040*
C5	0.49636 (15)	0.61909 (14)	0.2344 (2)	0.0337 (4)
H5	0.4537	0.6610	0.2646	0.040*
N7	0.92034 (13)	0.39582 (12)	0.30105 (18)	0.0364 (4)
H7	0.9131	0.4604	0.2998	0.044*
N9	1.13591 (12)	0.26242 (13)	0.12521 (19)	0.0367 (4)
H9A	1.1409	0.3272	0.1246	0.044*
N8	0.80738 (13)	0.38416 (13)	0.4302 (2)	0.0422 (4)
H8A	0.8043	0.4490	0.4319	0.051*
H8B	0.7726	0.3485	0.4708	0.051*
C7	0.86565 (14)	0.33931 (15)	0.3650 (2)	0.0329 (4)
N10	1.19013 (14)	0.19768 (13)	0.0747 (2)	0.0425 (4)
N12	1.08698 (13)	0.11357 (12)	0.15986 (19)	0.0374 (4)
C12	1.07262 (14)	0.21051 (14)	0.1769 (2)	0.0311 (4)
N11	1.16049 (14)	0.10833 (13)	0.0959 (2)	0.0429 (4)

C10	1.00037 (14)	0.25408 (14)	0.2397 (2)	0.0304 (4)
C9	0.94263 (15)	0.19202 (14)	0.3023 (2)	0.0339 (4)
H9	0.9504	0.1222	0.3025	0.041*
C11	0.98593 (16)	0.35597 (14)	0.2389 (2)	0.0363 (4)
H11	1.0211	0.3983	0.1957	0.044*
C8	0.87624 (15)	0.23312 (15)	0.3618 (2)	0.0337 (4)
H8	0.8378	0.1915	0.4004	0.040*
C11	0.84188 (4)	0.97883 (4)	0.45292 (7)	0.04632 (17)
C12	0.65611 (4)	0.24380 (4)	0.05567 (6)	0.04478 (17)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0407 (9)	0.0214 (8)	0.0500 (10)	0.0034 (7)	0.0270 (8)	0.0030 (7)
N3	0.0440 (10)	0.0261 (8)	0.0491 (10)	-0.0020 (7)	0.0273 (8)	-0.0008 (7)
C1	0.0332 (10)	0.0274 (9)	0.0362 (10)	0.0024 (7)	0.0157 (8)	0.0009 (7)
N2	0.0483 (11)	0.0305 (9)	0.0589 (11)	0.0027 (7)	0.0363 (9)	0.0028 (8)
N6	0.0456 (10)	0.0267 (8)	0.0507 (10)	-0.0021 (7)	0.0270 (8)	0.0009 (7)
N4	0.0518 (11)	0.0327 (9)	0.0561 (11)	-0.0061 (8)	0.0343 (9)	-0.0012 (8)
C6	0.0334 (10)	0.0265 (9)	0.0358 (10)	-0.0001 (7)	0.0152 (8)	-0.0004 (7)
N5	0.0529 (11)	0.0299 (9)	0.0581 (11)	-0.0034 (8)	0.0346 (9)	-0.0008 (8)
C4	0.0313 (9)	0.0259 (9)	0.0368 (10)	-0.0001 (7)	0.0149 (8)	0.0011 (7)
C2	0.0341 (10)	0.0284 (9)	0.0419 (11)	0.0044 (8)	0.0186 (8)	-0.0002 (8)
C3	0.0368 (10)	0.0238 (9)	0.0422 (11)	0.0017 (7)	0.0158 (8)	-0.0001 (8)
C5	0.0358 (10)	0.0281 (9)	0.0437 (11)	0.0035 (8)	0.0218 (8)	0.0018 (8)
N7	0.0468 (10)	0.0227 (8)	0.0486 (10)	0.0051 (7)	0.0279 (8)	0.0046 (7)
N9	0.0410 (10)	0.0264 (8)	0.0510 (10)	0.0036 (7)	0.0265 (8)	0.0027 (7)
N8	0.0473 (11)	0.0334 (9)	0.0586 (11)	0.0043 (8)	0.0345 (9)	0.0034 (8)
C7	0.0328 (10)	0.0327 (10)	0.0353 (10)	0.0024 (8)	0.0143 (8)	0.0018 (8)
N10	0.0473 (11)	0.0343 (9)	0.0569 (11)	0.0066 (8)	0.0320 (9)	0.0036 (8)
N12	0.0431 (10)	0.0293 (8)	0.0477 (10)	0.0038 (7)	0.0256 (8)	0.0015 (7)
C12	0.0348 (10)	0.0266 (9)	0.0339 (9)	0.0021 (7)	0.0142 (8)	0.0033 (7)
N11	0.0489 (11)	0.0344 (10)	0.0560 (11)	0.0059 (8)	0.0316 (9)	0.0010 (8)
C10	0.0328 (10)	0.0282 (9)	0.0325 (9)	0.0027 (7)	0.0139 (8)	0.0015 (7)
C9	0.0375 (10)	0.0236 (9)	0.0431 (11)	0.0009 (7)	0.0168 (9)	0.0031 (8)
C11	0.0427 (11)	0.0296 (10)	0.0446 (11)	0.0017 (8)	0.0253 (9)	0.0037 (8)
C8	0.0344 (10)	0.0298 (10)	0.0409 (10)	-0.0010 (8)	0.0178 (8)	0.0040 (8)
C11	0.0567 (3)	0.0300 (3)	0.0626 (4)	-0.0067 (2)	0.0336 (3)	-0.0052 (2)
C12	0.0502 (3)	0.0282 (3)	0.0624 (4)	0.0046 (2)	0.0273 (3)	0.0040 (2)

*Geometric parameters (Å, °)*

N1—C1	1.355 (2)	N7—C7	1.355 (2)
N1—C5	1.356 (2)	N7—C11	1.358 (2)
N1—H1	0.8600	N7—H7	0.8600
N3—N4	1.338 (2)	N9—N10	1.339 (2)
N3—C6	1.340 (2)	N9—C12	1.341 (2)
N3—H3A	0.8600	N9—H9A	0.8600

C1—N2	1.325 (2)	N8—C7	1.323 (2)
C1—C2	1.411 (3)	N8—H8A	0.8600
N2—H2A	0.8600	N8—H8B	0.8600
N2—H2B	0.8600	C7—C8	1.415 (3)
N6—C6	1.320 (2)	N10—N11	1.292 (2)
N6—N5	1.362 (2)	N12—C12	1.318 (2)
N4—N5	1.291 (2)	N12—N11	1.365 (2)
C6—C4	1.457 (2)	C12—C10	1.459 (2)
C4—C5	1.363 (3)	C10—C11	1.364 (3)
C4—C3	1.421 (3)	C10—C9	1.419 (3)
C2—C3	1.361 (3)	C9—C8	1.357 (3)
C2—H2	0.9300	C9—H9	0.9300
C3—H3	0.9300	C11—H11	0.9300
C5—H5	0.9300	C8—H8	0.9300
C1—N1—C5	123.42 (16)	C7—N7—C11	123.49 (17)
C1—N1—H1	118.3	C7—N7—H7	118.3
C5—N1—H1	118.3	C11—N7—H7	118.3
N4—N3—C6	109.55 (16)	N10—N9—C12	109.34 (16)
N4—N3—H3A	125.2	N10—N9—H9A	125.3
C6—N3—H3A	125.2	C12—N9—H9A	125.3
N2—C1—N1	119.62 (17)	C7—N8—H8A	120.0
N2—C1—C2	122.91 (17)	C7—N8—H8B	120.0
N1—C1—C2	117.47 (16)	H8A—N8—H8B	120.0
C1—N2—H2A	120.0	N8—C7—N7	119.82 (18)
C1—N2—H2B	120.0	N8—C7—C8	122.80 (18)
H2A—N2—H2B	120.0	N7—C7—C8	117.36 (17)
C6—N6—N5	105.81 (16)	N11—N10—N9	106.13 (15)
N5—N4—N3	105.89 (15)	C12—N12—N11	105.95 (16)
N6—C6—N3	107.67 (16)	N12—C12—N9	107.80 (16)
N6—C6—C4	126.56 (17)	N12—C12—C10	126.32 (17)
N3—C6—C4	125.76 (17)	N9—C12—C10	125.87 (17)
N4—N5—N6	111.08 (16)	N10—N11—N12	110.78 (16)
C5—C4—C3	117.77 (17)	C11—C10—C9	118.05 (17)
C5—C4—C6	120.74 (17)	C11—C10—C12	120.76 (17)
C3—C4—C6	121.49 (16)	C9—C10—C12	121.19 (17)
C3—C2—C1	119.91 (17)	C8—C9—C10	120.89 (17)
C3—C2—H2	120.0	C8—C9—H9	119.6
C1—C2—H2	120.0	C10—C9—H9	119.6
C2—C3—C4	120.90 (17)	N7—C11—C10	120.20 (17)
C2—C3—H3	119.5	N7—C11—H11	119.9
C4—C3—H3	119.5	C10—C11—H11	119.9
N1—C5—C4	120.49 (17)	C9—C8—C7	119.93 (17)
N1—C5—H5	119.8	C9—C8—H8	120.0
C4—C5—H5	119.8	C7—C8—H8	120.0
C5—N1—C1—N2	177.74 (19)	C11—N7—C7—N8	176.37 (19)
C5—N1—C1—C2	-1.8 (3)	C11—N7—C7—C8	-1.9 (3)

C6—N3—N4—N5	0.0 (2)	C12—N9—N10—N11	-0.2 (2)
N5—N6—C6—N3	0.1 (2)	N11—N12—C12—N9	-0.2 (2)
N5—N6—C6—C4	179.20 (18)	N11—N12—C12—C10	179.35 (18)
N4—N3—C6—N6	0.0 (2)	N10—N9—C12—N12	0.3 (2)
N4—N3—C6—C4	-179.15 (19)	N10—N9—C12—C10	-179.30 (18)
N3—N4—N5—N6	0.1 (2)	N9—N10—N11—N12	0.1 (2)
C6—N6—N5—N4	-0.1 (2)	C12—N12—N11—N10	0.1 (2)
N6—C6—C4—C5	175.7 (2)	N12—C12—C10—C11	-173.8 (2)
N3—C6—C4—C5	-5.4 (3)	N9—C12—C10—C11	5.6 (3)
N6—C6—C4—C3	-4.9 (3)	N12—C12—C10—C9	5.7 (3)
N3—C6—C4—C3	173.99 (19)	N9—C12—C10—C9	-174.82 (19)
N2—C1—C2—C3	-178.97 (19)	C11—C10—C9—C8	-1.1 (3)
N1—C1—C2—C3	0.6 (3)	C12—C10—C9—C8	179.35 (18)
C1—C2—C3—C4	0.9 (3)	C7—N7—C11—C10	-0.6 (3)
C5—C4—C3—C2	-1.2 (3)	C9—C10—C11—N7	2.1 (3)
C6—C4—C3—C2	179.48 (19)	C12—C10—C11—N7	-178.31 (18)
C1—N1—C5—C4	1.6 (3)	C10—C9—C8—C7	-1.4 (3)
C3—C4—C5—N1	0.0 (3)	N8—C7—C8—C9	-175.3 (2)
C6—C4—C5—N1	179.36 (18)	N7—C7—C8—C9	2.9 (3)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 $\cdots$ N5 <sup>i</sup>	0.86	2.59	3.317 (2)	143
N1—H1 $\cdots$ N6 <sup>i</sup>	0.86	2.04	2.894 (2)	171
N2—H2A $\cdots$ N5 <sup>i</sup>	0.86	2.15	2.972 (3)	160
N7—H7 $\cdots$ N11 <sup>ii</sup>	0.86	2.57	3.304 (2)	144
N7—H7 $\cdots$ N12 <sup>ii</sup>	0.86	2.06	2.913 (2)	169
N8—H8A $\cdots$ N11 <sup>ii</sup>	0.86	2.20	3.024 (3)	160
N2—H2B $\cdots$ C11 <sup>iii</sup>	0.86	2.36	3.2187 (18)	177
N3—H3A $\cdots$ C12 <sup>i</sup>	0.86	2.17	3.0047 (19)	165
N8—H8B $\cdots$ C12 <sup>iv</sup>	0.86	2.39	3.2432 (18)	172
N9—H9A $\cdots$ C11 <sup>v</sup>	0.86	2.18	3.0031 (18)	160

Symmetry codes: (i)  $-x+1, y+1/2, -z+1/2$ ; (ii)  $-x+2, y+1/2, -z+1/2$ ; (iii)  $x, -y+3/2, z-1/2$ ; (iv)  $x, -y+1/2, z+1/2$ ; (v)  $-x+2, y-1/2, -z+1/2$ .