

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1,3,4,6-Tetrachloro-7,7-bis(4-chlorophenyl)bicyclo[4.2.0]oct-3-ene-2,5-dione

Huayou Hu,\* Lei Li, Jun-Feng Ji and Zhi-Guo Shen

 School of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China  
 Correspondence e-mail: huayouhu@seu.edu.cn


Received 24 September 2008; accepted 26 September 2008

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.008$  Å;  $R$  factor = 0.066;  $wR$  factor = 0.192; data-to-parameter ratio = 14.3.

The title compound,  $\text{C}_{20}\text{H}_{10}\text{Cl}_6\text{O}_2$ , a quinone derivative, was obtained by the irradiation of 2,3,5,6-tetrachlorobenzoquinone and 4,4'-(ethene-1,1-diyl)bis(chlorobenzene). The six- and four-membered rings are fused in a *cis* configuration. The dihedral angle between them is  $53.4$  ( $3$ )°.

## Related literature

For related literature, see: Eckert & Goetz (1994); Miyashi *et al.* (1985); Schenk (1960); Xu, Song *et al.* (1994); Xu, Wang *et al.* (1994); Xue *et al.* (2000). For a related structure, see: Braun *et al.* (1999)



## Experimental

## Crystal data

$\text{C}_{20}\text{H}_{10}\text{Cl}_6\text{O}_2$	$\gamma = 102.68$ ( $3$ )°
$M_r = 494.98$	$V = 996.4$ ( $3$ ) Å <sup>3</sup>
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.6710$ ( $17$ ) Å	Mo $K\alpha$ radiation
$b = 9.6850$ ( $19$ ) Å	$\mu = 0.88$ mm <sup>-1</sup>
$c = 12.864$ ( $3$ ) Å	$T = 293$ ( $2$ ) K
$\alpha = 105.49$ ( $3$ )°	$0.30 \times 0.20 \times 0.10$ mm
$\beta = 97.11$ ( $3$ )°	

## Data collection

Enraf–Nonius CAD-4 diffractometer	3619 independent reflections
Absorption correction: $\psi$ scan (SHELXTL; Sheldrick, 2008)	2787 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.779$ , $T_{\max} = 0.917$	$R_{\text{int}} = 0.049$
3879 measured reflections	3 standard reflections every 200 reflections intensity decay: none

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.066$	253 parameters
$wR(F^2) = 0.192$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.66$ e Å <sup>-3</sup>
3619 reflections	$\Delta\rho_{\min} = -0.57$ e Å <sup>-3</sup>

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms, 1993); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Program for Young Excellent Talents in Southeast University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2797).

## References

- Braun, M., Christl, M., Peters, E.-M. & Peters, K. (1999). *J. Chem. Soc. Perkin Trans. 1*, pp. 2813–2820.
- Eckert, G. & Goetz, M. (1994). *J. Am. Chem. Soc.* **116**, 11999–12009.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. (1993). *XCAD4*. University of Marburg, Germany.
- Miyashi, T., Takahashi, Y., Mukai, T., Roth, H. D. & Schilling, M. L. M. (1985). *J. Am. Chem. Soc.* **107**, 1079–1080.
- Schenk, G. O. Z. (1960). *Electrochemistry*, **64**, 997–1011.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xu, J. H., Song, Y. L., Zhang, Z. G., Wang, L. C. & Xu, J. W. (1994). *Tetrahedron*, **50**, 1199–1210.
- Xu, J. H., Wang, L. C., Xu, J. W., Yan, B. Z. & Yuan, H. C. (1994). *J. Chem. Soc. Perkin Trans. 1*, pp. 571–577.
- Xue, J., Xu, J.-W., Yang, L. & Xu, J.-H. (2000). *J. Org. Chem.* **65**, 30–40.

## supporting information

*Acta Cryst.* (2008). E64, o2151 [doi:10.1107/S1600536808031139]

**1,3,4,6-Tetrachloro-7,7-bis(4-chlorophenyl)bicyclo[4.2.0]oct-3-ene-2,5-dione**

Huayou Hu, Lei Li, Jun-Feng Ji and Zhi-Guo Shen

**S1. Comment**

The reactions of the high potential 2,3,5,6-tetrachlorobenzoquinone with alkenes display varied reaction sites and regioselectivity, depending on the structure of the alkenes and reaction conditions (Schenk 1960; Miyashi *et al.* 1985; Eckert & Goetz 1994; Xu, Song *et al.* 1994; Xu, Wang *et al.* 1994). While irradiation of a benzene solution of 2,3,5,6-tetrachlorobenzoquinone and 4,4'-(ethene-1,1-diyl)bis(chlorobenzene) with light of wavelength longer than 400 nm resulted in formation products of the title compound as a yellow solid (Xue *et al.* 2000). The yellow crystals were obtained by recrystallization of these solids from petroleum ether-chloroform.


The title compound, C<sub>20</sub>H<sub>10</sub>Cl<sub>6</sub>O<sub>2</sub>, is a quinone derivative. In the quinone, the distances of the C=O bonds are 1.191 (7) and 1.199 (7) Å, which are considered to have full double-bond character. Meanwhile, the distances of C1—C2 and C5—C6 are, respectively, 1.478 (9) and 1.475 (8) Å, which are a little longer than that of C1=C6 (1.354 (9) Å), but shorter than those of C—C bonds (1.527 (8)–1.560 (7) Å). This shows that C1—C2 and C5—C6 bonds both have part double-bond character.

**S2. Experimental**

Irradiation of a benzene solution of 2,3,5,6-tetrachlorobenzoquinone (0.05 mol L<sup>-1</sup>) and 4,4'-(ethene-1,1-diyl)bis(chlorobenzene) (0.10 mol L<sup>-1</sup>) with light of wavelength longer than 400 nm for 10 h resulted in complete consumption of 2,3,5,6-tetrachlorobenzoquinone and the formation of products 1,3,4,6-tetrachloro-7,7-bis(4-chlorophenyl)bicyclo[4.2.0]oct-3-ene-2,5-dione. Recrystallization from petroleum ether (bp 60–90 °) and chloroform gave a slightly yellow crystal.

**S3. Refinement**

H atoms were positioned geometrically and refined using a riding model with C—H = 0.93–0.97 Å and with  $U_{\text{iso}}(\text{H}) = 1.2$  times  $U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of the title compound with 30% displacement ellipsoids.

### 1,3,4,6-Tetrachloro-7,7-bis(4-chlorophenyl)bicyclo[4.2.0]oct-3-ene-2,5-dione

#### Crystal data

$C_{20}H_{10}Cl_6O_2$

$M_r = 494.98$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 8.6710$  (17) Å

$b = 9.6850$  (19) Å

$c = 12.864$  (3) Å

$\alpha = 105.49$  (3)°

$\beta = 97.11$  (3)°

$\gamma = 102.68$  (3)°

$V = 996.4$  (3) Å<sup>3</sup>

$Z = 2$

$F(000) = 496$

$D_x = 1.650$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 25 reflections

$\theta = 10\text{--}13^\circ$

$\mu = 0.88$  mm<sup>-1</sup>

$T = 293$  K

Block, yellow

$0.30 \times 0.20 \times 0.10$  mm

*Data collection*

Enraf–Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan  
(*SHELXTL*; Sheldrick, 2008)

$T_{\min} = 0.779$ ,  $T_{\max} = 0.917$

3879 measured reflections

3619 independent reflections

2787 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.049$

$\theta_{\max} = 25.3^\circ$ ,  $\theta_{\min} = 1.7^\circ$

$h = -10 \rightarrow 10$

$k = -11 \rightarrow 11$

$l = 0 \rightarrow 15$

3 standard reflections every 200 reflections

intensity decay: none

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.066$

$wR(F^2) = 0.192$

$S = 1.01$

3619 reflections

253 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + 6P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.66 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.58 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.3758 (2)	0.1746 (2)	0.61309 (16)	0.0752 (6)
C12	0.0143 (2)	−0.07268 (16)	0.83090 (14)	0.0527 (4)
C13	0.4819 (2)	0.4713 (2)	0.81185 (19)	0.0761 (6)
C14	0.25830 (19)	0.20039 (18)	1.01117 (12)	0.0524 (4)
C15	−0.2014 (3)	0.5682 (2)	0.53637 (17)	0.0788 (6)
C16	−0.7475 (2)	−0.30216 (19)	0.73595 (17)	0.0653 (5)
O1	0.0754 (6)	−0.0161 (5)	0.6230 (4)	0.0625 (12)
O2	0.2265 (5)	0.4901 (4)	0.9419 (4)	0.0574 (11)
C1	0.2697 (7)	0.2059 (7)	0.7168 (5)	0.0444 (13)
C2	0.1230 (7)	0.0867 (6)	0.7046 (4)	0.0409 (13)
C3	0.0404 (6)	0.1008 (6)	0.8037 (4)	0.0365 (11)
C4	0.1223 (7)	0.2309 (6)	0.9110 (4)	0.0382 (12)
C5	0.2210 (6)	0.3667 (6)	0.8877 (4)	0.0384 (12)
C6	0.3165 (6)	0.3339 (6)	0.8013 (5)	0.0422 (13)
C7	−0.0468 (6)	0.2385 (6)	0.9327 (4)	0.0385 (12)

H7A	-0.0851	0.1819	0.9806	0.046*
H7B	-0.0580	0.3389	0.9577	0.046*
C8	-0.1175 (6)	0.1593 (6)	0.8105 (4)	0.0337 (11)
C9	-0.1295 (6)	0.2656 (6)	0.7408 (4)	0.0360 (11)
C10	-0.1477 (7)	0.2165 (6)	0.6274 (5)	0.0449 (13)
H10A	-0.1463	0.1193	0.5932	0.054*
C11	-0.1680 (9)	0.3087 (7)	0.5634 (5)	0.0554 (16)
H11A	-0.1766	0.2747	0.4876	0.067*
C12	-0.1752 (7)	0.4491 (6)	0.6133 (5)	0.0445 (13)
C13	-0.1599 (7)	0.5022 (6)	0.7252 (5)	0.0452 (13)
H13A	-0.1630	0.5991	0.7585	0.054*
C14	-0.1399 (7)	0.4087 (6)	0.7877 (5)	0.0459 (14)
H14A	-0.1332	0.4432	0.8633	0.055*
C15	-0.2792 (6)	0.0423 (6)	0.7864 (4)	0.0341 (11)
C16	-0.3207 (7)	-0.0869 (6)	0.6989 (5)	0.0441 (13)
H16A	-0.2506	-0.1029	0.6505	0.053*
C17	-0.4660 (7)	-0.1935 (7)	0.6820 (5)	0.0493 (14)
H17A	-0.4930	-0.2801	0.6229	0.059*
C18	-0.5684 (7)	-0.1687 (6)	0.7538 (5)	0.0431 (13)
C19	-0.5304 (7)	-0.0393 (7)	0.8409 (5)	0.0487 (14)
H19A	-0.6005	-0.0232	0.8893	0.058*
C20	-0.3883 (7)	0.0644 (6)	0.8548 (4)	0.0421 (13)
H20A	-0.3642	0.1527	0.9122	0.050*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0764 (12)	0.0926 (14)	0.0673 (11)	0.0278 (10)	0.0554 (10)	0.0190 (10)
C12	0.0647 (10)	0.0394 (8)	0.0665 (10)	0.0226 (7)	0.0264 (8)	0.0234 (7)
C13	0.0535 (10)	0.0653 (11)	0.1098 (16)	0.0045 (8)	0.0405 (10)	0.0250 (10)
C14	0.0565 (9)	0.0623 (9)	0.0436 (8)	0.0229 (7)	0.0147 (6)	0.0167 (7)
C15	0.1310 (18)	0.0587 (10)	0.0781 (12)	0.0495 (11)	0.0479 (12)	0.0402 (10)
C16	0.0518 (9)	0.0553 (10)	0.0947 (13)	0.0112 (7)	0.0326 (9)	0.0267 (9)
O1	0.080 (3)	0.054 (3)	0.049 (3)	0.019 (2)	0.038 (2)	-0.003 (2)
O2	0.069 (3)	0.037 (2)	0.064 (3)	0.014 (2)	0.032 (2)	0.003 (2)
C1	0.050 (3)	0.052 (3)	0.043 (3)	0.025 (3)	0.031 (3)	0.015 (3)
C2	0.048 (3)	0.049 (3)	0.039 (3)	0.030 (3)	0.026 (2)	0.014 (3)
C3	0.048 (3)	0.034 (3)	0.036 (3)	0.019 (2)	0.023 (2)	0.011 (2)
C4	0.047 (3)	0.034 (3)	0.035 (3)	0.013 (2)	0.018 (2)	0.005 (2)
C5	0.036 (3)	0.038 (3)	0.039 (3)	0.010 (2)	0.014 (2)	0.005 (2)
C6	0.032 (3)	0.047 (3)	0.052 (3)	0.014 (2)	0.015 (2)	0.017 (3)
C7	0.047 (3)	0.048 (3)	0.029 (3)	0.023 (2)	0.019 (2)	0.011 (2)
C8	0.042 (3)	0.038 (3)	0.031 (2)	0.021 (2)	0.021 (2)	0.012 (2)
C9	0.044 (3)	0.035 (3)	0.040 (3)	0.020 (2)	0.026 (2)	0.014 (2)
C10	0.067 (4)	0.037 (3)	0.040 (3)	0.028 (3)	0.023 (3)	0.010 (2)
C11	0.091 (5)	0.045 (3)	0.044 (3)	0.033 (3)	0.034 (3)	0.016 (3)
C12	0.047 (3)	0.036 (3)	0.058 (4)	0.014 (2)	0.023 (3)	0.018 (3)
C13	0.055 (3)	0.025 (3)	0.055 (3)	0.013 (2)	0.023 (3)	0.005 (2)

C14	0.060 (4)	0.039 (3)	0.046 (3)	0.021 (3)	0.031 (3)	0.010 (2)
C15	0.041 (3)	0.035 (3)	0.031 (3)	0.012 (2)	0.015 (2)	0.013 (2)
C16	0.048 (3)	0.043 (3)	0.044 (3)	0.015 (2)	0.029 (3)	0.006 (2)
C17	0.046 (3)	0.042 (3)	0.054 (4)	0.011 (3)	0.019 (3)	0.003 (3)
C18	0.038 (3)	0.047 (3)	0.051 (3)	0.014 (2)	0.013 (2)	0.022 (3)
C19	0.049 (3)	0.057 (4)	0.054 (3)	0.024 (3)	0.035 (3)	0.021 (3)
C20	0.050 (3)	0.043 (3)	0.039 (3)	0.020 (3)	0.021 (2)	0.011 (2)

*Geometric parameters (Å, °)*

C11—C1	1.712 (5)	C9—C14	1.384 (7)
C12—C3	1.779 (5)	C9—C10	1.385 (7)
C13—C6	1.693 (6)	C10—C11	1.390 (8)
C14—C4	1.766 (6)	C10—H10A	0.9300
C15—C12	1.740 (6)	C11—C12	1.359 (8)
C16—C18	1.734 (6)	C11—H11A	0.9300
O1—C2	1.190 (7)	C12—C13	1.372 (8)
O2—C5	1.200 (6)	C13—C14	1.385 (8)
C1—C6	1.354 (8)	C13—H13A	0.9300
C1—C2	1.478 (8)	C14—H14A	0.9300
C2—C3	1.528 (7)	C15—C20	1.382 (7)
C3—C4	1.560 (7)	C15—C16	1.383 (7)
C3—C8	1.596 (7)	C16—C17	1.393 (8)
C4—C5	1.526 (7)	C16—H16A	0.9300
C4—C7	1.540 (7)	C17—C18	1.370 (8)
C5—C6	1.474 (7)	C17—H17A	0.9300
C7—C8	1.532 (7)	C18—C19	1.383 (8)
C7—H7A	0.9700	C19—C20	1.365 (8)
C7—H7B	0.9700	C19—H19A	0.9300
C8—C15	1.534 (7)	C20—H20A	0.9300
C8—C9	1.545 (7)		
C6—C1—C2	123.4 (5)	C14—C9—C10	117.1 (5)
C6—C1—C11	121.3 (5)	C14—C9—C8	121.3 (5)
C2—C1—C11	115.2 (4)	C10—C9—C8	121.2 (5)
O1—C2—C1	121.5 (5)	C9—C10—C11	121.7 (5)
O1—C2—C3	122.3 (5)	C9—C10—H10A	119.1
C1—C2—C3	116.1 (5)	C11—C10—H10A	119.1
C2—C3—C4	118.3 (5)	C12—C11—C10	119.0 (6)
C2—C3—C8	123.0 (4)	C12—C11—H11A	120.5
C4—C3—C8	86.8 (4)	C10—C11—H11A	120.5
C2—C3—C12	106.0 (3)	C11—C12—C13	121.4 (5)
C4—C3—C12	110.3 (4)	C11—C12—C15	120.6 (5)
C8—C3—C12	111.5 (3)	C13—C12—C15	118.0 (4)
C5—C4—C7	115.4 (4)	C12—C13—C14	118.8 (5)
C5—C4—C3	112.4 (4)	C12—C13—H13A	120.6
C7—C4—C3	88.4 (4)	C14—C13—H13A	120.6
C5—C4—C14	103.3 (4)	C9—C14—C13	121.9 (5)

C7—C4—C14	118.9 (4)	C9—C14—H14A	119.0
C3—C4—C14	118.7 (4)	C13—C14—H14A	119.0
O2—C5—C6	123.6 (5)	C20—C15—C16	117.9 (5)
O2—C5—C4	121.0 (5)	C20—C15—C8	119.4 (5)
C6—C5—C4	115.2 (5)	C16—C15—C8	122.6 (4)
C1—C6—C5	122.0 (5)	C15—C16—C17	121.0 (5)
C1—C6—C13	122.7 (4)	C15—C16—H16A	119.5
C5—C6—C13	115.2 (4)	C17—C16—H16A	119.5
C8—C7—C4	89.8 (4)	C18—C17—C16	119.0 (5)
C8—C7—H7A	113.7	C18—C17—H17A	120.5
C4—C7—H7A	113.7	C16—C17—H17A	120.5
C8—C7—H7B	113.7	C17—C18—C19	120.9 (5)
C4—C7—H7B	113.7	C17—C18—C16	119.6 (5)
H7A—C7—H7B	110.9	C19—C18—C16	119.5 (4)
C7—C8—C15	114.9 (4)	C20—C19—C18	119.0 (5)
C7—C8—C9	114.1 (4)	C20—C19—H19A	120.5
C15—C8—C9	109.7 (4)	C18—C19—H19A	120.5
C7—C8—C3	87.4 (4)	C19—C20—C15	122.0 (5)
C15—C8—C3	117.2 (4)	C19—C20—H20A	119.0
C9—C8—C3	112.1 (4)	C15—C20—H20A	119.0
C6—C1—C2—O1	171.0 (6)	C4—C3—C8—C7	-20.4 (4)
C11—C1—C2—O1	-6.9 (8)	C12—C3—C8—C7	90.1 (4)
C6—C1—C2—C3	-11.0 (8)	C2—C3—C8—C15	100.7 (6)
C11—C1—C2—C3	171.1 (4)	C4—C3—C8—C15	-137.3 (4)
O1—C2—C3—C4	173.7 (5)	C12—C3—C8—C15	-26.7 (5)
C1—C2—C3—C4	-4.3 (7)	C2—C3—C8—C9	-27.4 (7)
O1—C2—C3—C8	-80.5 (7)	C4—C3—C8—C9	94.6 (4)
C1—C2—C3—C8	101.6 (6)	C12—C3—C8—C9	-154.9 (4)
O1—C2—C3—C12	49.4 (7)	C7—C8—C9—C14	-25.0 (7)
C1—C2—C3—C12	-128.6 (4)	C15—C8—C9—C14	105.5 (6)
C2—C3—C4—C5	29.4 (6)	C3—C8—C9—C14	-122.4 (5)
C8—C3—C4—C5	-96.7 (4)	C7—C8—C9—C10	162.0 (5)
C12—C3—C4—C5	151.6 (4)	C15—C8—C9—C10	-67.4 (6)
C2—C3—C4—C7	146.4 (5)	C3—C8—C9—C10	64.7 (7)
C8—C3—C4—C7	20.3 (4)	C14—C9—C10—C11	2.9 (9)
C12—C3—C4—C7	-91.5 (4)	C8—C9—C10—C11	176.2 (6)
C2—C3—C4—C14	-91.2 (5)	C9—C10—C11—C12	-2.1 (10)
C8—C3—C4—C14	142.7 (4)	C10—C11—C12—C13	1.2 (10)
C12—C3—C4—C14	30.9 (5)	C10—C11—C12—C15	-179.7 (5)
C7—C4—C5—O2	44.0 (7)	C11—C12—C13—C14	-1.2 (9)
C3—C4—C5—O2	143.4 (5)	C15—C12—C13—C14	179.7 (5)
C14—C4—C5—O2	-87.4 (6)	C10—C9—C14—C13	-3.0 (9)
C7—C4—C5—C6	-140.6 (5)	C8—C9—C14—C13	-176.2 (5)
C3—C4—C5—C6	-41.2 (6)	C12—C13—C14—C9	2.2 (9)
C14—C4—C5—C6	88.0 (5)	C7—C8—C15—C20	34.1 (7)
C2—C1—C6—C5	-1.7 (9)	C9—C8—C15—C20	-96.0 (5)
C11—C1—C6—C5	176.1 (4)	C3—C8—C15—C20	134.7 (5)

---

C2—C1—C6—C13	-178.1 (4)	C7—C8—C15—C16	-145.2 (5)
C11—C1—C6—C13	-0.3 (8)	C9—C8—C15—C16	84.8 (6)
O2—C5—C6—C1	-155.5 (6)	C3—C8—C15—C16	-44.6 (7)
C4—C5—C6—C1	29.3 (8)	C20—C15—C16—C17	-1.8 (9)
O2—C5—C6—C13	21.2 (8)	C8—C15—C16—C17	177.4 (5)
C4—C5—C6—C13	-154.1 (4)	C15—C16—C17—C18	0.0 (9)
C5—C4—C7—C8	93.0 (5)	C16—C17—C18—C19	1.0 (9)
C3—C4—C7—C8	-21.1 (4)	C16—C17—C18—C16	-178.5 (5)
C14—C4—C7—C8	-143.4 (4)	C17—C18—C19—C20	-0.1 (9)
C4—C7—C8—C15	139.6 (4)	C16—C18—C19—C20	179.4 (5)
C4—C7—C8—C9	-92.5 (5)	C18—C19—C20—C15	-1.8 (9)
C4—C7—C8—C3	20.7 (4)	C16—C15—C20—C19	2.8 (8)
C2—C3—C8—C7	-142.4 (5)	C8—C15—C20—C19	-176.5 (5)

---