

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Chloro-*N*-(3-phenylallylidene)aniline

Bing-Xiang Zhang

Department of Chemistry, Taishan University, 271021 Taian, Shandong, People's Republic of China

Correspondence e-mail: bezhbx@163.com


Received 10 November 2008; accepted 24 November 2008

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.026; wR factor = 0.066; data-to-parameter ratio = 14.2.

In the title molecule, $\text{C}_{15}\text{H}_{12}\text{ClN}$, the $\text{C}=\text{N}$ and $\text{C}=\text{C}$ bond lengths are 1.273 (2) and 1.324 (2) Å, respectively. The two aromatic rings form a dihedral angle of 3.27 (3)°.

Related literature

For a related structure, see Pu (2008). For general background, see: Garnovskii *et al.* (1993); Anderson *et al.* (1997); Musie *et al.* (2001); Paul *et al.* (2002). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{12}\text{ClN}$ $M_r = 241.71$ Orthorhombic, $Pna2_1$ $a = 7.7333$ (7) Å $b = 5.5957$ (5) Å $c = 29.383$ (3) Å $V = 1271.5$ (2) Å³ $Z = 4$ Mo $K\alpha$ radiation
 $\mu = 0.28$ mm⁻¹ $T = 295$ (2) K
 $0.15 \times 0.12 \times 0.08$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

 $T_{\min} = 0.960$, $T_{\max} = 0.978$ 6043 measured reflections
2187 independent reflections
2042 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.026$ $wR(F^2) = 0.066$ $S = 1.05$

2187 reflections

154 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.09$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Absolute structure: Flack (1983),

1031 Friedel pairs

Flack parameter: 0.07 (5)

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This project was supported by the Postgraduate Foundation of Taishan University (grant No. Y03-1-13).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2477).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1-19.
- Anderson, O. P., Cour, A. L., Findeisen, M., Hennig, L., Simonsen, O., Taylor, L. & Toftund, H. (1997). *J. Chem. Soc. Dalton Trans.* pp. 111-120.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876-881.
- Garnovskii, A. D., Nivorozhkin, A. L. & Minki, V. I. (1993). *Coord. Chem. Rev.* **126**, 1-69.
- Musie, G. T., Wei, M., Subramaniam, B. & Busch, D. H. (2001). *Inorg. Chem.* **40**, 3336-3341.
- Paul, S., Barik, A. K., Peng, S. M. & Kar, S. K. (2002). *Inorg. Chem.* **41**, 5803-5809.
- Pu, X.-H. (2008). *Acta Cryst.* **E64**, o1734.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.

supporting information

Acta Cryst. (2008). E64, o2478 [doi:10.1107/S1600536808039391]

4-Chloro-*N*-(3-phenylallylidene)aniline

Bing-Xiang Zhang

S1. Comment

Schiff base compounds have been of great interest for many years. These compounds play an important role in the development of coordination chemistry related to catalysis and enzymatic reactions, magnetism and molecular architectures (Garnovskii *et al.*, 1993; Anderson *et al.*, 1997; Musie *et al.*, 2001; Paul *et al.*, 2002; Pu, 2008). In order to search for new Schiff-bases with higher bioactivity, the title compound (I) was synthesized and its crystal structure determined.

In (I) (Fig. 1), the bond lengths and angles are in good agreement with the expected values (Allen *et al.*, 1987).

S2. Experimental

The title compound was synthesized by the reaction of 4-Chloro-phenylamine (1 mmol, 127.6 mg) with 3-Phenyl-propenal (1 mmol, 132.2 mg) in ethanol (20 ml) under reflux conditions (338 K) for 5 h. The solvent was removed and the solid product recrystallized from tetrahydrofuran. After five days yellow crystals suitable for X-ray diffraction study were obtained.

S3. Refinement

All H atoms were placed in idealized positions (C—H = 0.93 Å) and refined as riding atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of (I), with displacement ellipsoids drawn at the 30% probability level.

4-Chloro-N-(3-phenylallylidene)aniline

Crystal data

C₁₅H₁₂ClN $M_r = 241.71$ Orthorhombic, *Pna*2₁

Hall symbol: P 2c -2n

 $a = 7.7333$ (7) Å $b = 5.5957$ (5) Å $c = 29.383$ (3) Å $V = 1271.5$ (2) Å³ $Z = 4$ $F(000) = 504$ $D_x = 1.263$ Mg m⁻³Mo *K*α radiation, $\lambda = 0.71073$ Å

Cell parameters from 3328 reflections

 $\theta = 2.7$ – 27.2° $\mu = 0.28$ mm⁻¹ $T = 295$ K

Block, yellow

 $0.15 \times 0.12 \times 0.08$ mm

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 2005)

 $T_{\min} = 0.960$, $T_{\max} = 0.978$

6043 measured reflections

2187 independent reflections

2042 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.016$ $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.8^\circ$ $h = -9 \rightarrow 9$ $k = -6 \rightarrow 4$ $l = -32 \rightarrow 35$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.026$ $wR(F^2) = 0.066$ $S = 1.06$

2187 reflections

154 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0375P)^2 + 0.0706P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.001$ $\Delta\rho_{\max} = 0.09$ e Å⁻³ $\Delta\rho_{\min} = -0.15$ e Å⁻³Absolute structure: Flack (1983), 1031 Friedel
pairs

Absolute structure parameter: 0.07 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.65759 (7)	0.44278 (11)	0.969692 (18)	0.08065 (18)
N1	0.87735 (18)	0.5317 (3)	0.77774 (5)	0.0522 (3)
C1	0.7249 (2)	0.4739 (3)	0.91350 (6)	0.0509 (4)

C2	0.8238 (2)	0.6663 (3)	0.90138 (5)	0.0535 (4)
H2	0.8555	0.7795	0.9230	0.064*
C3	0.8761 (2)	0.6907 (3)	0.85660 (5)	0.0503 (4)
H3	0.9443	0.8203	0.8483	0.060*
C4	0.82813 (19)	0.5241 (3)	0.82396 (6)	0.0430 (3)
C5	0.7294 (2)	0.3300 (3)	0.83740 (6)	0.0483 (4)
H5	0.6984	0.2152	0.8160	0.058*
C6	0.6762 (2)	0.3037 (3)	0.88200 (6)	0.0524 (4)
H6	0.6088	0.1737	0.8906	0.063*
C7	0.9240 (2)	0.7278 (3)	0.75959 (6)	0.0499 (4)
H7	0.9253	0.8657	0.7773	0.060*
C8	0.9747 (2)	0.7431 (3)	0.71272 (5)	0.0477 (3)
H8	0.9650	0.6067	0.6948	0.057*
C9	1.0343 (2)	0.9406 (3)	0.69352 (5)	0.0504 (4)
H9	1.0425	1.0744	0.7122	0.060*
C10	1.0884 (2)	0.9711 (3)	0.64651 (5)	0.0421 (3)
C11	1.1805 (2)	1.1742 (3)	0.63356 (5)	0.0493 (4)
H11	1.2065	1.2901	0.6552	0.059*
C12	1.2339 (2)	1.2059 (3)	0.58897 (6)	0.0541 (4)
H12	1.2960	1.3419	0.5810	0.065*
C13	1.1957 (2)	1.0385 (3)	0.55664 (6)	0.0531 (4)
H13	1.2307	1.0608	0.5267	0.064*
C14	1.1043 (2)	0.8352 (3)	0.56881 (5)	0.0505 (4)
H14	1.0791	0.7201	0.5470	0.061*
C15	1.0509 (2)	0.8028 (3)	0.61284 (5)	0.0452 (3)
H15	0.9886	0.6664	0.6204	0.054*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0921 (4)	0.1047 (4)	0.0451 (2)	-0.0037 (3)	0.0100 (2)	0.0165 (3)
N1	0.0539 (8)	0.0588 (8)	0.0439 (8)	-0.0028 (6)	0.0035 (6)	0.0003 (6)
C1	0.0462 (9)	0.0647 (11)	0.0417 (8)	0.0070 (8)	0.0009 (7)	0.0096 (8)
C2	0.0597 (10)	0.0562 (10)	0.0448 (9)	-0.0048 (8)	-0.0048 (7)	-0.0015 (8)
C3	0.0522 (9)	0.0502 (9)	0.0487 (9)	-0.0114 (8)	-0.0012 (7)	0.0045 (8)
C4	0.0383 (7)	0.0467 (8)	0.0441 (8)	0.0020 (6)	-0.0021 (7)	0.0021 (7)
C5	0.0484 (9)	0.0431 (8)	0.0535 (9)	-0.0021 (7)	0.0010 (7)	-0.0011 (7)
C6	0.0496 (9)	0.0503 (9)	0.0573 (10)	-0.0071 (7)	0.0031 (8)	0.0092 (8)
C7	0.0480 (8)	0.0569 (9)	0.0448 (8)	0.0002 (7)	-0.0011 (7)	0.0019 (7)
C8	0.0464 (8)	0.0556 (9)	0.0412 (7)	-0.0002 (7)	-0.0005 (7)	-0.0001 (6)
C9	0.0534 (9)	0.0527 (9)	0.0450 (8)	-0.0019 (7)	-0.0028 (7)	-0.0021 (7)
C10	0.0392 (7)	0.0449 (8)	0.0423 (8)	0.0034 (6)	-0.0041 (6)	0.0052 (6)
C11	0.0499 (9)	0.0444 (8)	0.0537 (9)	-0.0028 (7)	-0.0055 (7)	0.0005 (7)
C12	0.0515 (10)	0.0483 (9)	0.0625 (11)	-0.0021 (7)	0.0037 (8)	0.0143 (8)
C13	0.0553 (10)	0.0565 (10)	0.0477 (10)	0.0064 (7)	0.0035 (7)	0.0099 (8)
C14	0.0584 (10)	0.0500 (9)	0.0432 (9)	0.0057 (7)	-0.0044 (7)	-0.0017 (7)
C15	0.0481 (8)	0.0426 (7)	0.0449 (8)	-0.0032 (7)	-0.0052 (7)	0.0041 (7)

Geometric parameters (Å, °)

C11—C1	1.7399 (17)	C8—C9	1.324 (2)
N1—C7	1.273 (2)	C8—H8	0.9300
N1—C4	1.411 (2)	C9—C10	1.454 (2)
C1—C2	1.368 (2)	C9—H9	0.9300
C1—C6	1.380 (3)	C10—C11	1.394 (2)
C2—C3	1.383 (2)	C10—C15	1.397 (2)
C2—H2	0.9300	C11—C12	1.385 (2)
C3—C4	1.388 (2)	C11—H11	0.9300
C3—H3	0.9300	C12—C13	1.367 (3)
C4—C5	1.385 (2)	C12—H12	0.9300
C5—C6	1.382 (2)	C13—C14	1.386 (3)
C5—H5	0.9300	C13—H13	0.9300
C6—H6	0.9300	C14—C15	1.370 (2)
C7—C8	1.434 (2)	C14—H14	0.9300
C7—H7	0.9300	C15—H15	0.9300
C7—N1—C4	120.37 (14)	C9—C8—H8	118.2
C2—C1—C6	121.41 (16)	C7—C8—H8	118.2
C2—C1—C11	119.54 (14)	C8—C9—C10	127.14 (15)
C6—C1—C11	119.05 (14)	C8—C9—H9	116.4
C1—C2—C3	119.25 (16)	C10—C9—H9	116.4
C1—C2—H2	120.4	C11—C10—C15	117.55 (14)
C3—C2—H2	120.4	C11—C10—C9	120.17 (14)
C2—C3—C4	120.87 (15)	C15—C10—C9	122.27 (14)
C2—C3—H3	119.6	C12—C11—C10	121.00 (15)
C4—C3—H3	119.6	C12—C11—H11	119.5
C5—C4—C3	118.49 (15)	C10—C11—H11	119.5
C5—C4—N1	116.52 (14)	C13—C12—C11	120.35 (16)
C3—C4—N1	124.95 (14)	C13—C12—H12	119.8
C6—C5—C4	121.20 (15)	C11—C12—H12	119.8
C6—C5—H5	119.4	C12—C13—C14	119.57 (15)
C4—C5—H5	119.4	C12—C13—H13	120.2
C5—C6—C1	118.77 (15)	C14—C13—H13	120.2
C5—C6—H6	120.6	C15—C14—C13	120.39 (15)
C1—C6—H6	120.6	C15—C14—H14	119.8
N1—C7—C8	122.08 (15)	C13—C14—H14	119.8
N1—C7—H7	119.0	C14—C15—C10	121.14 (14)
C8—C7—H7	119.0	C14—C15—H15	119.4
C9—C8—C7	123.66 (15)	C10—C15—H15	119.4
C6—C1—C2—C3	0.0 (3)	N1—C7—C8—C9	-175.51 (17)
C11—C1—C2—C3	-179.67 (13)	C7—C8—C9—C10	179.94 (15)
C1—C2—C3—C4	0.6 (2)	C8—C9—C10—C11	-166.99 (16)
C2—C3—C4—C5	-1.3 (2)	C8—C9—C10—C15	13.3 (3)
C2—C3—C4—N1	-178.92 (16)	C15—C10—C11—C12	-0.6 (2)
C7—N1—C4—C5	160.20 (15)	C9—C10—C11—C12	179.69 (15)

C7—N1—C4—C3	-22.1 (2)	C10—C11—C12—C13	0.5 (3)
C3—C4—C5—C6	1.3 (2)	C11—C12—C13—C14	-0.6 (3)
N1—C4—C5—C6	179.18 (14)	C12—C13—C14—C15	0.7 (2)
C4—C5—C6—C1	-0.7 (2)	C13—C14—C15—C10	-0.7 (2)
C2—C1—C6—C5	0.1 (2)	C11—C10—C15—C14	0.7 (2)
C11—C1—C6—C5	179.73 (13)	C9—C10—C15—C14	-179.60 (15)
C4—N1—C7—C8	-179.89 (14)		
