

2,2'-[Nonane-1,9-diylbis(nitrilomethylidyne)]diphenol

Norbani Abdullah, Kong Mun Lo, Hairul Anuar Tajuddin,
Jia Ti Tee and Seik Weng Ng*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 24 November 2008; accepted 26 November 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002 \text{ \AA}$; R factor = 0.040; wR factor = 0.177; data-to-parameter ratio = 18.2.

In the title Schiff base compound, $\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_2$, the complete molecule is generated by crystallographic twofold symmetry, with one C atom lying on the rotation axis. The nonane chain adopts a linear conformation and the hydroxy group forms an intramolecular O—H···N hydrogen bond to the imine group.

Related literature

For the effect of alkyl length on the optical properties of 2,2'-[alkyl-1,9-diylbis(nitrilomethylidyne)]diphenols, see: Kawasaki *et al.* (1996, 1999). For the reduction of the Schiff base to the secondary diamine, see: Csaszar (1984). For the structure of 2,2'-[hexane-1,6-diylbis(nitrilomethylidyne)]diphenol, see: Sheikhshoae & Sharif (2006).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_2$
 $M_r = 366.49$
Monoclinic, $C2/c$

$a = 43.6905 (10) \text{ \AA}$
 $b = 4.7258 (1) \text{ \AA}$
 $c = 9.8928 (2) \text{ \AA}$

$\beta = 96.935 (2)^\circ$
 $V = 2027.65 (8) \text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.08 \text{ mm}^{-1}$
 $T = 100 (2) \text{ K}$
 $0.40 \times 0.03 \times 0.02 \text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: none
8930 measured reflections

2317 independent reflections
1573 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.177$
 $S = 1.10$
2317 reflections
127 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1···N1	0.86 (1)	1.81 (2)	2.5755 (19)	148 (3)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the Fundamental Research Grant Scheme of the Ministry of Higher Education for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2862).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Csaszar, J. (1984). *Acta Phys. Chem.* **30**, 61.
- Kawasaki, T., Kamata, T., Ushijima, H., Kanakubo, M., Murata, S., Mizukami, F., Fujii, Y. & Usui, Y. (1999). *J. Chem. Soc. Perkin Trans. 2*, pp. 193–198.
- Kawasaki, T., Kamata, T., Ushijima, H., Murata, S., Mizukami, F., Fujii, Y. & Usui, Y. (1996). *Mol. Cryst. Liq. Cryst.* **286**, 579–584.
- Sheikhshoae, I. & Sharif, M. A. (2006). *Acta Cryst. E* **62**, o3563–o3565.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2008). *publCIF*. In preparation.

supporting information

Acta Cryst. (2008). E64, o2494 [doi:10.1107/S1600536808039858]

2,2'-[Nonane-1,9-diylbis(nitrilomethylidyne)]diphenol

Norbani Abdullah, Kong Mun Lo, Hairul Anuar Tajuddin, Jia Ti Tee and Seik Weng Ng

S1. Comment

For more details, see the Abstract. For the molecular structure, see Fig. 1. and for details of hydrogen bonding, see Table 1.

S2. Experimental

Salicylaldehyde (0.050 mol, 6.1 g) and sodium hydroxide (0.05 mol, 2.0 g) in methanol (125 ml) was added to 1,9-diaminononane (0.025 mol, 3.9 g) in methanol (125 ml). The solution was heated for 1 h. The solvent was evaporated and the product recrystallized from ethanol to yield yellow plates of (I). The rod used for data collection was cut from a plate.

S3. Refinement

The C-bound hydrogen atoms were placed at calculated positions ($C-H = 0.95-0.99 \text{ \AA}$) and refined as riding with $U_{\text{iso}}(H) = 1.2U_{\text{eq}}(C)$. The hydroxy H-atom was located in a difference Fourier map and was refined with a distance restraint of $O-H = 0.84 \pm 0.01 \text{ \AA}$.




Figure 1

View of the molecular structure of (I) at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius. The unlabelled atoms are generated by the symmetry operation $(1-x, y, 1/2-z)$.

2,2'-[Nonane-1,9-diylbis(nitrilomethylidyne)]diphenol

Crystal data

$C_{23}H_{30}N_2O_2$
 $M_r = 366.49$
Monoclinic, $C2/c$
Hall symbol: -C 2yc
 $a = 43.6905 (10) \text{ \AA}$
 $b = 4.7258 (1) \text{ \AA}$

$c = 9.8928 (2) \text{ \AA}$
 $\beta = 96.935 (2)^\circ$
 $V = 2027.65 (8) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 792$
 $D_x = 1.201 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 2058 reflections
 $\theta = 2.8\text{--}27.9^\circ$
 $\mu = 0.08 \text{ mm}^{-1}$

$T = 100 \text{ K}$
 Rod, yellow
 $0.40 \times 0.03 \times 0.02 \text{ mm}$

Data collection

Bruker SMART APEX CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 8930 measured reflections
 2317 independent reflections

1573 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 0.9^\circ$
 $h = -56 \rightarrow 56$
 $k = -6 \rightarrow 6$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.177$
 $S = 1.10$
 2317 reflections
 127 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.1065P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.63065 (3)	0.8920 (3)	0.86876 (12)	0.0364 (4)	
H1	0.6186 (5)	0.787 (5)	0.816 (2)	0.075 (8)*	
N1	0.61425 (3)	0.4953 (3)	0.69663 (13)	0.0283 (4)	
C1	0.66008 (4)	0.8348 (4)	0.84965 (15)	0.0300 (4)	
C2	0.68388 (4)	0.9845 (4)	0.92417 (17)	0.0382 (5)	
H2	0.6793	1.1251	0.9874	0.046*	
C3	0.71405 (4)	0.9284 (4)	0.90596 (19)	0.0410 (5)	
H3	0.7302	1.0303	0.9577	0.049*	
C4	0.72134 (4)	0.7262 (4)	0.81367 (19)	0.0388 (5)	
H4	0.7422	0.6916	0.8009	0.047*	
C5	0.69785 (4)	0.5759 (4)	0.74073 (18)	0.0344 (4)	
H5	0.7027	0.4358	0.6779	0.041*	
C6	0.66712 (4)	0.6252 (3)	0.75711 (15)	0.0281 (4)	
C7	0.64264 (4)	0.4541 (4)	0.68374 (15)	0.0282 (4)	
H7	0.6480	0.3082	0.6249	0.034*	
C8	0.59137 (4)	0.3110 (4)	0.62109 (16)	0.0295 (4)	
H8A	0.5792	0.2140	0.6854	0.035*	
H8B	0.6020	0.1649	0.5720	0.035*	
C9	0.56994 (4)	0.4833 (4)	0.51955 (16)	0.0290 (4)	
H9A	0.5590	0.6250	0.5696	0.035*	
H9B	0.5824	0.5867	0.4584	0.035*	

C10	0.54637 (4)	0.2996 (4)	0.43422 (16)	0.0295 (4)
H10A	0.5351	0.1845	0.4957	0.035*
H10B	0.5573	0.1683	0.3786	0.035*
C11	0.52325 (3)	0.4723 (4)	0.34047 (16)	0.0278 (4)
H11A	0.5347	0.5929	0.2818	0.033*
H11B	0.5119	0.5990	0.3968	0.033*
C12	0.5000	0.2941 (5)	0.2500	0.0291 (5)
H12A	0.4888	0.1707	0.3082	0.035*
H12B	0.5112	0.1707	0.1918	0.035*
				0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0363 (8)	0.0429 (8)	0.0299 (7)	0.0016 (6)	0.0041 (5)	-0.0045 (5)
N1	0.0256 (8)	0.0366 (8)	0.0213 (7)	0.0014 (6)	-0.0026 (5)	0.0018 (6)
C1	0.0343 (10)	0.0354 (9)	0.0194 (7)	0.0001 (7)	-0.0007 (7)	0.0058 (7)
C2	0.0489 (12)	0.0379 (10)	0.0259 (8)	-0.0070 (8)	-0.0036 (8)	0.0002 (8)
C3	0.0395 (11)	0.0436 (11)	0.0361 (10)	-0.0118 (8)	-0.0115 (8)	0.0069 (8)
C4	0.0286 (10)	0.0441 (11)	0.0415 (10)	-0.0031 (8)	-0.0053 (8)	0.0083 (8)
C5	0.0307 (10)	0.0396 (10)	0.0318 (9)	0.0013 (7)	-0.0006 (7)	0.0039 (7)
C6	0.0298 (9)	0.0331 (9)	0.0204 (7)	0.0006 (7)	-0.0018 (6)	0.0055 (6)
C7	0.0281 (9)	0.0352 (9)	0.0204 (7)	0.0029 (7)	-0.0008 (6)	0.0019 (6)
C8	0.0251 (9)	0.0352 (9)	0.0269 (8)	0.0007 (7)	-0.0017 (7)	0.0005 (7)
C9	0.0238 (9)	0.0354 (9)	0.0266 (8)	0.0014 (7)	-0.0018 (7)	0.0002 (7)
C10	0.0238 (9)	0.0334 (9)	0.0302 (8)	0.0020 (6)	-0.0012 (7)	-0.0005 (7)
C11	0.0231 (8)	0.0332 (9)	0.0265 (8)	0.0013 (6)	0.0005 (6)	-0.0006 (6)
C12	0.0230 (12)	0.0327 (12)	0.0306 (11)	0.000	-0.0010 (9)	0.000

Geometric parameters (\AA , $^\circ$)

O1—C1	1.349 (2)	C8—C9	1.524 (2)
O1—H1	0.86 (1)	C8—H8A	0.9900
N1—C7	1.277 (2)	C8—H8B	0.9900
N1—C8	1.461 (2)	C9—C10	1.522 (2)
C1—C2	1.393 (2)	C9—H9A	0.9900
C1—C6	1.408 (2)	C9—H9B	0.9900
C2—C3	1.377 (3)	C10—C11	1.523 (2)
C2—H2	0.9500	C10—H10A	0.9900
C3—C4	1.385 (3)	C10—H10B	0.9900
C3—H3	0.9500	C11—C12	1.524 (2)
C4—C5	1.378 (2)	C11—H11A	0.9900
C4—H4	0.9500	C11—H11B	0.9900
C5—C6	1.391 (2)	C12—C11 ⁱ	1.524 (2)
C5—H5	0.9500	C12—H12A	0.9900
C6—C7	1.462 (2)	C12—H12B	0.9900
C7—H7	0.9500		
C1—O1—H1	109.1 (19)	C9—C8—H8B	109.6

C7—N1—C8	118.06 (14)	H8A—C8—H8B	108.1
O1—C1—C2	119.15 (16)	C10—C9—C8	112.43 (14)
O1—C1—C6	121.25 (15)	C10—C9—H9A	109.1
C2—C1—C6	119.60 (16)	C8—C9—H9A	109.1
C3—C2—C1	119.86 (17)	C10—C9—H9B	109.1
C3—C2—H2	120.1	C8—C9—H9B	109.1
C1—C2—H2	120.1	H9A—C9—H9B	107.9
C2—C3—C4	121.26 (17)	C9—C10—C11	112.75 (14)
C2—C3—H3	119.4	C9—C10—H10A	109.0
C4—C3—H3	119.4	C11—C10—H10A	109.0
C5—C4—C3	118.99 (18)	C9—C10—H10B	109.0
C5—C4—H4	120.5	C11—C10—H10B	109.0
C3—C4—H4	120.5	H10A—C10—H10B	107.8
C4—C5—C6	121.38 (17)	C10—C11—C12	114.05 (15)
C4—C5—H5	119.3	C10—C11—H11A	108.7
C6—C5—H5	119.3	C12—C11—H11A	108.7
C5—C6—C1	118.89 (15)	C10—C11—H11B	108.7
C5—C6—C7	120.52 (15)	C12—C11—H11B	108.7
C1—C6—C7	120.54 (15)	H11A—C11—H11B	107.6
N1—C7—C6	121.81 (15)	C11—C12—C11 ⁱ	112.9 (2)
N1—C7—H7	119.1	C11—C12—H12A	109.0
C6—C7—H7	119.1	C11 ⁱ —C12—H12A	109.0
N1—C8—C9	110.26 (14)	C11—C12—H12B	109.0
N1—C8—H8A	109.6	C11 ⁱ —C12—H12B	109.0
C9—C8—H8A	109.6	H12A—C12—H12B	107.8
N1—C8—H8B	109.6		
O1—C1—C2—C3	179.92 (15)	C2—C1—C6—C7	176.40 (14)
C6—C1—C2—C3	0.6 (2)	C8—N1—C7—C6	-179.01 (14)
C1—C2—C3—C4	0.5 (3)	C5—C6—C7—N1	-179.81 (16)
C2—C3—C4—C5	-1.1 (3)	C1—C6—C7—N1	2.8 (2)
C3—C4—C5—C6	0.6 (3)	C7—N1—C8—C9	-117.30 (16)
C4—C5—C6—C1	0.5 (2)	N1—C8—C9—C10	177.84 (13)
C4—C5—C6—C7	-176.97 (15)	C8—C9—C10—C11	175.42 (14)
O1—C1—C6—C5	179.62 (14)	C9—C10—C11—C12	177.98 (12)
C2—C1—C6—C5	-1.0 (2)	C10—C11—C12—C11 ⁱ	178.76 (15)
O1—C1—C6—C7	-3.0 (2)		

Symmetry code: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^{\circ}$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1 \cdots N1	0.86 (1)	1.81 (2)	2.5755 (19)	148 (3)