

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[bis[3,10-bis(2-hydroxyethyl)-1,3,5,8,10,12-hexaazacyclotetradecane]tetra- μ -cyanido-tetracyanidodicycopper(II)molybdenum(IV)] tetrahydrate

Hu Zhou, Ying-Ying Chen, Wen-Yan Liu and Ai-Hua Yuan*

School of Materials Science and Engineering, Jiangsu University of Science and Technology, Zhenjiang 212003, People's Republic of China
Correspondence e-mail: aihuayuan@163.com


Received 11 October 2008; accepted 4 November 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.043; wR factor = 0.103; data-to-parameter ratio = 18.5.

In the title complex, $[[\text{Cu}_2\text{Mo}(\text{CN})_8(\text{C}_{12}\text{H}_{30}\text{N}_6\text{O}_2)_2]\cdot 4\text{H}_2\text{O}]_n$, the polyhedron around Mo has site symmetry $\bar{1}$ with a distorted square-antiprismatic shape, while the Cu atom (2 symmetry) is in a distorted axially elongated octahedral coordination environment. The uncoordinated water molecule is disordered over three sites with occupancies of 0.445 (7), 0.340 (7) and 0.215 (7). Mo and Cu atoms acting as basic components are connected by an Mo—CN—Cu—NC—Mo—linkage to form a distorted diamond-like network. Additional hydrogen bonding between the N—H groups and the water molecules stabilizes this arrangement.

Related literature

For background information, see: Larionova *et al.* (2004); Przychodzeń *et al.* (2006). For related structures, see: Chen *et al.* (2007); Zhou *et al.* (2007; 2008). For literature related to the synthesis, see: Suh & Kang (1988); Leipoldt *et al.* (1974).



Experimental

Crystal data

$[\text{Cu}_2\text{Mo}(\text{CN})_8(\text{C}_{12}\text{H}_{30}\text{N}_6\text{O}_2)_2]\cdot 4\text{H}_2\text{O}$ $Z = 4$
 $M_r = 1084.08$ Mo $K\alpha$ radiation
 Tetragonal, $I4_1/a$ $\mu = 0.93$ mm⁻¹
 $a = 20.0707$ (10) Å $T = 293$ (2) K
 $c = 15.4380$ (17) Å $0.32 \times 0.28 \times 0.26$ mm
 $V = 6218.9$ (8) Å³

Data collection

Bruker SMART APEXII diffractometer 16289 measured reflections
 Absorption correction: multi-scan 3053 independent reflections
 (SADABS; Bruker, 2002) 2459 reflections with $I > 2\sigma(I)$
 $T_{\min} = 0.75$, $T_{\max} = 0.79$ $R_{\text{int}} = 0.058$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$ 1 restraint
 $wR(F^2) = 0.103$ H-atom parameters constrained
 $S = 1.06$ $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 3053 reflections $\Delta\rho_{\text{min}} = -0.38$ e Å⁻³
 165 parameters

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H3C}\cdots\text{N2}^{\text{i}}$	0.93	2.44	3.261 (4)	147
$\text{N5}-\text{H5C}\cdots\text{O2}^{\text{ii}}$	0.93	2.38	3.291 (8)	165
$\text{O1}-\text{H1C}\cdots\text{O4}^{\text{iii}}$	0.85	2.48	3.149 (7)	136
$\text{O4}-\text{H4D}\cdots\text{N2}^{\text{iv}}$	0.85	2.11	2.729 (6)	129
$\text{O4}-\text{H4E}\cdots\text{O2}^{\text{v}}$	0.85	2.47	3.279 (9)	159

Symmetry codes: (i) $-x + 2, -y + 1, -z$; (ii) $y + \frac{3}{4}, -x + \frac{3}{4}, z - \frac{1}{4}$; (iii) $-y + \frac{3}{4}, x + \frac{1}{4}, -z + \frac{3}{4}$; (iv) $x - 1, y, z + 1$; (v) $y - \frac{1}{4}, -x + \frac{1}{4}, -z + \frac{3}{4}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work is supported by the University Natural Science Foundation of Jiangsu Province (grant No. 07KJB150030).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2114).

References

- Bruker (2002). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chen, Y. Y., Zhou, H., Shen, X. P., Lu, H. F. & Yuan, A. H. (2007). *J. Mol. Struct.* **839**, 64–68.
 Larionova, J., Willemin, S., Donnadiou, B., Henner, B., Guérin, C., Gillon, B. & Goujon, A. (2004). *J. Phys. Chem. Solids*, **65**, 677–691.
 Leipoldt, J. G., Bok, L. D. C. & Cillier, P. J. (1974). *Z. Anorg. Allg. Chem.* **409**, 343–344.
 Przychodzeń, P., Korzeniak, T., Podgajny, R. & Sieklucka, B. (2006). *Coord. Chem. Rev.* **250**, 2234–2260.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Suh, M. P. & Kang, S. G. (1988). *Inorg. Chem.* **27**, 2544–2546.
 Zhou, H., Chen, Y. Y., Yuan, A. H. & Shen, X. P. (2008). *Inorg. Chem. Commun.* **11**, 363–366.
 Zhou, H., Yuan, A. H., Shen, X. P., Chen, Y. Y., Price, D. J. & Kepert, C. J. (2007). *Inorg. Chem. Commun.* **10**, 940–943.

supporting information

Acta Cryst. (2008). E64, m1528 [doi:10.1107/S1600536808036131]

Poly[bis[3,10-bis(2-hydroxyethyl)-1,3,5,8,10,12-hexaazacyclotetradecane]tetra- μ -cyanido-tetracyanidodicopper(II)molybdenum(IV)] tetrahydrate]

Hu Zhou, Ying-Ying Chen, Wen-Yan Liu and Ai-Hua Yuan

S1. Comment

Recently, octacyanometalates $[M(\text{CN})_8]^{3-/4-}$ ($M = \text{Mo}, \text{W}$ and Nb) have been found to be versatile building blocks and investigated extensively (Larionova *et al.*, 2004; Przychodzeń *et al.*, 2006). For $[\text{CuL}]^{2+}/[M(\text{CN})_8]^{3-/4-}$ ($L =$ macrocyclic ligands) bimetallic systems, macrocyclic ligands block partly coordination sites of the metal ions and release their apical ones, which can be used to construct cyano-bridged bimetallic complexes with octacyanometalate $[M(\text{CN})_8]^{3-/4-}$ ions. In a recent study, we chose $[\text{CuL}]^{2+}$ ($L = 3,10$ -diethanol-1,3,5,8,10,12-hexaazacyclotetradecane) as building block to synthesize successfully an octacyanometalate-based bimetallic complex $[\text{CuL}]_2[\text{Mo}(\text{CN})_8] \cdot 4\text{H}_2\text{O}$, (I), with a distorted diamond network.

The title complex crystallizes in the tetragonal ($I4_1/a$) space group. As displayed in Fig. 1, the Mo atom is coordinated by eight CN groups in a distorted square antiprism. The Cu atom is in a distorted axially elongated octahedral coordination environment, in which four nitrogen atoms from the ligand (L) occupy the equatorial positions, while the axial sites are occupied by two nitrogen atoms from the bridging cyanide groups on different $[\text{Mo}(\text{CN})_8]^{4-}$ anions. The bonding parameters of the macrocyclic ligand L are reminiscent of those found in related complexes reported previously (Chen *et al.*, 2007).

As shown in Fig. 2, Mo and Cu atoms acting as basic components (linker and connector, respectively) are connected by the Mo—CN—Cu—NC—Mo— linkages to form a three-dimensional structure. The network is composed of $[\text{CuL}]^{2+}$ unit that is linked *via* cyanides to adjacent four-connected $[\text{Mo}(\text{CN})_8(\mu\text{-CN})_4]^{4-}$ centers. From a topological standpoint, each $[\text{Mo}(\text{CN})_8]$ unit is a tetrahedral four-connecting node. These nodes are linked to four adjacent $[\text{Mo}(\text{CN})_8]$ units by the $[\text{CuL}]$ units, acting as linear two-connectors. The result is a distorted diamond network (Fig. 2), which is similar to octacyanometalate-based bimetallic complexes reported previously (Zhou *et al.*, 2007; 2008).

S2. Experimental

Well shaped brown crystals of the title complex suitable for X-ray single-crystal structure determination were grown at room temperature by slow diffusion of a DMF solution (15 ml) of $[\text{CuL}](\text{ClO}_4)_2$ (0.30 mmol) (Suh and Kang, 1988) and an aqueous solution (15 ml) of $\text{K}_4[\text{Mo}(\text{CN})_8] \cdot 2\text{H}_2\text{O}$ (0.15 mmol) (Leipoldt *et al.*, 1974) in a U-shaped tube containing agar for about one month. The resulting crystals were collected, washed with H_2O and dried in air.

S3. Refinement

All hydrogen atoms except for hydrogen atoms bound to water molecules were calculated at idealized positions with C—H = 0.99, N—H = 0.93 and O—H = 0.85 Å and included in the refinement in a riding mode with U_{iso} for H assigned as 1.2 times U_{eq} of the attached atoms. The O atom of the water of hydration molecule was disordered over three sites with unequal site occupancy factors at locations O2, O3 and O4. The H atoms of the disordered water molecule were located

from difference maps and were included in the refinements at geometrically idealized positions with O–H distances 0.85 Å and U_{iso} assigned as 1.2 times U_{eq} of the attached O atoms.





Figure 1

ORTEP drawing of the title complex, showing the atom labeling. Hydrogen atoms and water molecules are omitted for clarity. Symmetry codes: A: $2-x, 1-y, -z$; B: $2-x, 0.5-y, z$; C: $1.25-y, -0.75+x, 0.25-z$.

**Figure 2**

Topological depiction of the title complex, where the Mo nodes and the Cu connectors are shown in pink and green colors, respectively.

Poly[[bis[3,10-bis(2-hydroxyethyl)-1,3,5,8,10,12-hexaazacyclotetradecane]tetra- μ -cyanido-tetracyanidodicopper(II)molybdenum(IV)] tetrahydrate]

Crystal data

$[\text{Cu}_2\text{Mo}(\text{CN})_8(\text{C}_{12}\text{H}_{30}\text{N}_6\text{O}_2)_2] \cdot 4\text{H}_2\text{O}$

$M_r = 1084.08$

Tetragonal, $I4_1/a$

Hall symbol: $-I\ 4ad$

$a = 20.0707$ (10) Å

$c = 15.4380$ (17) Å

$V = 6218.9$ (8) Å³

$Z = 4$

$F(000) = 2256$

$D_x = 1.158$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3093 reflections

$\theta = 2.6\text{--}21.2^\circ$

$\mu = 0.93$ mm⁻¹

$T = 293$ K

Block, brown

$0.32 \times 0.28 \times 0.26$ mm

Data collection

Bruker SMART APEXII
diffractometer

Radiation source: sealed tube

Graphite monochromator

ϕ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2002)

$T_{\min} = 0.75$, $T_{\max} = 0.79$

16289 measured reflections

3053 independent reflections

2459 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.058$

$\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -24 \rightarrow 24$

$k = -19 \rightarrow 24$

$l = -16 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.103$
 $S = 1.06$
 3053 reflections
 165 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0515P)^2 + 3.1936P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.38 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.96688 (13)	0.34611 (15)	0.08263 (19)	0.0392 (6)	
C2	1.06251 (16)	0.27104 (17)	0.0121 (2)	0.0485 (7)	
C3	0.86789 (19)	0.5451 (2)	0.0135 (2)	0.0615 (9)	
H3A	0.8364	0.5817	0.0269	0.074*	
H3B	0.8476	0.5025	0.0322	0.074*	
C4	0.92199 (19)	0.5426 (2)	0.1513 (2)	0.0610 (9)	
H4A	0.8863	0.5716	0.1746	0.073*	
H4B	0.9081	0.4957	0.1597	0.073*	
C5	1.0360 (2)	0.5073 (2)	0.1850 (2)	0.0627 (10)	
H5A	1.0195	0.4618	0.1971	0.075*	
H5B	1.0715	0.5172	0.2277	0.075*	
C6	1.1181 (2)	0.4567 (2)	0.0813 (2)	0.0649 (10)	
H6A	1.1023	0.4126	0.1013	0.078*	
H6B	1.1591	0.4682	0.1137	0.078*	
C7	0.99694 (18)	0.62268 (18)	0.2106 (2)	0.0570 (9)	
H7A	0.9623	0.6491	0.1805	0.068*	
H7B	1.0397	0.6317	0.1808	0.068*	
C8	1.0028 (2)	0.6495 (2)	0.3018 (3)	0.0694 (11)	
H8A	1.0505	0.6519	0.3175	0.083*	
H8B	0.9848	0.6954	0.3030	0.083*	
Cu2	1.0000	0.5000	0.0000	0.04929 (18)	
Mo1	1.0000	0.2500	0.1250	0.03165 (14)	
N1	0.94948 (13)	0.39753 (13)	0.05790 (17)	0.0467 (6)	
N2	1.09533 (14)	0.28414 (15)	-0.04505 (19)	0.0554 (7)	
N3	0.92993 (13)	0.55566 (15)	0.05812 (19)	0.0539 (7)	

H3C	0.9417	0.6002	0.0514	0.065*	
N4	0.98158 (17)	0.55421 (16)	0.1980 (3)	0.0737 (9)	
N5	1.06603 (14)	0.50813 (15)	0.09613 (16)	0.0500 (7)	
H5C	1.0866	0.5493	0.0896	0.060*	
O1	0.9722 (2)	0.6148 (2)	0.3594 (3)	0.1107 (12)	
H1C	0.9917	0.5774	0.3651	0.133*	
O2	0.1088 (4)	0.4119 (4)	0.3446 (5)	0.072 (3)	0.340 (7)
H2A	0.0857	0.4473	0.3454	0.087*	0.340 (7)
H2B	0.0994	0.3893	0.3896	0.087*	0.340 (7)
O3	0.2794 (8)	0.5343 (7)	0.0656 (10)	0.088 (6)	0.215 (7)
H3D	0.2957	0.5708	0.0836	0.105*	0.215 (7)
H3E	0.3107	0.5076	0.0526	0.105*	0.215 (7)
O4	0.2103 (3)	0.2935 (3)	0.8613 (3)	0.065 (2)	0.445 (7)
H4D	0.1814	0.3166	0.8882	0.078*	0.445 (7)
H4E	0.1970	0.2534	0.8577	0.078*	0.445 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0211 (12)	0.0461 (17)	0.0503 (17)	0.0025 (11)	-0.0005 (11)	0.0068 (13)
C2	0.0437 (17)	0.0534 (19)	0.0485 (18)	-0.0006 (14)	0.0066 (14)	-0.0020 (15)
C3	0.058 (2)	0.064 (2)	0.063 (2)	-0.0125 (18)	-0.0127 (17)	0.0055 (18)
C4	0.058 (2)	0.083 (3)	0.0428 (19)	-0.0050 (19)	0.0110 (15)	0.0042 (17)
C5	0.068 (2)	0.079 (3)	0.0419 (18)	0.0000 (19)	-0.0184 (17)	-0.0064 (17)
C6	0.072 (2)	0.079 (3)	0.044 (2)	-0.007 (2)	-0.0283 (18)	0.0000 (17)
C7	0.060 (2)	0.056 (2)	0.0550 (19)	-0.0147 (16)	0.0152 (16)	-0.0383 (16)
C8	0.071 (2)	0.068 (2)	0.069 (3)	-0.003 (2)	0.019 (2)	-0.024 (2)
Cu2	0.0514 (3)	0.0560 (3)	0.0405 (3)	-0.0020 (3)	-0.0059 (2)	0.0008 (2)
Mo1	0.01805 (15)	0.01805 (15)	0.0589 (3)	0.000	0.000	0.000
N1	0.0459 (14)	0.0452 (14)	0.0490 (15)	0.0031 (11)	-0.0036 (12)	0.0120 (12)
N2	0.0549 (17)	0.0586 (17)	0.0528 (17)	-0.0083 (14)	0.0019 (14)	0.0287 (14)
N3	0.0405 (14)	0.0671 (18)	0.0540 (16)	0.0017 (13)	0.0018 (12)	0.0123 (14)
N4	0.071 (2)	0.0478 (18)	0.103 (3)	-0.0030 (15)	0.018 (2)	-0.0012 (17)
N5	0.0521 (16)	0.0572 (17)	0.0406 (14)	0.0061 (13)	-0.0163 (12)	-0.0079 (12)
O1	0.115 (3)	0.113 (3)	0.104 (3)	0.004 (2)	0.011 (2)	0.006 (2)
O2	0.060 (5)	0.092 (6)	0.065 (5)	0.023 (4)	-0.024 (4)	-0.028 (4)
O3	0.097 (11)	0.063 (8)	0.103 (11)	-0.006 (7)	0.041 (9)	-0.006 (8)
O4	0.064 (4)	0.091 (5)	0.041 (3)	-0.014 (3)	0.016 (2)	0.009 (3)

Geometric parameters (Å, °)

C1—N1	1.154 (4)	C8—O1	1.285 (5)
C1—Mo1	2.143 (3)	C8—H8A	0.9900
C2—N2	1.132 (4)	C8—H8B	0.9900
C2—Mo1	2.188 (3)	Cu2—N5 ⁱ	1.996 (2)
C3—N3	1.439 (4)	Cu2—N5	1.996 (2)
C3—C6 ⁱ	1.490 (5)	Cu2—N3	2.008 (3)
C3—H3A	0.9900	Cu2—N3 ⁱ	2.008 (3)

C3—H3B	0.9900	Cu2—N1 ⁱ	2.461 (3)
C4—N4	1.416 (5)	Cu2—N1	2.461 (3)
C4—N3	1.470 (5)	Mo1—C1 ⁱⁱ	2.143 (3)
C4—H4A	0.9900	Mo1—C1 ⁱⁱⁱ	2.143 (3)
C4—H4B	0.9900	Mo1—C1 ^{iv}	2.143 (3)
C5—N4	1.457 (5)	Mo1—C2 ⁱⁱⁱ	2.188 (3)
C5—N5	1.499 (5)	Mo1—C2 ^{iv}	2.188 (3)
C5—H5A	0.9900	Mo1—C2 ⁱⁱ	2.188 (3)
C5—H5B	0.9900	N3—H3C	0.9300
C6—N5	1.487 (5)	N5—H5C	0.9300
C6—C3 ⁱ	1.490 (5)	O1—H1C	0.8501
C6—H6A	0.9900	O2—H2A	0.8500
C6—H6B	0.9900	O2—H2B	0.8500
C7—N4	1.422 (4)	O3—H3D	0.8500
C7—C8	1.512 (5)	O3—H3E	0.8500
C7—H7A	0.9900	O4—H4D	0.8498
C7—H7B	0.9900	O4—H4E	0.8498
N1—C1—Mo1	178.4 (3)	N3 ⁱ —Cu2—N1	89.19 (10)
N2—C2—Mo1	177.5 (3)	N1 ⁱ —Cu2—N1	180.00 (11)
N3—C3—C6 ⁱ	108.1 (3)	C1 ⁱⁱ —Mo1—C1 ⁱⁱⁱ	95.35 (5)
N3—C3—H3A	110.1	C1 ⁱⁱ —Mo1—C1	144.45 (16)
C6 ⁱ —C3—H3A	110.1	C1 ⁱⁱⁱ —Mo1—C1	95.35 (5)
N3—C3—H3B	110.1	C1 ⁱⁱ —Mo1—C1 ^{iv}	95.35 (5)
C6 ⁱ —C3—H3B	110.1	C1 ⁱⁱⁱ —Mo1—C1 ^{iv}	144.45 (16)
H3A—C3—H3B	108.4	C1—Mo1—C1 ^{iv}	95.35 (5)
N4—C4—N3	112.2 (3)	C1 ⁱⁱ —Mo1—C2 ⁱⁱⁱ	145.00 (12)
N4—C4—H4A	109.2	C1 ⁱⁱⁱ —Mo1—C2 ⁱⁱⁱ	76.18 (12)
N3—C4—H4A	109.2	C1—Mo1—C2 ⁱⁱⁱ	70.56 (12)
N4—C4—H4B	109.2	C1 ^{iv} —Mo1—C2 ⁱⁱⁱ	75.69 (12)
N3—C4—H4B	109.2	C1 ⁱⁱ —Mo1—C2 ^{iv}	70.56 (12)
H4A—C4—H4B	107.9	C1 ⁱⁱⁱ —Mo1—C2 ^{iv}	75.69 (12)
N4—C5—N5	114.8 (3)	C1—Mo1—C2 ^{iv}	144.99 (12)
N4—C5—H5A	108.6	C1 ^{iv} —Mo1—C2 ^{iv}	76.18 (12)
N5—C5—H5A	108.6	C2 ⁱⁱⁱ —Mo1—C2 ^{iv}	74.44 (17)
N4—C5—H5B	108.6	C1 ⁱⁱ —Mo1—C2	75.69 (12)
N5—C5—H5B	108.6	C1 ⁱⁱⁱ —Mo1—C2	144.99 (12)
H5A—C5—H5B	107.5	C1—Mo1—C2	76.17 (12)
N5—C6—C3 ⁱ	107.5 (3)	C1 ^{iv} —Mo1—C2	70.55 (12)
N5—C6—H6A	110.2	C2 ⁱⁱⁱ —Mo1—C2	129.35 (11)
C3 ⁱ —C6—H6A	110.2	C2 ^{iv} —Mo1—C2	129.35 (11)
N5—C6—H6B	110.2	C1 ⁱⁱ —Mo1—C2 ⁱⁱ	76.17 (12)
C3 ⁱ —C6—H6B	110.2	C1 ⁱⁱⁱ —Mo1—C2 ⁱⁱ	70.55 (12)
H6A—C6—H6B	108.5	C1—Mo1—C2 ⁱⁱ	75.69 (12)
N4—C7—C8	119.2 (4)	C1 ^{iv} —Mo1—C2 ⁱⁱ	144.99 (12)
N4—C7—H7A	107.5	C2 ⁱⁱⁱ —Mo1—C2 ⁱⁱ	129.36 (11)
C8—C7—H7A	107.5	C2 ^{iv} —Mo1—C2 ⁱⁱ	129.35 (11)
N4—C7—H7B	107.5	C2—Mo1—C2 ⁱⁱ	74.44 (17)

C8—C7—H7B	107.5	C1—N1—Cu2	137.9 (2)
H7A—C7—H7B	107.0	C3—N3—C4	110.4 (3)
O1—C8—C7	114.5 (4)	C3—N3—Cu2	108.1 (2)
O1—C8—H8A	108.6	C4—N3—Cu2	114.4 (2)
C7—C8—H8A	108.6	C3—N3—H3C	107.9
O1—C8—H8B	108.6	C4—N3—H3C	107.9
C7—C8—H8B	108.6	Cu2—N3—H3C	107.9
H8A—C8—H8B	107.6	C4—N4—C7	114.3 (3)
N5 ⁱ —Cu2—N5	180.0	C4—N4—C5	117.2 (3)
N5 ⁱ —Cu2—N3	84.99 (12)	C7—N4—C5	118.8 (3)
N5—Cu2—N3	95.01 (12)	C6—N5—C5	114.6 (3)
N5 ⁱ —Cu2—N3 ⁱ	95.01 (12)	C6—N5—Cu2	107.20 (19)
N5—Cu2—N3 ⁱ	84.99 (12)	C5—N5—Cu2	114.4 (2)
N3—Cu2—N3 ⁱ	180.0	C6—N5—H5C	106.7
N5 ⁱ —Cu2—N1 ⁱ	94.12 (11)	C5—N5—H5C	106.7
N5—Cu2—N1 ⁱ	85.88 (11)	Cu2—N5—H5C	106.7
N3—Cu2—N1 ⁱ	89.19 (10)	C8—O1—H1C	109.3
N3 ⁱ —Cu2—N1 ⁱ	90.81 (10)	H2A—O2—H2B	108.2
N5 ⁱ —Cu2—N1	85.88 (11)	H3D—O3—H3E	109.5
N5—Cu2—N1	94.12 (11)	H4D—O4—H4E	109.5
N3—Cu2—N1	90.81 (10)		
N4—C7—C8—O1	-22.2 (6)	N3—C4—N4—C5	-70.7 (4)
N5 ⁱ —Cu2—N1—C1	129.7 (4)	C8—C7—N4—C4	120.0 (4)
N5—Cu2—N1—C1	-50.3 (4)	C8—C7—N4—C5	-94.9 (4)
N3—Cu2—N1—C1	-145.4 (4)	N5—C5—N4—C4	66.8 (4)
N3 ⁱ —Cu2—N1—C1	34.6 (4)	N5—C5—N4—C7	-77.3 (4)
C6 ⁱ —C3—N3—C4	-166.9 (3)	C3 ⁱ —C6—N5—C5	168.3 (3)
C6 ⁱ —C3—N3—Cu2	-41.1 (3)	C3 ⁱ —C6—N5—Cu2	40.2 (3)
N4—C4—N3—C3	-178.4 (3)	N4—C5—N5—C6	-175.7 (3)
N4—C4—N3—Cu2	59.5 (4)	N4—C5—N5—Cu2	-51.3 (4)
N5 ⁱ —Cu2—N3—C3	14.8 (2)	N3—Cu2—N5—C6	165.6 (2)
N5—Cu2—N3—C3	-165.2 (2)	N3 ⁱ —Cu2—N5—C6	-14.4 (2)
N1 ⁱ —Cu2—N3—C3	109.0 (2)	N1 ⁱ —Cu2—N5—C6	-105.5 (2)
N1—Cu2—N3—C3	-71.0 (2)	N1—Cu2—N5—C6	74.5 (2)
N5 ⁱ —Cu2—N3—C4	138.2 (3)	N3—Cu2—N5—C5	37.4 (3)
N5—Cu2—N3—C4	-41.8 (3)	N3 ⁱ —Cu2—N5—C5	-142.6 (3)
N1 ⁱ —Cu2—N3—C4	-127.6 (2)	N1 ⁱ —Cu2—N5—C5	126.3 (3)
N1—Cu2—N3—C4	52.4 (2)	N1—Cu2—N5—C5	-53.7 (3)
N3—C4—N4—C7	75.0 (4)		

Symmetry codes: (i) $-x+2, -y+1, -z$; (ii) $-x+2, -y+1/2, z$; (iii) $-y+5/4, x-3/4, -z+1/4$; (iv) $y+3/4, -x+5/4, -z+1/4$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3C \cdots N2 ⁱ	0.93	2.44	3.261 (4)	147
N5—H5C \cdots O2 ^v	0.93	2.38	3.291 (8)	165

O1—H1C···O4 ^{vi}	0.85	2.48	3.149 (7)	136
O4—H4D···N2 ^{vii}	0.85	2.11	2.729 (6)	129
O4—H4E···O2 ^{viii}	0.85	2.47	3.279 (9)	159

Symmetry codes: (i) $-x+2, -y+1, -z$; (v) $y+3/4, -x+3/4, z-1/4$; (vi) $-y+5/4, x+1/4, -z+5/4$; (vii) $x-1, y, z+1$; (viii) $y-1/4, -x+1/4, -z+5/4$.