

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 2-(4-chlorophenyl)-3-(2,4-difluorophenoxy)acrylate

Hai-Bin Gong,* Jie Wang, Ying Liu and Lei Wang

Xuzhou Central Hospital, Xuzhou Cardiovascular Disease Institute, Xuzhou 221009, People's Republic of China

Correspondence e-mail: adler_20008@yahoo.com.cn


Received 31 October 2008; accepted 11 November 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.065; wR factor = 0.189; data-to-parameter ratio = 13.4.

In the molecule of the title compound, $\text{C}_{17}\text{H}_{13}\text{ClF}_2\text{O}_3$, the dihedral angles formed by the aromatic rings of the chlorobenzene and difluorobenzene groups with the plane of the acrylate unit are 48.85 (12) and 9.07 (14)°, respectively. In the crystal structure, molecules are linked by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen-bond interactions, forming chains along the c axis.

Related literature

For the synthesis and crystal structures of related compounds, see: Li, Xue *et al.* (2008); Li, Wang & Jian (2008); Lin & Jian (2008); Liu *et al.* (2008). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{13}\text{ClF}_2\text{O}_3$
 $M_r = 338.72$
 Monoclinic, $P2_1/c$
 $a = 16.275$ (3) Å
 $b = 7.503$ (2) Å
 $c = 13.812$ (3) Å
 $\beta = 111.11$ (3)°

$V = 1573.4$ (7) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.28$ mm⁻¹
 $T = 298$ (2) K
 $0.30 \times 0.10 \times 0.10$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\min} = 0.922$, $T_{\max} = 0.973$

2956 measured reflections
 2823 independent reflections
 1566 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.189$
 $S = 1.02$
 2823 reflections

210 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{O1}^i$	0.93	2.51	3.321 (4)	146

 Symmetry code: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This work was supported by the National Natural Science Foundation of China (No. 30572073), the Natural Science Foundation of Jiangsu Province of China (No. DK2005428), the Medical Science and Technology Development Foundation, Jiangsu Province Department of Health (No. K200402), and the Social Development Foundation of Xuzhou (No. X2003025).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2266).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Li, Y.-F., Wang, L.-T. & Jian, F.-F. (2008). *Acta Cryst.* **E64**, o2140.
 Li, H.-Q., Xue, J.-Y., Shi, L., Gui, S.-Y. & Zhu, H.-L. (2008). *Eur. J. Med. Chem.* **43**, 662–667.
 Lin, J. & Jian, F.-F. (2008). *Acta Cryst.* **E64**, o2130.
 Liu, X.-H., Cui, P., Song, B.-A., Bhadury, P. S., Zhu, H.-L. & Wang, S.-F. (2008). *Bioorg. Med. Chem.* **16**, 4075–4082.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2008). E64, o2373 [doi:10.1107/S1600536808037318]

Ethyl 2-(4-chlorophenyl)-3-(2,4-difluorophenoxy)acrylate**Hai-Bin Gong, Jie Wang, Ying Liu and Lei Wang****S1. Comment**

Recently, the synthesis and structure of a number of etheric compounds have been widely investigated (Li, Xue *et al.*, 2008; Li, Wang & Jian, 2008; Lin & Jian, 2008; Liu *et al.*, 2008). We report herein the crystal structure of the new title compound.


In the molecule of the title compound (Fig. 1), the dihedral angles between the aromatic rings of the chlorobenzene and difluorobenzene groups with the plane of the acrylate unit are 48.85 (12) and 9.07 (14)° respectively. All the bond lengths (Allen *et al.*, 1987) and angles are not unusual. In the crystal structure, molecules are linked by weak intermolecular C—H···O hydrogen interactions forming chains along the *c* axis (Table 1).

S2. Experimental

An equimolar solution of ethyl 3-bromo-2-(4-chlorophenyl)acrylate and 2,4-difluorophenol in chloroform was left to react overnight at room temperature. Block-shaped crystals suitable for X-ray analysis were obtained by slow evaporation of the solvent in air for five days.

S3. Refinement

H atoms were included in the riding model approximation with C—H = 0.93–0.97 Å and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $1.5 U_{\text{eq}}(\text{C})$ for methyl H atoms.

**Figure 1**

The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme.

Ethyl 2-(4-chlorophenyl)-3-(2,4-difluorophenoxy)acrylate

Crystal data

$C_{17}H_{13}ClF_2O_3$
 $M_r = 338.72$
 Monoclinic, $P2_1/c$
 Hall symbol: $-P\ 2_1/c$
 $a = 16.275\ (3)\ \text{\AA}$
 $b = 7.503\ (2)\ \text{\AA}$
 $c = 13.812\ (3)\ \text{\AA}$
 $\beta = 111.11\ (3)^\circ$
 $V = 1573.4\ (7)\ \text{\AA}^3$
 $Z = 4$

$F(000) = 696$
 $D_x = 1.430\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 1002 reflections
 $\theta = 2.4\text{--}24.5^\circ$
 $\mu = 0.28\ \text{mm}^{-1}$
 $T = 298\ \text{K}$
 Block, colorless
 $0.30 \times 0.10 \times 0.10\ \text{mm}$

Data collection

Bruker SMART 1000 CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2001)
 $T_{\min} = 0.922$, $T_{\max} = 0.973$

2956 measured reflections
 2823 independent reflections
 1566 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$
 $\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 1.3^\circ$
 $h = -19 \rightarrow 18$
 $k = -9 \rightarrow 10$
 $l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.189$
 $S = 1.02$
 2823 reflections
 210 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0904P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXTL* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.018 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0187 (2)	0.7796 (5)	0.9699 (3)	0.0450 (9)
C2	1.1022 (2)	0.7724 (5)	1.0453 (3)	0.0473 (9)
C3	1.1764 (2)	0.8337 (6)	1.0319 (3)	0.0581 (11)
H3	1.2315	0.8266	1.0844	0.070*
C4	1.1654 (3)	0.9066 (6)	0.9368 (3)	0.0597 (11)
C5	1.0851 (3)	0.9178 (6)	0.8598 (3)	0.0621 (12)
H5	1.0797	0.9691	0.7964	0.074*
C6	1.0121 (3)	0.8535 (6)	0.8755 (3)	0.0561 (11)
H6	0.9574	0.8596	0.8222	0.067*
C7	0.7054 (2)	0.6469 (5)	0.8670 (3)	0.0423 (9)
C8	0.6325 (2)	0.7234 (5)	0.8808 (3)	0.0540 (10)
H8	0.6376	0.7663	0.9459	0.065*
C9	0.5527 (2)	0.7366 (6)	0.7996 (3)	0.0571 (11)
H9	0.5045	0.7876	0.8100	0.069*
C10	0.5451 (2)	0.6742 (5)	0.7036 (3)	0.0543 (11)
C11	0.6147 (3)	0.5952 (6)	0.6881 (3)	0.0572 (11)
H11	0.6086	0.5513	0.6229	0.069*
C12	0.6947 (2)	0.5804 (5)	0.7696 (3)	0.0482 (10)
H12	0.7418	0.5252	0.7588	0.058*
C13	0.8635 (2)	0.7022 (5)	0.9293 (3)	0.0465 (9)
H13	0.8526	0.7373	0.8612	0.056*
C14	0.7936 (2)	0.6482 (5)	0.9511 (3)	0.0426 (9)
C15	0.8065 (2)	0.5946 (5)	1.0579 (3)	0.0483 (10)

C16	0.7414 (3)	0.4744 (6)	1.1715 (3)	0.0627 (12)
H16A	0.7580	0.5754	1.2185	0.075*
H16B	0.7858	0.3824	1.1975	0.075*
C17	0.6537 (3)	0.4055 (7)	1.1643 (3)	0.0716 (13)
H17A	0.6093	0.4922	1.1303	0.107*
H17B	0.6538	0.3831	1.2327	0.107*
H17C	0.6415	0.2968	1.1250	0.107*
C11	0.44478 (7)	0.69754 (19)	0.60039 (9)	0.0891 (5)
F1	1.10978 (13)	0.7012 (3)	1.13855 (15)	0.0663 (7)
F2	1.23761 (16)	0.9699 (4)	0.9209 (2)	0.0858 (9)
O3	0.94846 (18)	0.7113 (4)	0.9954 (2)	0.0723 (9)
O1	0.87704 (17)	0.6033 (4)	1.13049 (18)	0.0666 (9)
O2	0.73412 (16)	0.5282 (4)	1.06775 (17)	0.0525 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.043 (2)	0.048 (2)	0.043 (2)	−0.0032 (18)	0.0138 (16)	−0.0068 (18)
C2	0.047 (2)	0.057 (2)	0.0348 (19)	0.0045 (19)	0.0106 (16)	−0.0028 (18)
C3	0.042 (2)	0.078 (3)	0.049 (2)	−0.007 (2)	0.0090 (17)	−0.010 (2)
C4	0.053 (2)	0.070 (3)	0.062 (3)	−0.008 (2)	0.028 (2)	−0.011 (2)
C5	0.071 (3)	0.074 (3)	0.044 (2)	−0.001 (2)	0.024 (2)	0.001 (2)
C6	0.049 (2)	0.073 (3)	0.040 (2)	−0.005 (2)	0.0084 (17)	−0.002 (2)
C7	0.042 (2)	0.043 (2)	0.039 (2)	−0.0041 (17)	0.0112 (15)	0.0051 (16)
C8	0.048 (2)	0.062 (3)	0.046 (2)	0.002 (2)	0.0106 (17)	−0.004 (2)
C9	0.044 (2)	0.063 (3)	0.060 (3)	0.007 (2)	0.0137 (19)	0.004 (2)
C10	0.044 (2)	0.056 (3)	0.048 (2)	−0.003 (2)	−0.0007 (17)	0.0099 (19)
C11	0.059 (2)	0.067 (3)	0.038 (2)	−0.006 (2)	0.0081 (18)	−0.001 (2)
C12	0.048 (2)	0.054 (2)	0.041 (2)	−0.0005 (19)	0.0140 (16)	−0.0013 (18)
C13	0.045 (2)	0.055 (2)	0.0329 (18)	0.0044 (19)	0.0056 (16)	0.0002 (17)
C14	0.042 (2)	0.047 (2)	0.0327 (18)	0.0010 (17)	0.0072 (15)	−0.0008 (16)
C15	0.049 (2)	0.052 (2)	0.039 (2)	−0.0023 (19)	0.0113 (18)	−0.0016 (18)
C16	0.069 (3)	0.076 (3)	0.038 (2)	−0.006 (2)	0.0140 (19)	0.002 (2)
C17	0.075 (3)	0.084 (3)	0.057 (3)	−0.013 (3)	0.025 (2)	−0.001 (2)
C11	0.0572 (7)	0.1132 (11)	0.0671 (8)	0.0017 (7)	−0.0137 (5)	0.0169 (7)
F1	0.0541 (13)	0.0976 (19)	0.0380 (12)	−0.0022 (13)	0.0054 (10)	0.0107 (12)
F2	0.0689 (16)	0.119 (2)	0.0845 (18)	−0.0260 (16)	0.0462 (14)	−0.0119 (17)
O3	0.0592 (18)	0.086 (2)	0.0644 (19)	−0.0014 (17)	0.0134 (15)	−0.0021 (17)
O1	0.0509 (16)	0.099 (2)	0.0377 (15)	−0.0153 (16)	0.0018 (12)	0.0063 (15)
O2	0.0487 (15)	0.0678 (18)	0.0382 (13)	−0.0050 (14)	0.0123 (11)	0.0020 (13)

Geometric parameters (Å, °)

C1—C6	1.384 (5)	C10—C11	1.363 (5)
C1—C2	1.385 (5)	C10—C11	1.747 (4)
C1—O3	1.409 (4)	C11—C12	1.385 (5)
C2—F1	1.358 (4)	C11—H11	0.9300
C2—C3	1.365 (5)	C12—H12	0.9300

C3—C4	1.373 (6)	C13—C14	1.340 (5)
C3—H3	0.9300	C13—O3	1.356 (4)
C4—F2	1.357 (4)	C13—H13	0.9300
C4—C5	1.359 (5)	C14—C15	1.469 (5)
C5—C6	1.370 (5)	C15—O1	1.224 (4)
C5—H5	0.9300	C15—O2	1.331 (4)
C6—H6	0.9300	C16—O2	1.452 (4)
C7—C12	1.386 (5)	C16—C17	1.486 (5)
C7—C8	1.392 (5)	C16—H16A	0.9700
C7—C14	1.486 (4)	C16—H16B	0.9700
C8—C9	1.380 (5)	C17—H17A	0.9600
C8—H8	0.9300	C17—H17B	0.9600
C9—C10	1.368 (5)	C17—H17C	0.9600
C9—H9	0.9300		
C6—C1—C2	116.5 (3)	C10—C11—C12	119.9 (4)
C6—C1—O3	125.9 (3)	C10—C11—H11	120.1
C2—C1—O3	117.6 (3)	C12—C11—H11	120.1
F1—C2—C3	118.6 (3)	C11—C12—C7	120.8 (4)
F1—C2—C1	117.2 (3)	C11—C12—H12	119.6
C3—C2—C1	124.1 (4)	C7—C12—H12	119.6
C2—C3—C4	116.6 (4)	C14—C13—O3	127.3 (3)
C2—C3—H3	121.7	C14—C13—H13	116.3
C4—C3—H3	121.7	O3—C13—H13	116.3
F2—C4—C5	119.7 (4)	C13—C14—C15	118.8 (3)
F2—C4—C3	118.2 (4)	C13—C14—C7	119.1 (3)
C5—C4—C3	122.1 (4)	C15—C14—C7	122.1 (3)
C4—C5—C6	119.8 (4)	O1—C15—O2	122.7 (3)
C4—C5—H5	120.1	O1—C15—C14	124.1 (4)
C6—C5—H5	120.1	O2—C15—C14	113.2 (3)
C5—C6—C1	120.9 (4)	O2—C16—C17	107.1 (3)
C5—C6—H6	119.6	O2—C16—H16A	110.3
C1—C6—H6	119.6	C17—C16—H16A	110.3
C12—C7—C8	117.8 (3)	O2—C16—H16B	110.3
C12—C7—C14	120.6 (3)	C17—C16—H16B	110.3
C8—C7—C14	121.4 (3)	H16A—C16—H16B	108.5
C9—C8—C7	121.1 (4)	C16—C17—H17A	109.5
C9—C8—H8	119.4	C16—C17—H17B	109.5
C7—C8—H8	119.4	H17A—C17—H17B	109.5
C10—C9—C8	119.6 (4)	C16—C17—H17C	109.5
C10—C9—H9	120.2	H17A—C17—H17C	109.5
C8—C9—H9	120.2	H17B—C17—H17C	109.5
C11—C10—C9	120.7 (3)	C13—O3—C1	124.9 (3)
C11—C10—C11	120.0 (3)	C15—O2—C16	116.4 (3)
C9—C10—C11	119.2 (3)		
C6—C1—C2—F1	179.6 (3)	C10—C11—C12—C7	-0.7 (6)
O3—C1—C2—F1	-0.6 (5)	C8—C7—C12—C11	2.1 (6)

C6—C1—C2—C3	0.4 (6)	C14—C7—C12—C11	-173.4 (4)
O3—C1—C2—C3	-179.8 (4)	O3—C13—C14—C15	1.2 (6)
F1—C2—C3—C4	-179.2 (3)	O3—C13—C14—C7	-179.6 (3)
C1—C2—C3—C4	0.0 (6)	C12—C7—C14—C13	44.5 (5)
C2—C3—C4—F2	179.6 (4)	C8—C7—C14—C13	-130.9 (4)
C2—C3—C4—C5	0.1 (7)	C12—C7—C14—C15	-136.4 (4)
F2—C4—C5—C6	179.8 (4)	C8—C7—C14—C15	48.3 (5)
C3—C4—C5—C6	-0.7 (7)	C13—C14—C15—O1	3.5 (6)
C4—C5—C6—C1	1.1 (7)	C7—C14—C15—O1	-175.7 (4)
C2—C1—C6—C5	-1.0 (6)	C13—C14—C15—O2	-174.1 (3)
O3—C1—C6—C5	179.3 (4)	C7—C14—C15—O2	6.8 (5)
C12—C7—C8—C9	-1.6 (6)	C14—C13—O3—C1	-175.8 (4)
C14—C7—C8—C9	173.9 (4)	C6—C1—O3—C13	2.4 (6)
C7—C8—C9—C10	-0.2 (6)	C2—C1—O3—C13	-177.4 (4)
C8—C9—C10—C11	1.6 (6)	O1—C15—O2—C16	3.3 (6)
C8—C9—C10—C11	-178.1 (3)	C14—C15—O2—C16	-179.1 (3)
C9—C10—C11—C12	-1.2 (6)	C17—C16—O2—C15	179.9 (3)
C11—C10—C11—C12	178.6 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6 \cdots O1 ⁱ	0.93	2.51	3.321 (4)	146

Symmetry code: (i) $x, -y+3/2, z-1/2$.