

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(4-Chlorophenyl)-2-[(3-phenyl-isoquinolin-1-yl)sulfanyl]ethanone

P. Manivel,^a Venkatesha R. Hathwar,^b P. Nithya,^a R. Subashini^a and F. Nawaz Khan^{a*}

^aChemistry Division, School of Science and Humanities, VIT University, Vellore 632 014, Tamil Nadu, India, and ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India
Correspondence e-mail: nawaz_f@yahoo.co.in


Received 27 December 2008; accepted 5 January 2009

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.071; wR factor = 0.125; data-to-parameter ratio = 13.7.

The title compound, $\text{C}_{23}\text{H}_{16}\text{ClNOS}$, exhibits dihedral angles of 11.73 (1) and 66.07 (1)°, respectively, between the mean plane of the isoquinoline system and the attached phenyl ring, and between the isoquinoline system and the chlorophenyl ring. The dihedral angle between the phenyl and chlorophenyl rings is 54.66 (1)°.

Related literature

For general background, see: Cremlyn *et al.* (1996); Carreno (1995); Kondo *et al.* (2000); Mosberg & Omnaas (1985); McReynolds *et al.* (2004). For related crystal structures, see: Hathwar *et al.* (2008); Manivel *et al.* (2009).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{16}\text{ClNOS}$
 $M_r = 389.89$
Orthorhombic, $Pbca$
 $a = 9.5874$ (14) Å
 $b = 17.888$ (3) Å
 $c = 22.025$ (3) Å
 $V = 3777.3$ (10) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.33$ mm⁻¹
 $T = 290$ (2) K
 $0.26 \times 0.20 \times 0.14$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.901$, $T_{\max} = 0.956$
25674 measured reflections
3332 independent reflections
1915 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.116$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.125$
 $S = 1.04$
3332 reflections
244 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.23$ e Å⁻³
 $\Delta\rho_{\min} = -0.23$ e Å⁻³

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *CAMERON* (Watkin *et al.*, 1993); software used to prepare material for publication: *SHELXL97*.

We thank the Department of Science and Technology, India, for use of the CCD facility set up under the IRHPA-DST program at IISc. We thank Prof T. N. Guru Row, IISc, Bangalore, for useful crystallographic discussions. FNK thanks the DST for Fast Track Proposal funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2532).

References

- Bruker (2004). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Carreno, M. C. (1995). *Chem. Rev.* **95**, 1717–1760.
Cremlyn, R. J. (1996). In *An Introduction to Organosulfur Chemistry*. New York: John Wiley.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Hathwar, V. R., Prabakaran, K., Subashini, R., Manivel, P. & Khan, F. N. (2008). *Acta Cryst.* **E64**, o2295.
Kondo, T. & Mitsudo, T. (2000). *Chem. Rev.* **100**, 3205–3220.
Manivel, P., Hathwar, V. R., Nithya, P., Prabakaran, K. & Khan, F. N. (2009). *Acta Cryst.* **E65**, o137–138.
McReynolds, M. D., Dougherty, J. M. & Hanson, P. R. (2004). *Chem. Rev.* **104**, 2239–2258.
Mosberg, H. I. & Omnaas, J. R. (1985). *J. Am. Chem. Soc.* **107**, 2986–2987.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Watkin, D. J., Pearce, L. & Prout, C. K. (1993). *CAMERON*. Chemical Crystallography Laboratory, University of Oxford, England.

supporting information

Acta Cryst. (2009). E65, o254 [doi:10.1107/S1600536809000257]

1-(4-Chlorophenyl)-2-[(3-phenylisoquinolin-1-yl)sulfanyl]ethanone

P. Manivel, Venkatesha R. Hathwar, P. Nithya, R. Subashini and F. Nawaz Khan

S1. Comment

Organic thioethers are useful synthetic intermediates, key reagents in organic synthesis, bio-organic, medicinal, and heterocyclic chemistry (Cremlyn, 1996 and McReynolds *et al.*, 2004). They are also useful as heteroatomic functional groups in organic synthesis, for example, chiral sulphoxides can be generated by the oxidation of thioethers which are useful as auxiliaries in asymmetric syntheses (Carreno, 1995). Many syntheses have been reported for the preparation of thioethers in literature (Kondo *et al.*, 2000) whereas commonly used method is the alkylation of thiols (Mosberg *et al.*, 1985 and references there in).

The compound (I) forms the dihedral angles of 11.73 (1)° and 66.07 (1)° between mean plane of isoquinoline moiety and phenyl ring and mean plane of isoquinoline moiety and chlorophenyl ring respectively. The crystal packing is stabilized by C—H···O inter molecular Hydrogen bonds (Figure 2).

S2. Experimental

3-Phenylisoquinoline-1-thiol and 2-bromo-1-(4-chlorophenyl)ethanone were mixed in the ratio 1:1.05 equivalents with ethanol in a round bottom flask. Then it was heated under nitrogen atmosphere on an oil bath at 323 K. After 2 h, the products were filtered and dissolved in chloroform. Further, it was washed with water, dried and concentrated. The single-crystal for X-ray structure analysis was obtained from ether solution by slow evaporation.

S3. Refinement

All the H atoms in (I) were positioned geometrically and refined using a riding model with C—H bond lengths of 0.93 Å and 0.97 Å for aromatic and for methylene H atoms respectively and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for all carbon bound H atoms.




Figure 1

ORTEP diagram of the asymmetric unit of (I) with 50% probability displacement ellipsoids.




Figure 2

The crystal packing diagram of (I). The dotted lines indicate intermolecular C—H...O hydrogen bonds. All H atoms have been omitted for clarity.

1-(4-Chlorophenyl)-2-[(3-phenylisoquinolin-1-yl)sulfanyl]ethanone

Crystal data

$C_{23}H_{16}ClNOS$

$M_r = 389.89$

Orthorhombic, *Pbca*

Hall symbol: $-P\ 2ac\ 2ab$

$a = 9.5874\ (14)\ \text{\AA}$

$b = 17.888\ (3)\ \text{\AA}$

$c = 22.025\ (3)\ \text{\AA}$

$V = 3777.3\ (10)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1616$

$D_x = 1.371\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 871 reflections

$\theta = 1.9\text{--}25.0^\circ$

$\mu = 0.33\ \text{mm}^{-1}$

$T = 290\ \text{K}$

Rod, colourless

$0.26 \times 0.20 \times 0.14\ \text{mm}$

Data collection

Bruker SMART CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.901$, $T_{\max} = 0.956$

25674 measured reflections
 3332 independent reflections
 1915 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.116$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -11 \rightarrow 11$
 $k = -21 \rightarrow 21$
 $l = -24 \rightarrow 26$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.125$
 $S = 1.04$
 3332 reflections
 244 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0308P)^2 + 0.7191P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	-0.16890 (17)	0.07371 (9)	0.25163 (6)	0.1375 (6)
S1	-0.09127 (10)	0.34062 (5)	-0.06818 (5)	0.0545 (3)
O1	0.0295 (2)	0.19469 (13)	-0.02148 (12)	0.0588 (7)
N1	0.0851 (3)	0.36685 (14)	0.02265 (13)	0.0406 (7)
C1	0.0416 (3)	0.39062 (18)	-0.03051 (16)	0.0401 (9)
C2	0.1941 (3)	0.40302 (18)	0.05067 (15)	0.0390 (9)
C3	0.2597 (4)	0.46147 (19)	0.02355 (16)	0.0460 (9)
H3	0.3348	0.4841	0.0430	0.055*
C4	0.2805 (4)	0.54850 (19)	-0.06405 (19)	0.0554 (10)
H4	0.3564	0.5724	-0.0463	0.067*
C5	0.2337 (5)	0.5720 (2)	-0.1189 (2)	0.0644 (12)
H5	0.2784	0.6111	-0.1387	0.077*
C6	0.1184 (5)	0.5375 (2)	-0.14581 (18)	0.0665 (12)
H6	0.0862	0.5543	-0.1832	0.080*
C7	0.0532 (4)	0.47981 (19)	-0.11778 (17)	0.0570 (11)
H7	-0.0237	0.4575	-0.1360	0.068*

C8	0.1007 (4)	0.45349 (17)	-0.06155 (16)	0.0410 (8)
C9	0.2160 (4)	0.48838 (17)	-0.03338 (16)	0.0428 (9)
C10	0.2362 (4)	0.3727 (2)	0.11049 (16)	0.0439 (9)
C11	0.3618 (4)	0.3927 (2)	0.13781 (17)	0.0656 (12)
H11	0.4201	0.4267	0.1184	0.079*
C12	0.4012 (5)	0.3632 (3)	0.19275 (19)	0.0801 (13)
H12	0.4854	0.3779	0.2100	0.096*
C13	0.3193 (5)	0.3129 (3)	0.22247 (19)	0.0778 (13)
H13	0.3463	0.2933	0.2598	0.093*
C14	0.1957 (5)	0.2919 (2)	0.19589 (19)	0.0748 (13)
H14	0.1390	0.2570	0.2152	0.090*
C15	0.1543 (4)	0.3218 (2)	0.14102 (17)	0.0594 (11)
H15	0.0694	0.3072	0.1243	0.071*
C16	-0.1538 (3)	0.28339 (18)	-0.00757 (16)	0.0486 (10)
H16A	-0.1685	0.3150	0.0276	0.058*
H16B	-0.2440	0.2634	-0.0193	0.058*
C17	-0.0618 (4)	0.21887 (18)	0.01117 (18)	0.0454 (10)
C18	-0.0910 (4)	0.18457 (17)	0.07171 (18)	0.0470 (9)
C19	-0.2116 (4)	0.1996 (2)	0.10416 (18)	0.0529 (10)
H19	-0.2767	0.2330	0.0885	0.064*
C20	-0.2360 (4)	0.1656 (3)	0.15937 (19)	0.0693 (12)
H20	-0.3179	0.1754	0.1806	0.083*
C21	-0.1389 (6)	0.1174 (3)	0.18275 (19)	0.0769 (13)
C22	-0.0177 (5)	0.1022 (2)	0.1514 (2)	0.0762 (13)
H22	0.0480	0.0695	0.1675	0.091*
C23	0.0055 (4)	0.1354 (2)	0.0962 (2)	0.0630 (11)
H23	0.0869	0.1249	0.0750	0.076*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.1541 (14)	0.1891 (16)	0.0694 (9)	-0.0056 (13)	-0.0200 (9)	0.0362 (10)
S1	0.0462 (6)	0.0542 (5)	0.0632 (7)	-0.0085 (5)	-0.0116 (6)	-0.0032 (5)
O1	0.0391 (15)	0.0504 (15)	0.087 (2)	0.0030 (12)	0.0112 (15)	-0.0136 (14)
N1	0.0357 (17)	0.0379 (15)	0.0481 (19)	-0.0030 (14)	0.0007 (16)	-0.0072 (14)
C1	0.032 (2)	0.038 (2)	0.050 (2)	0.0031 (16)	0.0014 (18)	-0.0074 (18)
C2	0.036 (2)	0.037 (2)	0.045 (2)	0.0020 (17)	0.0029 (17)	-0.0125 (17)
C3	0.044 (2)	0.040 (2)	0.054 (2)	-0.0063 (17)	-0.005 (2)	-0.0126 (19)
C4	0.055 (3)	0.040 (2)	0.071 (3)	-0.0047 (19)	0.010 (2)	-0.005 (2)
C5	0.074 (3)	0.043 (2)	0.076 (3)	-0.001 (2)	0.017 (3)	0.006 (2)
C6	0.084 (3)	0.058 (3)	0.058 (3)	0.005 (2)	0.003 (3)	0.011 (2)
C7	0.060 (3)	0.048 (2)	0.063 (3)	0.000 (2)	-0.004 (2)	0.003 (2)
C8	0.042 (2)	0.0353 (18)	0.046 (2)	0.0077 (17)	0.0068 (19)	-0.0053 (18)
C9	0.043 (2)	0.0303 (19)	0.055 (2)	-0.0010 (17)	0.007 (2)	-0.0096 (18)
C10	0.040 (2)	0.047 (2)	0.045 (2)	-0.0009 (18)	0.001 (2)	-0.0121 (19)
C11	0.057 (3)	0.087 (3)	0.054 (3)	-0.015 (2)	-0.002 (2)	0.004 (2)
C12	0.064 (3)	0.119 (4)	0.057 (3)	-0.018 (3)	-0.013 (3)	0.003 (3)
C13	0.082 (4)	0.101 (4)	0.050 (3)	-0.005 (3)	-0.010 (3)	0.006 (3)

C14	0.084 (3)	0.078 (3)	0.063 (3)	-0.016 (3)	-0.009 (3)	0.019 (3)
C15	0.060 (3)	0.059 (3)	0.058 (3)	-0.009 (2)	-0.011 (2)	0.005 (2)
C16	0.033 (2)	0.044 (2)	0.068 (3)	-0.0058 (17)	-0.0008 (19)	-0.0072 (19)
C17	0.031 (2)	0.0346 (19)	0.071 (3)	-0.0067 (17)	-0.005 (2)	-0.011 (2)
C18	0.039 (2)	0.0350 (18)	0.067 (3)	-0.0037 (18)	-0.005 (2)	-0.012 (2)
C19	0.047 (3)	0.053 (2)	0.058 (3)	-0.002 (2)	-0.008 (2)	-0.008 (2)
C20	0.059 (3)	0.091 (3)	0.058 (3)	-0.009 (3)	-0.006 (3)	-0.013 (3)
C21	0.084 (4)	0.090 (3)	0.057 (3)	-0.012 (3)	-0.019 (3)	-0.003 (3)
C22	0.077 (4)	0.062 (3)	0.090 (4)	0.004 (3)	-0.030 (3)	0.007 (3)
C23	0.050 (3)	0.050 (2)	0.089 (3)	0.002 (2)	-0.008 (2)	-0.001 (2)

Geometric parameters (Å, °)

C11—C21	1.731 (4)	C11—C12	1.373 (5)
S1—C1	1.764 (3)	C11—H11	0.9300
S1—C16	1.786 (3)	C12—C13	1.362 (5)
O1—C17	1.212 (4)	C12—H12	0.9300
N1—C1	1.314 (4)	C13—C14	1.374 (5)
N1—C2	1.375 (4)	C13—H13	0.9300
C1—C8	1.433 (4)	C14—C15	1.379 (5)
C2—C3	1.359 (4)	C14—H14	0.9300
C2—C10	1.481 (4)	C15—H15	0.9300
C3—C9	1.407 (4)	C16—C17	1.510 (4)
C3—H3	0.9300	C16—H16A	0.9700
C4—C5	1.355 (5)	C16—H16B	0.9700
C4—C9	1.412 (4)	C17—C18	1.494 (5)
C4—H4	0.9300	C18—C23	1.385 (5)
C5—C6	1.398 (5)	C18—C19	1.385 (5)
C5—H5	0.9300	C19—C20	1.380 (5)
C6—C7	1.355 (5)	C19—H19	0.9300
C6—H6	0.9300	C20—C21	1.369 (6)
C7—C8	1.401 (4)	C20—H20	0.9300
C7—H7	0.9300	C21—C22	1.380 (6)
C8—C9	1.413 (4)	C22—C23	1.371 (5)
C10—C15	1.378 (4)	C22—H22	0.9300
C10—C11	1.392 (5)	C23—H23	0.9300
C1—S1—C16	100.46 (17)	C11—C12—H12	119.4
C1—N1—C2	119.3 (3)	C12—C13—C14	118.2 (4)
N1—C1—C8	123.6 (3)	C12—C13—H13	120.9
N1—C1—S1	119.0 (3)	C14—C13—H13	120.9
C8—C1—S1	117.3 (3)	C13—C14—C15	121.0 (4)
C3—C2—N1	121.1 (3)	C13—C14—H14	119.5
C3—C2—C10	123.1 (3)	C15—C14—H14	119.5
N1—C2—C10	115.8 (3)	C10—C15—C14	121.3 (4)
C2—C3—C9	121.2 (3)	C10—C15—H15	119.4
C2—C3—H3	119.4	C14—C15—H15	119.4
C9—C3—H3	119.4	C17—C16—S1	116.5 (2)

C5—C4—C9	121.1 (4)	C17—C16—H16A	108.2
C5—C4—H4	119.4	S1—C16—H16A	108.2
C9—C4—H4	119.4	C17—C16—H16B	108.2
C4—C5—C6	120.3 (4)	S1—C16—H16B	108.2
C4—C5—H5	119.9	H16A—C16—H16B	107.3
C6—C5—H5	119.9	O1—C17—C18	121.2 (3)
C7—C6—C5	120.5 (4)	O1—C17—C16	122.2 (3)
C7—C6—H6	119.8	C18—C17—C16	116.6 (3)
C5—C6—H6	119.8	C23—C18—C19	118.6 (4)
C6—C7—C8	120.6 (4)	C23—C18—C17	118.9 (4)
C6—C7—H7	119.7	C19—C18—C17	122.5 (3)
C8—C7—H7	119.7	C20—C19—C18	120.7 (4)
C7—C8—C9	119.6 (3)	C20—C19—H19	119.6
C7—C8—C1	123.8 (3)	C18—C19—H19	119.6
C9—C8—C1	116.5 (3)	C21—C20—C19	119.6 (4)
C3—C9—C4	123.8 (3)	C21—C20—H20	120.2
C3—C9—C8	118.2 (3)	C19—C20—H20	120.2
C4—C9—C8	118.0 (3)	C20—C21—C22	120.6 (4)
C15—C10—C11	116.9 (4)	C20—C21—C11	120.1 (4)
C15—C10—C2	121.4 (3)	C22—C21—C11	119.3 (4)
C11—C10—C2	121.8 (3)	C23—C22—C21	119.7 (4)
C12—C11—C10	121.3 (4)	C23—C22—H22	120.2
C12—C11—H11	119.3	C21—C22—H22	120.2
C10—C11—H11	119.3	C22—C23—C18	120.8 (4)
C13—C12—C11	121.3 (4)	C22—C23—H23	119.6
C13—C12—H12	119.4	C18—C23—H23	119.6
C2—N1—C1—C8	-0.6 (4)	C3—C2—C10—C11	-12.3 (5)
C2—N1—C1—S1	176.3 (2)	N1—C2—C10—C11	165.9 (3)
C16—S1—C1—N1	14.4 (3)	C15—C10—C11—C12	-0.4 (6)
C16—S1—C1—C8	-168.5 (2)	C2—C10—C11—C12	-178.8 (4)
C1—N1—C2—C3	-1.8 (4)	C10—C11—C12—C13	0.4 (7)
C1—N1—C2—C10	179.9 (3)	C11—C12—C13—C14	0.3 (7)
N1—C2—C3—C9	1.8 (5)	C12—C13—C14—C15	-1.0 (7)
C10—C2—C3—C9	180.0 (3)	C11—C10—C15—C14	-0.3 (6)
C9—C4—C5—C6	1.0 (6)	C2—C10—C15—C14	178.1 (3)
C4—C5—C6—C7	-0.8 (6)	C13—C14—C15—C10	1.0 (6)
C5—C6—C7—C8	-0.3 (6)	C1—S1—C16—C17	-73.9 (3)
C6—C7—C8—C9	1.2 (5)	S1—C16—C17—O1	-18.5 (4)
C6—C7—C8—C1	-176.9 (3)	S1—C16—C17—C18	163.1 (2)
N1—C1—C8—C7	-178.9 (3)	O1—C17—C18—C23	13.8 (5)
S1—C1—C8—C7	4.2 (4)	C16—C17—C18—C23	-167.7 (3)
N1—C1—C8—C9	2.9 (5)	O1—C17—C18—C19	-165.7 (3)
S1—C1—C8—C9	-174.0 (2)	C16—C17—C18—C19	12.8 (4)
C2—C3—C9—C4	-179.5 (3)	C23—C18—C19—C20	-0.8 (5)
C2—C3—C9—C8	0.7 (5)	C17—C18—C19—C20	178.7 (3)
C5—C4—C9—C3	-180.0 (3)	C18—C19—C20—C21	1.0 (6)
C5—C4—C9—C8	-0.2 (5)	C19—C20—C21—C22	-0.5 (6)

C7—C8—C9—C3	178.9 (3)	C19—C20—C21—C11	-179.4 (3)
C1—C8—C9—C3	-2.8 (4)	C20—C21—C22—C23	-0.2 (6)
C7—C8—C9—C4	-0.9 (4)	C11—C21—C22—C23	178.7 (3)
C1—C8—C9—C4	177.3 (3)	C21—C22—C23—C18	0.4 (6)
C3—C2—C10—C15	169.4 (3)	C19—C18—C23—C22	0.1 (5)
N1—C2—C10—C15	-12.4 (4)	C17—C18—C23—C22	-179.4 (3)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C4—H4...O1 ⁱ	0.93	2.51	3.322 (4)	147
C16—H16B...O1 ⁱⁱ	0.97	2.47	3.128 (4)	125

Symmetry codes: (i) $-x+1/2, y+1/2, z$; (ii) $x-1/2, -y+1/2, -z$.