

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 2-[2-((Z)-{1-*trans*-[2-(4-fluoro-3-methylphenyl)-2-methylcyclopropyl]-ethylidene}aminoxymethyl)phenyl]-2-[(*E*)-methoxyimino]acetate

Chunfeng Song,^a Ronald Ross,^b Steven Shaber^b and Bin Li^{c*}

^aMedical College of Chifeng University, Chifeng 024000, People's Republic of China, ^bDow AgroSciences, 9330 Zionsville Road, Indianapolis, Indiana 46268, USA, and ^cAgrochemicals Division, Shenyang Research Institute of Chemical Industry, Shenyang 110021, People's Republic of China

Correspondence e-mail: libin1@sinochem.com


Received 3 March 2009; accepted 12 March 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.046; wR factor = 0.140; data-to-parameter ratio = 14.1.

The title compound, $\text{C}_{24}\text{H}_{27}\text{FN}_2\text{O}_4$, is an important intermediate in the synthesis of fungicidal strobilurin-type compounds. In the crystal structure, the oxime bond attached to the cyclopropane ring adopts a *Z* configuration, while the oxime bond attached to the benzene ring adopts an *E* configuration. The fluoromethylphenyl group adopts a *trans* configuration with respect to the remainder of the molecule, and its mean plane forms a dihedral angle of $56.1(1)^\circ$ with the plane of the cyclopropane ring.

Related literature

For synthesis details and related literature, see: Li *et al.* (2008); Ross *et al.* (2001).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{27}\text{FN}_2\text{O}_4$
 $M_r = 426.48$
 Triclinic, $P\bar{1}$
 $a = 7.819(3)$ Å
 $b = 10.901(4)$ Å
 $c = 14.565(5)$ Å
 $\alpha = 105.134(4)^\circ$
 $\beta = 92.014(4)^\circ$
 $\gamma = 104.718(4)^\circ$
 $V = 1152.3(7)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.34 \times 0.30 \times 0.24$ mm

Data collection

Bruker APEXII CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 2003)
 $T_{\min} = 0.754$, $T_{\max} = 0.979$
 6297 measured reflections
 4026 independent reflections
 2977 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.140$
 $S = 1.05$
 4026 reflections
 285 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: B12355).

References

- Bruker (2003). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Li, B., Xiang, D. & Si, N. G. (2008). Int. Patent No. WO2008/104101A1.
 Ross, R., Nguyen, D. & Szapacs, E. (2001). Int. Patent No. WO01/87826A1.
 Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o796 [doi:10.1107/S1600536809009064]

Methyl 2-[2-((*Z*)-{1-*trans*-[2-(4-fluoro-3-methylphenyl)-2-methylcyclopropyl]-ethylidene}aminooxymethyl)phenyl]-2-[(*E*)-methoxyimino]acetate

Chunfeng Song, Ronald Ross, Steven Shaber and Bin Li

S1. Comment

Some cyclopropyl-containing Strobilurin-type compounds have been reported as a new class of fungicides characterized by their broad spectrum and high levels of fungicidal activity (Ross *et al.*, 2001; Li *et al.*, 2008). The title compound is an intermediate for preparing such compounds and the crystal structure is important to identify the configuration of possible products.

S2. Experimental

The title compound was prepared according to a published procedure (Li *et al.*, 2008) as follows: *trans* 2-(4-fluoro-3-methylphenyl)-2-methyl-1-acetylcyclopropane (26.8 g, 13 mmol) and methyl *E*-2-(aminooxymethyl)phenyl glyoxylate *O*-methyloxime (34.0 g, 14.3 mmol) were added into methanol (500 ml) and acetic acid (0.7 ml). The mixture was stirred for 14 h and washed with water and brine, dried by anhydrous magnesium sulfate, then evaporated to give an oily product. This product was separated with silica gel chromatography to give major product A (90%) and minor product B (10%). The product B was dissolved in acetone and left to stand at room temperature. Block crystals suitable for X-ray analyses were obtained.

¹H NMR(CDCI₃): δ 7.78–7.81 (m, 1H), 7.523–7.492 (m, 1H), 7.412–7.382 (m, 2H), 7.212–7.195 (m, 1H), 7.177–7.035 (m, 2H), 6.785–6.753 (t, 1H), 5.044–5.038 (d, 2H), 3.948 (s, 3H), 3.705 (s, 3H), 2.055–1.986 (m, 4H), 1.868 (s, 3H), 1.205–1.189 (m, 4H).

S3. Refinement

Although all H atoms were visible in difference Fourier maps, they were finally placed in geometrically calculated positions, with C—H distances in the range 0.93–0.98 Å, and included in the final refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$.




Figure 1

Molecular structure of the title compound with 30% probability displacement ellipsoids for non-H atoms.

Methyl 2-[2-((Z)-{1-*trans*-[2-(4-fluoro-3-methylphenyl)-2-methylcyclopropyl]ethylidene}aminooxymethyl)phenyl]-2-[(E)-methoxyimino]acetate

Crystal data

$C_{24}H_{27}FN_2O_4$

$M_r = 426.48$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.819 (3) \text{ \AA}$

$b = 10.901 (4) \text{ \AA}$

$c = 14.565 (5) \text{ \AA}$

$\alpha = 105.134 (4)^\circ$

$\beta = 92.014 (4)^\circ$

$\gamma = 104.718 (4)^\circ$

$V = 1152.3 (7) \text{ \AA}^3$

$Z = 2$

$F(000) = 452$

$D_x = 1.229 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2258 reflections

$\theta = 2.7\text{--}25.2^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, colorless

$0.34 \times 0.30 \times 0.24 \text{ mm}$

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2003)

$T_{\min} = 0.754$, $T_{\max} = 0.979$

6297 measured reflections

4026 independent reflections

2977 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.0^\circ$

$h = -9 \rightarrow 9$

$k = -9 \rightarrow 12$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.140$
 $S = 1.05$
 4026 reflections
 285 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0682P)^2 + 0.2537P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.15 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	1.35756 (18)	0.82907 (15)	0.97952 (11)	0.0942 (5)
O1	0.56929 (18)	0.50914 (14)	0.67273 (10)	0.0625 (4)
O2	0.6063 (2)	0.39143 (18)	0.88366 (11)	0.0903 (6)
O3	0.33762 (19)	0.25037 (15)	0.86384 (10)	0.0680 (4)
O4	0.41061 (16)	0.08197 (13)	0.59611 (9)	0.0600 (4)
N1	0.4299 (2)	0.55373 (18)	0.63920 (12)	0.0604 (4)
N2	0.37516 (19)	0.14127 (15)	0.68779 (10)	0.0509 (4)
C1	1.0499 (3)	0.9394 (2)	0.8529 (2)	0.0836 (7)
H1	1.0399	1.0172	0.8411	0.100*
C2	1.1977 (3)	0.9404 (2)	0.9074 (2)	0.0879 (8)
H2	1.2875	1.0183	0.9325	0.105*
C3	1.2108 (3)	0.8262 (2)	0.92420 (16)	0.0680 (6)
C4	1.0856 (3)	0.7074 (2)	0.88688 (14)	0.0588 (5)
C5	1.1107 (4)	0.5818 (3)	0.9027 (2)	0.0895 (8)
H5A	1.2018	0.5563	0.8658	0.134*
H5B	1.0012	0.5131	0.8829	0.134*
H5C	1.1452	0.5958	0.9693	0.134*
C6	0.9380 (3)	0.7098 (2)	0.83200 (14)	0.0558 (5)
H6	0.8506	0.6309	0.8053	0.067*
C7	0.9156 (3)	0.8237 (2)	0.81544 (15)	0.0590 (5)
C8	0.7563 (3)	0.8232 (2)	0.75450 (16)	0.0637 (5)
C9	0.7882 (4)	0.8208 (4)	0.6527 (2)	0.1044 (10)
H9A	0.8837	0.8960	0.6523	0.157*
H9B	0.6821	0.8235	0.6189	0.157*
H9C	0.8192	0.7413	0.6219	0.157*

C10	0.6289 (3)	0.8960 (3)	0.8011 (2)	0.0918 (8)
H10A	0.6542	0.9400	0.8691	0.110*
H10B	0.5744	0.9411	0.7642	0.110*
C11	0.5746 (3)	0.7483 (2)	0.77037 (15)	0.0624 (5)
H11	0.5744	0.7077	0.8230	0.075*
C12	0.4369 (3)	0.6710 (2)	0.68858 (15)	0.0609 (5)
C13	0.2966 (3)	0.7290 (3)	0.65974 (19)	0.0849 (7)
H13A	0.3490	0.7967	0.6302	0.127*
H13B	0.2447	0.7664	0.7153	0.127*
H13C	0.2061	0.6610	0.6152	0.127*
C14	0.5777 (3)	0.3937 (2)	0.60237 (15)	0.0659 (6)
H14A	0.6049	0.4141	0.5427	0.079*
H14B	0.4643	0.3271	0.5910	0.079*
C15	0.7210 (3)	0.34381 (19)	0.63878 (14)	0.0552 (5)
C16	0.8912 (3)	0.3751 (2)	0.61019 (18)	0.0790 (7)
H16	0.9156	0.4278	0.5688	0.095*
C17	1.0237 (3)	0.3291 (3)	0.6423 (2)	0.0924 (9)
H17	1.1361	0.3505	0.6219	0.111*
C18	0.9913 (3)	0.2521 (3)	0.7040 (2)	0.0822 (8)
H18	1.0807	0.2205	0.7251	0.099*
C19	0.8259 (3)	0.2222 (2)	0.73409 (16)	0.0629 (5)
H19	0.8041	0.1715	0.7770	0.075*
C20	0.6902 (2)	0.26630 (17)	0.70163 (13)	0.0482 (4)
C21	0.5128 (2)	0.22899 (18)	0.73554 (12)	0.0460 (4)
C22	0.4921 (3)	0.3000 (2)	0.83565 (13)	0.0536 (5)
C23	0.3080 (4)	0.3151 (3)	0.95956 (16)	0.0839 (7)
H23A	0.3990	0.3128	1.0046	0.126*
H23B	0.1937	0.2702	0.9735	0.126*
H23C	0.3116	0.4052	0.9642	0.126*
C24	0.2488 (3)	-0.0023 (2)	0.54163 (15)	0.0660 (6)
H24A	0.1929	-0.0644	0.5749	0.099*
H24B	0.2740	-0.0491	0.4803	0.099*
H24C	0.1707	0.0499	0.5329	0.099*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0727 (9)	0.0917 (10)	0.1010 (10)	0.0112 (7)	-0.0317 (8)	0.0138 (8)
O1	0.0619 (8)	0.0581 (8)	0.0631 (8)	0.0179 (7)	-0.0104 (7)	0.0099 (7)
O2	0.0780 (11)	0.0935 (12)	0.0624 (9)	-0.0017 (10)	0.0000 (8)	-0.0159 (9)
O3	0.0682 (9)	0.0725 (10)	0.0519 (8)	0.0143 (8)	0.0142 (7)	0.0015 (7)
O4	0.0439 (7)	0.0660 (9)	0.0507 (7)	0.0069 (6)	0.0036 (6)	-0.0089 (6)
N1	0.0528 (10)	0.0679 (12)	0.0589 (10)	0.0149 (8)	-0.0040 (8)	0.0179 (9)
N2	0.0423 (8)	0.0574 (9)	0.0468 (8)	0.0151 (7)	0.0040 (7)	0.0025 (7)
C1	0.0752 (16)	0.0552 (13)	0.111 (2)	0.0105 (12)	-0.0123 (14)	0.0179 (13)
C2	0.0710 (16)	0.0578 (15)	0.113 (2)	0.0013 (12)	-0.0212 (14)	0.0066 (14)
C3	0.0567 (12)	0.0704 (15)	0.0662 (13)	0.0114 (11)	-0.0096 (10)	0.0086 (11)
C4	0.0593 (12)	0.0604 (12)	0.0536 (11)	0.0124 (10)	0.0020 (9)	0.0150 (9)

C5	0.0882 (18)	0.0775 (17)	0.1009 (19)	0.0139 (14)	-0.0201 (15)	0.0343 (15)
C6	0.0532 (11)	0.0544 (12)	0.0532 (11)	0.0056 (9)	0.0018 (9)	0.0133 (9)
C7	0.0538 (11)	0.0591 (12)	0.0617 (12)	0.0129 (9)	0.0047 (9)	0.0152 (10)
C8	0.0556 (12)	0.0641 (13)	0.0752 (14)	0.0126 (10)	0.0068 (10)	0.0293 (11)
C9	0.0740 (17)	0.153 (3)	0.0969 (19)	0.0095 (17)	0.0048 (14)	0.074 (2)
C10	0.0781 (16)	0.0698 (16)	0.118 (2)	0.0295 (13)	-0.0085 (15)	0.0025 (15)
C11	0.0602 (12)	0.0647 (13)	0.0608 (12)	0.0187 (10)	0.0047 (10)	0.0138 (10)
C12	0.0527 (11)	0.0747 (15)	0.0586 (12)	0.0183 (10)	0.0055 (9)	0.0231 (11)
C13	0.0769 (16)	0.107 (2)	0.0785 (16)	0.0456 (15)	0.0001 (13)	0.0204 (14)
C14	0.0794 (15)	0.0581 (13)	0.0561 (12)	0.0167 (11)	-0.0040 (10)	0.0123 (10)
C15	0.0523 (11)	0.0469 (11)	0.0556 (11)	0.0063 (8)	0.0017 (9)	0.0035 (9)
C16	0.0671 (15)	0.0739 (15)	0.0773 (15)	-0.0045 (12)	0.0139 (12)	0.0112 (12)
C17	0.0399 (12)	0.096 (2)	0.111 (2)	0.0015 (12)	0.0135 (13)	-0.0070 (17)
C18	0.0427 (12)	0.0842 (17)	0.105 (2)	0.0210 (12)	-0.0052 (12)	-0.0011 (15)
C19	0.0465 (11)	0.0586 (12)	0.0778 (14)	0.0172 (9)	-0.0082 (10)	0.0082 (10)
C20	0.0395 (9)	0.0443 (10)	0.0517 (10)	0.0107 (8)	-0.0021 (8)	-0.0005 (8)
C21	0.0400 (9)	0.0493 (10)	0.0470 (10)	0.0164 (8)	-0.0018 (8)	0.0071 (8)
C22	0.0516 (11)	0.0551 (11)	0.0494 (10)	0.0159 (9)	-0.0017 (9)	0.0061 (9)
C23	0.1056 (19)	0.0876 (18)	0.0553 (13)	0.0333 (15)	0.0254 (13)	0.0059 (12)
C24	0.0537 (12)	0.0676 (13)	0.0585 (12)	0.0052 (10)	-0.0089 (9)	-0.0006 (10)

Geometric parameters (Å, °)

F1—C3	1.368 (2)	C10—C11	1.496 (3)
O1—C14	1.418 (2)	C10—H10A	0.970
O1—N1	1.421 (2)	C10—H10B	0.970
O2—C22	1.192 (2)	C11—C12	1.487 (3)
O3—C22	1.319 (2)	C11—H11	0.980
O3—C23	1.445 (3)	C12—C13	1.496 (3)
O4—N2	1.3944 (19)	C13—H13A	0.960
O4—C24	1.425 (2)	C13—H13B	0.960
N1—C12	1.279 (3)	C13—H13C	0.960
N2—C21	1.278 (2)	C14—C15	1.501 (3)
C1—C2	1.375 (3)	C14—H14A	0.970
C1—C7	1.385 (3)	C14—H14B	0.970
C1—H1	0.930	C15—C20	1.388 (3)
C2—C3	1.358 (3)	C15—C16	1.394 (3)
C2—H2	0.930	C16—C17	1.376 (4)
C3—C4	1.371 (3)	C16—H16	0.930
C4—C6	1.390 (3)	C17—C18	1.371 (4)
C4—C5	1.505 (3)	C17—H17	0.930
C5—H5A	0.960	C18—C19	1.368 (3)
C5—H5B	0.960	C18—H18	0.930
C5—H5C	0.960	C19—C20	1.387 (3)
C6—C7	1.377 (3)	C19—H19	0.930
C6—H6	0.930	C20—C21	1.483 (3)
C7—C8	1.502 (3)	C21—C22	1.497 (3)
C8—C10	1.500 (3)	C23—H23A	0.960

C8—C9	1.507 (3)	C23—H23B	0.960
C8—C11	1.508 (3)	C23—H23C	0.960
C9—H9A	0.960	C24—H24A	0.960
C9—H9B	0.960	C24—H24B	0.960
C9—H9C	0.960	C24—H24C	0.960
C14—O1—N1	107.55 (14)	C8—C11—H11	114.3
C22—O3—C23	116.14 (17)	N1—C12—C11	123.38 (19)
N2—O4—C24	109.21 (14)	N1—C12—C13	116.2 (2)
C12—N1—O1	111.23 (16)	C11—C12—C13	120.5 (2)
C21—N2—O4	111.21 (14)	C12—C13—H13A	109.5
C2—C1—C7	120.8 (2)	C12—C13—H13B	109.5
C2—C1—H1	119.6	H13A—C13—H13B	109.5
C7—C1—H1	119.6	C12—C13—H13C	109.5
C3—C2—C1	119.1 (2)	H13A—C13—H13C	109.5
C3—C2—H2	120.4	H13B—C13—H13C	109.5
C1—C2—H2	120.4	O1—C14—C15	107.78 (16)
C2—C3—F1	118.4 (2)	O1—C14—H14A	110.2
C2—C3—C4	123.2 (2)	C15—C14—H14A	110.2
F1—C3—C4	118.4 (2)	O1—C14—H14B	110.2
C3—C4—C6	116.2 (2)	C15—C14—H14B	110.2
C3—C4—C5	121.4 (2)	H14A—C14—H14B	108.5
C6—C4—C5	122.4 (2)	C20—C15—C16	117.8 (2)
C4—C5—H5A	109.5	C20—C15—C14	121.82 (18)
C4—C5—H5B	109.5	C16—C15—C14	120.4 (2)
H5A—C5—H5B	109.5	C17—C16—C15	121.1 (3)
C4—C5—H5C	109.5	C17—C16—H16	119.5
H5A—C5—H5C	109.5	C15—C16—H16	119.5
H5B—C5—H5C	109.5	C18—C17—C16	120.5 (2)
C7—C6—C4	122.83 (19)	C18—C17—H17	119.7
C7—C6—H6	118.6	C16—C17—H17	119.7
C4—C6—H6	118.6	C19—C18—C17	119.2 (2)
C6—C7—C1	117.8 (2)	C19—C18—H18	120.4
C6—C7—C8	121.80 (19)	C17—C18—H18	120.4
C1—C7—C8	120.3 (2)	C18—C19—C20	121.0 (2)
C10—C8—C7	118.3 (2)	C18—C19—H19	119.5
C10—C8—C9	118.5 (2)	C20—C19—H19	119.5
C7—C8—C9	114.35 (19)	C19—C20—C15	120.29 (18)
C10—C8—C11	59.65 (15)	C19—C20—C21	118.64 (18)
C7—C8—C11	118.65 (18)	C15—C20—C21	121.07 (16)
C9—C8—C11	116.8 (2)	N2—C21—C20	126.38 (16)
C8—C9—H9A	109.5	N2—C21—C22	116.44 (16)
C8—C9—H9B	109.5	C20—C21—C22	117.18 (15)
H9A—C9—H9B	109.5	O2—C22—O3	124.65 (18)
C8—C9—H9C	109.5	O2—C22—C21	122.16 (19)
H9A—C9—H9C	109.5	O3—C22—C21	113.20 (16)
H9B—C9—H9C	109.5	O3—C23—H23A	109.5
C11—C10—C8	60.42 (15)	O3—C23—H23B	109.5

C11—C10—H10A	117.7	H23A—C23—H23B	109.5
C8—C10—H10A	117.7	O3—C23—H23C	109.5
C11—C10—H10B	117.7	H23A—C23—H23C	109.5
C8—C10—H10B	117.7	H23B—C23—H23C	109.5
H10A—C10—H10B	114.9	O4—C24—H24A	109.5
C12—C11—C10	122.1 (2)	O4—C24—H24B	109.5
C12—C11—C8	121.18 (19)	H24A—C24—H24B	109.5
C10—C11—C8	59.93 (16)	O4—C24—H24C	109.5
C12—C11—H11	114.3	H24A—C24—H24C	109.5
C10—C11—H11	114.3	H24B—C24—H24C	109.5
C14—O1—N1—C12	-166.73 (18)	C10—C11—C12—N1	154.4 (2)
C24—O4—N2—C21	-172.57 (17)	C8—C11—C12—N1	82.6 (3)
C7—C1—C2—C3	0.0 (4)	C10—C11—C12—C13	-25.6 (3)
C1—C2—C3—F1	-179.2 (2)	C8—C11—C12—C13	-97.4 (3)
C1—C2—C3—C4	2.0 (4)	N1—O1—C14—C15	-178.28 (16)
C2—C3—C4—C6	-1.9 (4)	O1—C14—C15—C20	82.5 (2)
F1—C3—C4—C6	179.34 (18)	O1—C14—C15—C16	-97.2 (2)
C2—C3—C4—C5	176.4 (3)	C20—C15—C16—C17	0.8 (3)
F1—C3—C4—C5	-2.4 (3)	C14—C15—C16—C17	-179.4 (2)
C3—C4—C6—C7	-0.3 (3)	C15—C16—C17—C18	-0.5 (4)
C5—C4—C6—C7	-178.6 (2)	C16—C17—C18—C19	-0.6 (4)
C4—C6—C7—C1	2.2 (3)	C17—C18—C19—C20	1.3 (3)
C4—C6—C7—C8	178.95 (19)	C18—C19—C20—C15	-1.0 (3)
C2—C1—C7—C6	-2.1 (4)	C18—C19—C20—C21	179.03 (19)
C2—C1—C7—C8	-178.9 (2)	C16—C15—C20—C19	-0.1 (3)
C6—C7—C8—C10	115.1 (2)	C14—C15—C20—C19	-179.79 (17)
C1—C7—C8—C10	-68.2 (3)	C16—C15—C20—C21	179.88 (17)
C6—C7—C8—C9	-98.0 (3)	C14—C15—C20—C21	0.1 (3)
C1—C7—C8—C9	78.7 (3)	O4—N2—C21—C20	-0.3 (3)
C6—C7—C8—C11	46.2 (3)	O4—N2—C21—C22	-179.87 (15)
C1—C7—C8—C11	-137.1 (2)	C19—C20—C21—N2	-104.2 (2)
C7—C8—C10—C11	-108.4 (2)	C15—C20—C21—N2	75.9 (3)
C9—C8—C10—C11	106.1 (2)	C19—C20—C21—C22	75.4 (2)
C8—C10—C11—C12	-110.1 (2)	C15—C20—C21—C22	-104.5 (2)
C10—C8—C11—C12	111.6 (3)	C23—O3—C22—O2	1.0 (3)
C7—C8—C11—C12	-140.6 (2)	C23—O3—C22—C21	-179.41 (17)
C9—C8—C11—C12	2.7 (3)	N2—C21—C22—O2	-173.2 (2)
C7—C8—C11—C10	107.8 (2)	C20—C21—C22—O2	7.2 (3)
C9—C8—C11—C10	-108.8 (3)	N2—C21—C22—O3	7.2 (2)
O1—N1—C12—C11	-0.1 (3)	C20—C21—C22—O3	-172.41 (16)
O1—N1—C12—C13	179.85 (18)		
