

## N-(Pyrimidin-2-yl)aniline

**Edura Badaruddin, Nasir Shah Bakhtiar, Zaharah Aiyub,  
 Zanariah Abdullah and Seik Weng Ng\***

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
 Correspondence e-mail: seikweng@um.edu.my


Received 2 March 2009; accepted 3 March 2009

Key indicators: single-crystal X-ray study;  $T = 123\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  
 $R$  factor = 0.039;  $wR$  factor = 0.108; data-to-parameter ratio = 16.3.

There are two molecules in the asymmetric unit of the title compound,  $\text{C}_{10}\text{H}_9\text{N}_3$ , with inter-ring dihedral angles of  $31.1(1)$  and  $35.3(1)^\circ$ . The bridging  $\text{C}-\text{N}-\text{C}$  bond angles are  $128.2(1)$  and  $129.1(1)^\circ$ . In the crystal, the two independent molecules are linked into a dimer by two  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonds.

### Related literature

For the structure of 4-chloro-*N*-(pyrimidin-2-yl)aniline, see: Maizathul Akmam *et al.* (2009).



### Experimental

#### Crystal data

$\text{C}_{10}\text{H}_9\text{N}_3$	$b = 9.9382(2)\text{ \AA}$
$M_r = 171.20$	$c = 10.2038(2)\text{ \AA}$
Triclinic, $P\bar{1}$	$\alpha = 93.186(1)^\circ$
$a = 8.8792(2)\text{ \AA}$	$\beta = 103.665(1)^\circ$

$\gamma = 97.780(1)^\circ$   
 $V = 863.28(3)\text{ \AA}^3$   
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.08\text{ mm}^{-1}$   
 $T = 123\text{ K}$   
 $0.35 \times 0.20 \times 0.10\text{ mm}$

#### Data collection

Bruker SMART APEX  
 diffractometer  
 Absorption correction: none  
 8238 measured reflections

3950 independent reflections  
 3144 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.108$   
 $S = 1.03$   
 3950 reflections  
 243 parameters  
 2 restraints

H atoms treated by a mixture of  
 independent and constrained  
 refinement  
 $\Delta\rho_{\text{max}} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.23\text{ e \AA}^{-3}$

**Table 1**  
 Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1 $\cdots$ N5	0.89 (1)	2.10 (1)	2.972 (1)	164 (1)
N4—H4 $\cdots$ N2	0.89 (1)	2.15 (1)	3.020 (1)	165 (1)

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

The authors thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2492).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Maizathul Akmam, A. B., Abdullah, Z. & Ng, S. W. (2009). *Acta Cryst. E65*, o94.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

# supporting information

*Acta Cryst.* (2009). E65, o703 [doi:10.1107/S1600536809007685]

## N-(Pyrimidin-2-yl)aniline


**Edura Badaruddin, Nasir Shah Bakhtiar, Zaharah Aiyub, Zanariah Abdullah and Seik Weng Ng**

### S1. Experimental

2-Chloropyrimidine (0.05 mol), aniline (0.05 mol) and ethanol (5 ml) were heated at 423–433 K for 3 h. The product was dissolved in water and the solution extracted with ether. The ether phase was dried over sodium sulfate; the evaporation of the solvent gave well shaped colorless crystals along with some unidentified brown material.

### S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 Å) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H})$  set to  $1.2U_{\text{eq}}(\text{C})$ . The amino H-atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H  $0.88 \pm 0.01$  Å; their isotropic temperature factors were refined.



**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of the two independent molecules of  $\text{C}_{10}\text{H}_9\text{N}_3$  at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius. Dashed lines denote hydrogen bonds.

**N-(Pyrimidin-2-yl)aniline***Crystal data*

$C_{10}H_9N_3$   
 $M_r = 171.20$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 8.8792 (2)$  Å  
 $b = 9.9382 (2)$  Å  
 $c = 10.2038 (2)$  Å  
 $\alpha = 93.186 (1)^\circ$   
 $\beta = 103.665 (1)^\circ$   
 $\gamma = 97.780 (1)^\circ$   
 $V = 863.28 (3)$  Å<sup>3</sup>

$Z = 4$   
 $F(000) = 360$   
 $D_x = 1.317 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 3015 reflections  
 $\theta = 2.7\text{--}28.3^\circ$   
 $\mu = 0.08 \text{ mm}^{-1}$   
 $T = 123$  K  
Prism, colorless  
 $0.35 \times 0.20 \times 0.10$  mm

*Data collection*

Bruker SMART APEX  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
8238 measured reflections  
3950 independent reflections

3144 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.020$   
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.1^\circ$   
 $h = -11 \rightarrow 11$   
 $k = -12 \rightarrow 12$   
 $l = -13 \rightarrow 12$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.108$   
 $S = 1.03$   
3950 reflections  
243 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.056P)^2 + 0.1316P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.20 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.23 \text{ e } \text{\AA}^{-3}$

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.63241 (11)	0.69760 (10)	0.39973 (10)	0.0229 (2)
H1	0.5458 (13)	0.7328 (15)	0.4021 (15)	0.042 (4)*
N2	0.60532 (11)	0.62876 (10)	0.60461 (10)	0.0235 (2)
N3	0.82465 (11)	0.58169 (10)	0.51919 (10)	0.0257 (2)
N4	0.33475 (12)	0.78540 (10)	0.59499 (10)	0.0256 (2)
H4	0.4034 (15)	0.7306 (13)	0.5833 (14)	0.038 (4)*
N5	0.37902 (11)	0.86480 (10)	0.39948 (10)	0.0256 (2)
N6	0.18856 (11)	0.94997 (10)	0.50126 (10)	0.0245 (2)
C1	0.69680 (13)	0.72230 (11)	0.28811 (11)	0.0212 (2)
C2	0.59235 (14)	0.72964 (12)	0.16374 (12)	0.0259 (3)
H2	0.4827	0.7163	0.1571	0.031*
C3	0.64691 (15)	0.75616 (13)	0.04969 (12)	0.0305 (3)

H3	0.5748	0.7621	-0.0342	0.037*
C4	0.80703 (15)	0.77406 (13)	0.05803 (13)	0.0304 (3)
H4A	0.8449	0.7896	-0.0203	0.036*
C5	0.91044 (14)	0.76895 (12)	0.18192 (12)	0.0273 (3)
H5	1.0200	0.7824	0.1882	0.033*
C6	0.85757 (13)	0.74464 (11)	0.29708 (12)	0.0234 (2)
H6	0.9304	0.7432	0.3816	0.028*
C7	0.69174 (13)	0.63447 (11)	0.51132 (11)	0.0208 (2)
C8	0.66215 (14)	0.56772 (12)	0.71431 (12)	0.0269 (3)
H8	0.6055	0.5618	0.7825	0.032*
C9	0.79969 (15)	0.51238 (14)	0.73406 (13)	0.0321 (3)
H9	0.8393	0.4706	0.8140	0.039*
C10	0.87631 (14)	0.52129 (13)	0.63096 (13)	0.0309 (3)
H10	0.9700	0.4826	0.6402	0.037*
C11	0.27870 (12)	0.77031 (12)	0.71236 (11)	0.0219 (2)
C12	0.28250 (13)	0.64404 (12)	0.76560 (12)	0.0245 (3)
H12	0.3171	0.5728	0.7199	0.029*
C13	0.23654 (14)	0.62121 (13)	0.88390 (12)	0.0281 (3)
H13	0.2399	0.5349	0.9191	0.034*
C14	0.18554 (15)	0.72450 (14)	0.95093 (13)	0.0322 (3)
H14	0.1532	0.7093	1.0319	0.039*
C15	0.18214 (16)	0.84974 (14)	0.89877 (13)	0.0336 (3)
H15	0.1470	0.9204	0.9447	0.040*
C16	0.22895 (14)	0.87466 (12)	0.78070 (12)	0.0274 (3)
H16	0.2271	0.9617	0.7469	0.033*
C17	0.29837 (13)	0.87077 (11)	0.49672 (11)	0.0220 (2)
C18	0.34955 (14)	0.95120 (12)	0.30445 (12)	0.0277 (3)
H18	0.4052	0.9520	0.2356	0.033*
C19	0.24158 (14)	1.03965 (12)	0.30177 (12)	0.0281 (3)
H19	0.2233	1.1016	0.2342	0.034*
C20	0.16165 (14)	1.03296 (12)	0.40278 (12)	0.0271 (3)
H20	0.0840	1.0901	0.4020	0.033*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0200 (5)	0.0288 (5)	0.0229 (5)	0.0082 (4)	0.0084 (4)	0.0053 (4)
N2	0.0250 (5)	0.0245 (5)	0.0232 (5)	0.0057 (4)	0.0090 (4)	0.0032 (4)
N3	0.0237 (5)	0.0303 (5)	0.0251 (5)	0.0093 (4)	0.0068 (4)	0.0043 (4)
N4	0.0280 (5)	0.0276 (5)	0.0273 (5)	0.0120 (4)	0.0134 (4)	0.0068 (4)
N5	0.0250 (5)	0.0288 (5)	0.0260 (5)	0.0065 (4)	0.0102 (4)	0.0052 (4)
N6	0.0223 (5)	0.0262 (5)	0.0260 (5)	0.0066 (4)	0.0062 (4)	0.0031 (4)
C1	0.0246 (5)	0.0192 (5)	0.0215 (5)	0.0043 (4)	0.0086 (4)	0.0019 (4)
C2	0.0239 (6)	0.0279 (6)	0.0260 (6)	0.0052 (5)	0.0055 (5)	0.0043 (5)
C3	0.0364 (7)	0.0325 (7)	0.0221 (6)	0.0062 (5)	0.0054 (5)	0.0046 (5)
C4	0.0393 (7)	0.0294 (6)	0.0262 (6)	0.0035 (5)	0.0159 (5)	0.0041 (5)
C5	0.0273 (6)	0.0255 (6)	0.0319 (7)	0.0025 (5)	0.0140 (5)	0.0014 (5)
C6	0.0230 (5)	0.0241 (6)	0.0232 (6)	0.0032 (4)	0.0063 (4)	0.0007 (4)

C7	0.0208 (5)	0.0207 (5)	0.0206 (5)	0.0022 (4)	0.0055 (4)	-0.0003 (4)
C8	0.0319 (6)	0.0291 (6)	0.0220 (6)	0.0050 (5)	0.0107 (5)	0.0036 (5)
C9	0.0345 (7)	0.0394 (7)	0.0246 (6)	0.0114 (6)	0.0061 (5)	0.0114 (5)
C10	0.0263 (6)	0.0364 (7)	0.0317 (7)	0.0122 (5)	0.0055 (5)	0.0073 (5)
C11	0.0182 (5)	0.0257 (6)	0.0228 (6)	0.0043 (4)	0.0060 (4)	0.0035 (4)
C12	0.0245 (6)	0.0243 (6)	0.0259 (6)	0.0054 (4)	0.0080 (5)	0.0010 (5)
C13	0.0299 (6)	0.0269 (6)	0.0294 (6)	0.0042 (5)	0.0096 (5)	0.0075 (5)
C14	0.0363 (7)	0.0406 (7)	0.0259 (6)	0.0105 (6)	0.0160 (5)	0.0087 (5)
C15	0.0417 (7)	0.0368 (7)	0.0295 (7)	0.0172 (6)	0.0166 (6)	0.0036 (5)
C16	0.0328 (6)	0.0253 (6)	0.0275 (6)	0.0097 (5)	0.0104 (5)	0.0048 (5)
C17	0.0199 (5)	0.0228 (6)	0.0232 (6)	0.0025 (4)	0.0060 (4)	0.0017 (4)
C18	0.0291 (6)	0.0300 (6)	0.0254 (6)	0.0028 (5)	0.0102 (5)	0.0049 (5)
C19	0.0329 (6)	0.0254 (6)	0.0257 (6)	0.0047 (5)	0.0054 (5)	0.0074 (5)
C20	0.0268 (6)	0.0257 (6)	0.0287 (6)	0.0082 (5)	0.0042 (5)	0.0019 (5)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

N1—C7	1.3613 (14)	C5—H5	0.9500
N1—C1	1.4090 (14)	C6—H6	0.9500
N1—H1	0.892 (9)	C8—C9	1.3829 (17)
N2—C8	1.3294 (15)	C8—H8	0.9500
N2—C7	1.3555 (14)	C9—C10	1.3819 (17)
N3—C10	1.3322 (15)	C9—H9	0.9500
N3—C7	1.3423 (14)	C10—H10	0.9500
N4—C17	1.3584 (15)	C11—C16	1.3950 (16)
N4—C11	1.4093 (14)	C11—C12	1.3963 (16)
N4—H4	0.894 (8)	C12—C13	1.3831 (16)
N5—C18	1.3339 (15)	C12—H12	0.9500
N5—C17	1.3570 (14)	C13—C14	1.3859 (17)
N6—C20	1.3338 (15)	C13—H13	0.9500
N6—C17	1.3407 (14)	C14—C15	1.3818 (18)
C1—C6	1.3949 (15)	C14—H14	0.9500
C1—C2	1.3958 (16)	C15—C16	1.3883 (17)
C2—C3	1.3873 (16)	C15—H15	0.9500
C2—H2	0.9500	C16—H16	0.9500
C3—C4	1.3903 (18)	C18—C19	1.3824 (17)
C3—H3	0.9500	C18—H18	0.9500
C4—C5	1.3839 (18)	C19—C20	1.3824 (17)
C4—H4A	0.9500	C19—H19	0.9500
C5—C6	1.3860 (16)	C20—H20	0.9500
C7—N1—C1	128.19 (9)	C10—C9—H9	121.9
C7—N1—H1	114.6 (10)	C8—C9—H9	121.9
C1—N1—H1	117.0 (10)	N3—C10—C9	122.94 (11)
C8—N2—C7	115.61 (10)	N3—C10—H10	118.5
C10—N3—C7	116.06 (10)	C9—C10—H10	118.5
C17—N4—C11	129.13 (10)	C16—C11—C12	119.15 (10)
C17—N4—H4	115.5 (9)	C16—C11—N4	124.17 (10)

C11—N4—H4	115.4 (9)	C12—C11—N4	116.58 (10)
C18—N5—C17	115.71 (10)	C13—C12—C11	120.91 (11)
C20—N6—C17	115.97 (10)	C13—C12—H12	119.5
C6—C1—C2	119.10 (10)	C11—C12—H12	119.5
C6—C1—N1	123.56 (10)	C12—C13—C14	119.88 (11)
C2—C1—N1	117.30 (10)	C12—C13—H13	120.1
C3—C2—C1	120.70 (11)	C14—C13—H13	120.1
C3—C2—H2	119.7	C15—C14—C13	119.35 (11)
C1—C2—H2	119.7	C15—C14—H14	120.3
C2—C3—C4	120.04 (11)	C13—C14—H14	120.3
C2—C3—H3	120.0	C14—C15—C16	121.49 (11)
C4—C3—H3	120.0	C14—C15—H15	119.3
C5—C4—C3	119.13 (11)	C16—C15—H15	119.3
C5—C4—H4A	120.4	C15—C16—C11	119.20 (11)
C3—C4—H4A	120.4	C15—C16—H16	120.4
C4—C5—C6	121.39 (11)	C11—C16—H16	120.4
C4—C5—H5	119.3	N6—C17—N5	125.94 (11)
C6—C5—H5	119.3	N6—C17—N4	119.51 (10)
C5—C6—C1	119.59 (11)	N5—C17—N4	114.55 (10)
C5—C6—H6	120.2	N5—C18—C19	122.92 (11)
C1—C6—H6	120.2	N5—C18—H18	118.5
N3—C7—N2	125.92 (10)	C19—C18—H18	118.5
N3—C7—N1	119.10 (10)	C18—C19—C20	116.34 (11)
N2—C7—N1	114.96 (10)	C18—C19—H19	121.8
N2—C8—C9	123.22 (11)	C20—C19—H19	121.8
N2—C8—H8	118.4	N6—C20—C19	123.03 (11)
C9—C8—H8	118.4	N6—C20—H20	118.5
C10—C9—C8	116.22 (11)	C19—C20—H20	118.5
C7—N1—C1—C6	-31.08 (18)	C17—N4—C11—C16	-28.92 (18)
C7—N1—C1—C2	151.47 (11)	C17—N4—C11—C12	154.69 (11)
C6—C1—C2—C3	1.23 (17)	C16—C11—C12—C13	0.54 (17)
N1—C1—C2—C3	178.80 (10)	N4—C11—C12—C13	177.12 (10)
C1—C2—C3—C4	0.81 (18)	C11—C12—C13—C14	0.15 (18)
C2—C3—C4—C5	-1.85 (18)	C12—C13—C14—C15	-0.37 (19)
C3—C4—C5—C6	0.85 (18)	C13—C14—C15—C16	-0.1 (2)
C4—C5—C6—C1	1.19 (17)	C14—C15—C16—C11	0.8 (2)
C2—C1—C6—C5	-2.21 (17)	C12—C11—C16—C15	-1.00 (18)
N1—C1—C6—C5	-179.62 (10)	N4—C11—C16—C15	-177.30 (11)
C10—N3—C7—N2	-1.59 (17)	C20—N6—C17—N5	-2.37 (17)
C10—N3—C7—N1	-179.97 (10)	C20—N6—C17—N4	178.70 (10)
C8—N2—C7—N3	1.75 (16)	C18—N5—C17—N6	3.36 (17)
C8—N2—C7—N1	-179.81 (10)	C18—N5—C17—N4	-177.67 (10)
C1—N1—C7—N3	-3.22 (17)	C11—N4—C17—N6	-3.96 (18)
C1—N1—C7—N2	178.23 (10)	C11—N4—C17—N5	176.99 (11)
C7—N2—C8—C9	-0.28 (17)	C17—N5—C18—C19	-1.45 (17)
N2—C8—C9—C10	-1.14 (19)	N5—C18—C19—C20	-1.09 (18)
C7—N3—C10—C9	-0.06 (18)	C17—N6—C20—C19	-0.57 (17)

---

C8—C9—C10—N3

1.3 (2)

C18—C19—C20—N6

2.18 (18)

---

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1···N5	0.89 (1)	2.10 (1)	2.972 (1)	164 (1)
N4—H4···N2	0.89 (1)	2.15 (1)	3.020 (1)	165 (1)

---