

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

**(E)-3-[4-(Pentyloxy)phenyl]-1-phenylprop-2-en-1-one**Asghar Abbas,<sup>a</sup> M. Khawar Rauf,<sup>a</sup> Michael Bolte<sup>b</sup> and Aurangzeb Hasan<sup>a\*</sup><sup>a</sup>Department of Chemistry, Quaid-i-Azam University Islamabad, 45320-Pakistan, and <sup>b</sup>Institut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Str. 7, 60438 Frankfurt/Main, Germany

Correspondence e-mail: flavonoids@hotmail.com


Received 28 April 2009; accepted 4 May 2009

Key indicators: single-crystal X-ray study;  $T = 173$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.101; data-to-parameter ratio = 14.6.

The title compound,  $\text{C}_{20}\text{H}_{22}\text{O}_2$ , crystallizes with two independent molecules in the asymmetric unit. In each molecule, all the non-H atoms lie in a common plane (r.m.s. deviations of 0.098 and 0.079 Å). There is a  $\pi$ - $\pi$  stacking interaction in the crystal structure. The central aromatic rings of the two molecules, which are stacked head-to-tail one above the other, are separated by centroid-to-centroid distances of 3.872 (13) and 3.999 (10) Å.

## Related literature

For background information on chalcones and their properties, see: Achanta *et al.* (2006); Zhang *et al.* (2009); Tran *et al.* (2009); Yagura *et al.* (2008); Sarissky *et al.* (2008); Tang *et al.* (2008); Srivastava *et al.* (2008); For bond-length data, see: Allen *et al.* (1987). For related structures, see: Rosli *et al.* (2006); Harrison *et al.* (2006). For the synthesis, see: Wattanasin & Murphy (1980).



## Experimental

## Crystal data

$\text{C}_{20}\text{H}_{22}\text{O}_2$   
 $M_r = 294.38$   
 Monoclinic,  $P2_1/c$   
 $a = 7.4881$  (4) Å  
 $b = 21.3067$  (11) Å

$c = 20.8328$  (11) Å  
 $\beta = 93.974$  (4)°  
 $V = 3315.8$  (3) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation

$\mu = 0.07$  mm<sup>-1</sup>  
 $T = 173$  K

0.38 × 0.22 × 0.22 mm

## Data collection

Stoe IPDS II two-circle-diffractometer  
 Absorption correction: none  
 25934 measured reflections

5819 independent reflections  
 3319 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.086$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.101$   
 $S = 0.82$   
 5819 reflections

398 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.15$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

Table 1

Selected torsion angles (°).

C11—C1—C2—C3	−177.21 (17)	C11A—C1A—C2A—C3A	−170.05 (18)
C2—C1—C11—C12	−13.4 (3)	C2A—C1A—C11A—C12A	−9.5 (3)
C2—C1—C11—C16	167.85 (17)	C2A—C1A—C11A—C16A	171.52 (17)

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL-Plus* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

AA is grateful to the Higher Education Commission of Pakistan for financial support for his PhD program under scholarship No. IIC-0317109.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2111).

## References

- Achanta, G., Modzelewska, A., Feng, L., Khan, S. R. & Huang, P. (2006). *Mol. Pharmacol.* **70**, 426–433.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Harrison, W. T. A., Yathirajan, H. S., Sarojini, B. K., Narayana, B. & Vijaya Raj, K. K. (2006). *Acta Cryst.* **E62**, o1578–o1579.
- Rosli, M. M., Patil, P. S., Fun, H.-K., Razak, I. A. & Dharmaprakash, S. M. (2006). *Acta Cryst.* **E62**, o1466–o1468.
- Sarissky, M., Pilatova, M., Perjesi, P., Mojzic, J. & Sulla, I. (2008). *Neuro Oncology*, **10**, 1112–1112.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Srivastava, S. S., Joshi, S., Singh, A. R., Yadav, S., Saxena, A. S., Ram, V. J., Chandra, S. & Saxena, J. K. (2008). *Med. Chem. Res.* **17**, 234–244.
- Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.
- Tang, Y. X., Simoneau, A. R., Xie, J., Shahandeh, B. & Zi, X. L. (2008). *Cancer Prevent. Res.* **1**, 439–451.
- Tran, T. D., Park, H., Kim, H. P., Ecker, G. F. & Thai, K. M. (2009). *Bioorg. Med. Chem. Lett.* **19**, 1650–1653.
- Wattanasin, S. & Murphy, W. S. (1980). *Synthesis*, 647–650.
- Yagura, T., Motomiya, T., Ito, M., Honda, G., Iida, A., Kiuchi, F., Tokuda, H. & Nishino, H. (2008). *J. Nat. Med.* **62**, 174–178.
- Zhang, X. D., Hu, X., Hou, A. J. & Wang, H. Y. (2009). *Biol. Pharm. Bull.* **32**, 86–90.

## supporting information

*Acta Cryst.* (2009). E65, o1280 [doi:10.1107/S1600536809016754]

**(E)-3-[4-(Pentyloxy)phenyl]-1-phenylprop-2-en-1-one**

**Asghar Abbas, M. Khawar Rauf, Michael Bolte and Aurangzeb Hasan**

**S1. Comment**

Benzylideneacetophenones ( $\alpha, \beta$ -Unsaturated ketones) comprise a class of synthetic and naturally occurring compounds belonging to the flavonoid family of compounds, commonly known as "*chalcones*". Chalcones and their derivatives are used as precursors for the synthesis of a variety of bioactive organic compounds including heterocyclic compounds. Chalcone analogues have been reported to exhibit potent anticancer activity through inhibition of the proteasome (Achanta *et al.*, 2006), tyrosinase (Zhang *et al.*, 2009) and prostaglandin E-2 (Tran *et al.*, 2009). Naringenin chalcone, the aglycone of isosalipurposide (Yagura *et al.*, 2008) have strong anti-proliferative activity. Some synthetic (Sarissky *et al.*, 2008) and naturally occurring chalcones (Tang *et al.*, 2008) have shown good anticancer activity against Ben-Men-1 human benign meningioma cell line and bladder cancer. Oxygenated chalcones and bischalcones (Srivastava *et al.*, 2008) are a new class of inhibitors of DNA topoisomerase-II of malarial parasites. Here, we report on the crystal structure of the title compound, which was prepared and used as a precursor for the synthesis of heterocyclic compounds.

The molecular structure of the title compound is shown in Fig. 1. The geometrical parameters are normal (Allen *et al.*, 1987) and consistent with those of recently reported chalcone derivatives (Rosli *et al.*, 2006). The asymmetric unit consists of two independent conformers distinctly twisted about the C11—C1 / C11A—C1A and the C1—C2 / C1A—C2A bonds (Table 1), as was also observed for 2-bromo-1-chlorophenyl-3-(4-methoxyphenyl)-2-propen-1-one (Harrison *et al.*, 2006). The dihedral angle between the benzene ring mean planes (C11—C16) and (C21—C26), and (C11A—C16A) and (C21A—C26A) are 12.33 (4) and 7.63 (2)°, respectively. Atoms C1 and O1 deviate from the mean plane (C11—C16) by 0.035 (3) and 0.291 (3) Å, respectively. While atoms C1A and O1A deviated from the mean plane (C11A—C16A) by 0.033 (3) and 0.209 (3) Å, respectively.


In the crystal structure of the title compound the two independent molecules stack head-to-tail (Fig. 2). The central aromatic rings of the two molecules are separated by centroid-to-centroid distances of ca. 3.872 and 3.999 Å.

**S2. Experimental**


The title compound was synthesized by base catalyzed Claisen–Schmidt condensation reaction (Wattanasin *et al.*, 1980). A mixture of acetophenone (1.20 g, 0.01 mol) and 4-(pentyloxy)benzaldehyde (1.92 g, 0.01 mol) was dissolved in ethanol (50 ml) and then 20 ml of an aqueous solution of potassium hydroxide (5%) was added. The mixture was stirred for 3–4 hr, neutralized with dilute HCl and left to stand for 12 hr. The resulting crude solid mass was collected by filtration and recrystallized from ethanol, yielding crystals of the title compound. Full spectroscopic and physical characterization will be reported elsewhere.

**S3. Refinement**


Hydrogen atoms were located in a difference Fourier map but they were included in calculated positions [C-H = 0.95 - 0.99 Å] and refined as riding [ $U_{\text{iso}}(\text{H}) = 1.2$  or  $1.5U_{\text{eq}}(\text{parent C-atom})$ ]

**Figure 1**

a. Molecular structure of molecule A of the title compound in the asymmetric unit. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

b. Molecular structure of the molecule B of the title compound in the asymmetric unit. Displacement ellipsoids are drawn at the 50% probability level.



**Figure 3**

Crystal packing of the title compound with the  $\pi$ - $\pi$ -stacking interactions shown as dashed lines. The hydrogen atoms have been removed for clarity.

**(E)-3-[4-(pentyloxy)phenyl]-1-phenylprop-2-en-1-one**

*Crystal data*

$C_{20}H_{22}O_2$

$M_r = 294.38$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 7.4881(4) \text{ \AA}$

$b = 21.3067(11) \text{ \AA}$

$c = 20.8328(11) \text{ \AA}$

$\beta = 93.974(4)^\circ$

$V = 3315.8(3) \text{ \AA}^3$

$Z = 8$

$F(000) = 1264$

$D_x = 1.179 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 10828 reflections

$\theta = 2.2\text{--}25.4^\circ$

$\mu = 0.07 \text{ mm}^{-1}$

$T = 173$  K  $0.38 \times 0.22 \times 0.22$  mm  
 Needle, light yellow

*Data collection*

STOE IPDS II two-circle-diffractometer	3319 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.086$
Graphite monochromator	$\theta_{\text{max}} = 25.0^\circ$ , $\theta_{\text{min}} = 2.2^\circ$
$\omega$ scans	$h = -8 \rightarrow 8$
25934 measured reflections	$k = -25 \rightarrow 25$
5819 independent reflections	$l = -24 \rightarrow 24$

*Refinement*

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0461P)^2]$
$wR(F^2) = 0.101$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.82$	$(\Delta/\sigma)_{\text{max}} < 0.001$
5819 reflections	$\Delta\rho_{\text{max}} = 0.15 \text{ e } \text{\AA}^{-3}$
398 parameters	$\Delta\rho_{\text{min}} = -0.19 \text{ e } \text{\AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL</i> ,
Primary atom site location: structure-invariant direct methods	$F_c^* = kFc[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.0076 (8)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.3753 (2)	0.28357 (6)	0.50789 (6)	0.0563 (4)
O2	0.36921 (18)	0.62946 (6)	0.31330 (6)	0.0438 (3)
C1	0.4030 (2)	0.27970 (9)	0.45038 (8)	0.0374 (4)
C2	0.4045 (2)	0.33580 (9)	0.40938 (8)	0.0377 (4)
H2	0.4199	0.3314	0.3647	0.045*
C3	0.3843 (2)	0.39303 (9)	0.43424 (9)	0.0391 (4)
H3	0.3695	0.3943	0.4791	0.047*
C4	0.3530 (3)	0.68540 (8)	0.35105 (8)	0.0404 (5)
H4A	0.4567	0.6891	0.3830	0.049*
H4B	0.2425	0.6836	0.3745	0.049*
C5	0.3465 (3)	0.74114 (9)	0.30660 (8)	0.0410 (5)
H5A	0.2375	0.7387	0.2769	0.049*
H5B	0.4519	0.7403	0.2804	0.049*

---

C6	0.3451 (3)	0.80230 (9)	0.34415 (8)	0.0407 (5)
H6A	0.2424	0.8018	0.3716	0.049*
H6B	0.4558	0.8048	0.3729	0.049*
C7	0.3323 (3)	0.86063 (9)	0.30226 (9)	0.0485 (5)
H7A	0.4293	0.8596	0.2723	0.058*
H7B	0.2168	0.8600	0.2761	0.058*
C8	0.3457 (3)	0.92128 (10)	0.34037 (12)	0.0625 (6)
H8A	0.3372	0.9570	0.3107	0.094*
H8B	0.4608	0.9227	0.3658	0.094*
H8C	0.2480	0.9233	0.3692	0.094*
C11	0.4342 (2)	0.21639 (9)	0.42220 (8)	0.0344 (4)
C12	0.4295 (2)	0.20512 (9)	0.35622 (8)	0.0400 (4)
H12	0.4078	0.2388	0.3269	0.048*
C13	0.4562 (3)	0.14507 (9)	0.33307 (9)	0.0481 (5)
H13	0.4541	0.1379	0.2880	0.058*
C14	0.4859 (3)	0.09584 (10)	0.37528 (10)	0.0527 (5)
H14	0.5021	0.0546	0.3593	0.063*
C15	0.4920 (3)	0.10636 (10)	0.44094 (10)	0.0544 (6)
H15	0.5140	0.0725	0.4701	0.065*
C16	0.4661 (3)	0.16618 (9)	0.46385 (9)	0.0463 (5)
H16	0.4701	0.1731	0.5090	0.056*
C21	0.3818 (2)	0.45381 (9)	0.40194 (8)	0.0366 (4)
C22	0.4064 (2)	0.46178 (9)	0.33614 (8)	0.0419 (5)
H22	0.4264	0.4261	0.3103	0.050*
C23	0.4021 (3)	0.52028 (9)	0.30860 (9)	0.0438 (5)
H23	0.4190	0.5246	0.2640	0.053*
C24	0.3730 (2)	0.57350 (9)	0.34540 (8)	0.0373 (4)
C25	0.3491 (3)	0.56698 (9)	0.41056 (8)	0.0422 (5)
H25	0.3308	0.6029	0.4363	0.051*
C26	0.3522 (3)	0.50777 (9)	0.43742 (9)	0.0431 (5)
H26	0.3334	0.5036	0.4818	0.052*
O1A	0.8034 (2)	0.81245 (6)	0.24815 (6)	0.0549 (4)
O2A	0.90510 (18)	0.45190 (6)	0.41506 (5)	0.0432 (3)
C1A	0.8272 (2)	0.80873 (9)	0.30728 (8)	0.0386 (4)
C2A	0.8414 (2)	0.74769 (9)	0.33972 (8)	0.0383 (4)
H2A	0.8414	0.7464	0.3853	0.046*
C3A	0.8545 (2)	0.69393 (9)	0.30771 (8)	0.0386 (4)
H3A	0.8536	0.6969	0.2622	0.046*
C4A	0.9039 (3)	0.39658 (8)	0.37602 (8)	0.0407 (4)
H4A1	1.0081	0.3967	0.3493	0.049*
H4A2	0.7933	0.3951	0.3471	0.049*
C5A	0.9124 (3)	0.34046 (9)	0.42018 (9)	0.0422 (5)
H5A1	0.8057	0.3402	0.4457	0.051*
H5A2	1.0198	0.3438	0.4505	0.051*
C6A	0.9201 (3)	0.27920 (8)	0.38292 (9)	0.0448 (5)
H6A1	0.8104	0.2755	0.3538	0.054*
H6A2	1.0237	0.2806	0.3559	0.054*
C7A	0.9362 (3)	0.22144 (10)	0.42529 (11)	0.0575 (6)

H7A1	0.8292	0.2185	0.4505	0.069*
H7A2	1.0424	0.2259	0.4560	0.069*
C8A	0.9535 (3)	0.16114 (10)	0.38681 (14)	0.0798 (8)
H8A1	0.9630	0.1253	0.4163	0.120*
H8A2	1.0609	0.1633	0.3626	0.120*
H8A3	0.8477	0.1561	0.3568	0.120*
C11A	0.8432 (2)	0.86778 (9)	0.34646 (8)	0.0364 (4)
C12A	0.8455 (3)	0.86841 (9)	0.41317 (8)	0.0441 (5)
H12A	0.8404	0.8300	0.4361	0.053*
C13A	0.8553 (3)	0.92462 (10)	0.44644 (10)	0.0533 (5)
H13A	0.8548	0.9246	0.4920	0.064*
C14A	0.8656 (3)	0.98053 (10)	0.41389 (10)	0.0525 (5)
H14A	0.8725	1.0190	0.4370	0.063*
C15A	0.8661 (3)	0.98060 (9)	0.34781 (10)	0.0512 (5)
H15A	0.8751	1.0192	0.3254	0.061*
C16A	0.8533 (3)	0.92468 (9)	0.31398 (9)	0.0437 (5)
H16A	0.8514	0.9251	0.2683	0.052*
C21A	0.8699 (2)	0.63097 (9)	0.33500 (8)	0.0367 (4)
C22A	0.8770 (3)	0.62003 (9)	0.40134 (8)	0.0412 (5)
H22A	0.8715	0.6546	0.4299	0.049*
C23A	0.8918 (3)	0.56028 (9)	0.42609 (9)	0.0431 (5)
H23A	0.8994	0.5542	0.4714	0.052*
C24A	0.8957 (2)	0.50891 (8)	0.38534 (8)	0.0361 (4)
C25A	0.8902 (2)	0.51804 (9)	0.31898 (8)	0.0405 (5)
H25A	0.8943	0.4833	0.2906	0.049*
C26A	0.8787 (3)	0.57867 (9)	0.29517 (8)	0.0420 (5)
H26A	0.8766	0.5849	0.2500	0.050*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0865 (11)	0.0481 (9)	0.0348 (7)	0.0088 (8)	0.0074 (7)	-0.0041 (6)
O2	0.0578 (8)	0.0340 (8)	0.0403 (7)	-0.0030 (6)	0.0070 (6)	-0.0019 (6)
C1	0.0362 (10)	0.0384 (11)	0.0372 (10)	0.0006 (8)	0.0002 (8)	-0.0023 (8)
C2	0.0400 (11)	0.0369 (11)	0.0365 (9)	0.0005 (9)	0.0039 (8)	-0.0026 (8)
C3	0.0373 (10)	0.0402 (12)	0.0398 (10)	-0.0013 (9)	0.0016 (8)	-0.0026 (8)
C4	0.0469 (11)	0.0327 (11)	0.0419 (10)	-0.0040 (9)	0.0052 (8)	-0.0031 (8)
C5	0.0434 (11)	0.0382 (11)	0.0415 (10)	-0.0026 (9)	0.0034 (8)	0.0014 (8)
C6	0.0416 (11)	0.0379 (11)	0.0428 (10)	-0.0014 (9)	0.0033 (8)	0.0011 (8)
C7	0.0478 (12)	0.0419 (12)	0.0556 (12)	-0.0010 (10)	0.0019 (9)	0.0054 (9)
C8	0.0582 (14)	0.0398 (13)	0.0881 (16)	0.0016 (11)	-0.0051 (12)	0.0027 (12)
C11	0.0310 (9)	0.0323 (10)	0.0401 (9)	0.0001 (8)	0.0037 (7)	-0.0001 (8)
C12	0.0456 (11)	0.0365 (11)	0.0380 (9)	0.0028 (9)	0.0036 (8)	-0.0008 (8)
C13	0.0592 (13)	0.0392 (12)	0.0470 (11)	0.0030 (10)	0.0114 (9)	-0.0082 (9)
C14	0.0604 (14)	0.0357 (12)	0.0638 (13)	0.0038 (10)	0.0168 (10)	-0.0028 (10)
C15	0.0715 (15)	0.0389 (12)	0.0540 (12)	0.0060 (11)	0.0131 (11)	0.0108 (10)
C16	0.0553 (12)	0.0422 (12)	0.0425 (10)	0.0025 (10)	0.0104 (9)	0.0017 (9)
C21	0.0343 (10)	0.0344 (11)	0.0409 (10)	-0.0019 (8)	0.0031 (8)	-0.0026 (8)

C22	0.0464 (11)	0.0390 (12)	0.0410 (10)	-0.0002 (9)	0.0079 (8)	-0.0077 (9)
C23	0.0549 (12)	0.0406 (12)	0.0369 (10)	-0.0022 (10)	0.0099 (9)	-0.0036 (9)
C24	0.0380 (10)	0.0343 (11)	0.0400 (10)	-0.0042 (8)	0.0046 (8)	0.0012 (8)
C25	0.0527 (12)	0.0341 (11)	0.0405 (10)	-0.0031 (9)	0.0080 (8)	-0.0071 (8)
C26	0.0545 (12)	0.0387 (12)	0.0368 (10)	-0.0040 (10)	0.0077 (9)	-0.0038 (8)
O1A	0.0860 (11)	0.0418 (8)	0.0360 (7)	0.0002 (8)	-0.0008 (7)	0.0018 (6)
O2A	0.0588 (9)	0.0313 (8)	0.0397 (7)	0.0033 (6)	0.0043 (6)	0.0012 (6)
C1A	0.0390 (11)	0.0364 (11)	0.0404 (10)	0.0007 (9)	0.0022 (8)	0.0015 (8)
C2A	0.0435 (11)	0.0351 (11)	0.0364 (9)	0.0001 (9)	0.0035 (8)	0.0001 (8)
C3A	0.0416 (11)	0.0381 (11)	0.0360 (9)	0.0008 (9)	0.0015 (8)	0.0020 (8)
C4A	0.0464 (11)	0.0333 (11)	0.0418 (10)	0.0007 (9)	-0.0011 (8)	-0.0022 (8)
C5A	0.0406 (11)	0.0402 (12)	0.0460 (11)	-0.0001 (9)	0.0042 (8)	0.0038 (9)
C6A	0.0455 (11)	0.0323 (11)	0.0557 (11)	-0.0029 (9)	-0.0018 (9)	0.0051 (9)
C7A	0.0435 (12)	0.0408 (13)	0.0879 (15)	-0.0026 (10)	0.0012 (11)	0.0214 (11)
C8A	0.0639 (16)	0.0321 (13)	0.140 (2)	-0.0054 (12)	-0.0151 (15)	0.0103 (14)
C11A	0.0342 (10)	0.0334 (11)	0.0416 (10)	-0.0008 (8)	0.0034 (8)	0.0013 (8)
C12A	0.0568 (13)	0.0344 (11)	0.0408 (10)	-0.0011 (10)	0.0012 (9)	0.0023 (8)
C13A	0.0693 (15)	0.0438 (13)	0.0464 (11)	-0.0047 (11)	0.0003 (10)	-0.0061 (10)
C14A	0.0574 (13)	0.0383 (13)	0.0618 (14)	-0.0052 (10)	0.0042 (10)	-0.0089 (10)
C15A	0.0555 (13)	0.0343 (12)	0.0639 (13)	-0.0018 (10)	0.0055 (10)	0.0062 (10)
C16A	0.0493 (12)	0.0364 (11)	0.0453 (11)	-0.0014 (9)	0.0035 (9)	0.0052 (9)
C21A	0.0387 (10)	0.0334 (11)	0.0381 (10)	0.0021 (8)	0.0033 (8)	-0.0004 (8)
C22A	0.0544 (12)	0.0344 (11)	0.0353 (9)	0.0040 (9)	0.0067 (8)	-0.0049 (8)
C23A	0.0582 (13)	0.0377 (12)	0.0343 (9)	0.0056 (9)	0.0098 (8)	0.0011 (8)
C24A	0.0386 (11)	0.0334 (11)	0.0364 (10)	0.0035 (8)	0.0026 (8)	0.0016 (8)
C25A	0.0498 (12)	0.0326 (11)	0.0387 (10)	0.0029 (9)	0.0009 (8)	-0.0051 (8)
C26A	0.0538 (12)	0.0390 (12)	0.0330 (9)	0.0031 (9)	0.0025 (8)	-0.0008 (8)

*Geometric parameters (Å, °)*

O1—C1	1.2326 (19)	O1A—C1A	1.235 (2)
O2—C24	1.366 (2)	O2A—C24A	1.363 (2)
O2—C4	1.438 (2)	O2A—C4A	1.432 (2)
C1—C2	1.470 (3)	C1A—C2A	1.466 (3)
C1—C11	1.496 (3)	C1A—C11A	1.500 (3)
C2—C3	1.338 (3)	C2A—C3A	1.332 (2)
C2—H2	0.9500	C2A—H2A	0.9500
C3—C21	1.459 (3)	C3A—C21A	1.459 (3)
C3—H3	0.9500	C3A—H3A	0.9500
C4—C5	1.505 (2)	C4A—C5A	1.507 (2)
C4—H4A	0.9900	C4A—H4A1	0.9900
C4—H4B	0.9900	C4A—H4A2	0.9900
C5—C6	1.520 (3)	C5A—C6A	1.522 (3)
C5—H5A	0.9900	C5A—H5A1	0.9900
C5—H5B	0.9900	C5A—H5A2	0.9900
C6—C7	1.518 (3)	C6A—C7A	1.515 (3)
C6—H6A	0.9900	C6A—H6A1	0.9900
C6—H6B	0.9900	C6A—H6A2	0.9900



C7—C8	1.516 (3)	C7A—C8A	1.525 (3)
C7—H7A	0.9900	C7A—H7A1	0.9900
C7—H7B	0.9900	C7A—H7A2	0.9900
C8—H8A	0.9800	C8A—H8A1	0.9800
C8—H8B	0.9800	C8A—H8A2	0.9800
C8—H8C	0.9800	C8A—H8A3	0.9800
C11—C16	1.388 (3)	C11A—C12A	1.389 (2)
C11—C12	1.393 (2)	C11A—C16A	1.393 (3)
C12—C13	1.387 (3)	C12A—C13A	1.383 (3)
C12—H12	0.9500	C12A—H12A	0.9500
C13—C14	1.377 (3)	C13A—C14A	1.376 (3)
C13—H13	0.9500	C13A—H13A	0.9500
C14—C15	1.384 (3)	C14A—C15A	1.377 (3)
C14—H14	0.9500	C14A—H14A	0.9500
C15—C16	1.379 (3)	C15A—C16A	1.384 (3)
C15—H15	0.9500	C15A—H15A	0.9500
C16—H16	0.9500	C16A—H16A	0.9500
C21—C26	1.393 (3)	C21A—C26A	1.394 (3)
C21—C22	1.406 (2)	C21A—C22A	1.399 (2)
C22—C23	1.372 (3)	C22A—C23A	1.375 (3)
C22—H22	0.9500	C22A—H22A	0.9500
C23—C24	1.394 (3)	C23A—C24A	1.387 (2)
C23—H23	0.9500	C23A—H23A	0.9500
C24—C25	1.388 (2)	C24A—C25A	1.394 (2)
C25—C26	1.380 (3)	C25A—C26A	1.384 (3)
C25—H25	0.9500	C25A—H25A	0.9500
C26—H26	0.9500	C26A—H26A	0.9500
C24—O2—C4	117.09 (13)	C24A—O2A—C4A	118.52 (13)
O1—C1—C2	121.27 (17)	O1A—C1A—C2A	121.18 (17)
O1—C1—C11	118.89 (16)	O1A—C1A—C11A	119.30 (17)
C2—C1—C11	119.83 (15)	C2A—C1A—C11A	119.52 (16)
C3—C2—C1	120.71 (16)	C3A—C2A—C1A	122.50 (16)
C3—C2—H2	119.6	C3A—C2A—H2A	118.8
C1—C2—H2	119.6	C1A—C2A—H2A	118.8
C2—C3—C21	128.96 (17)	C2A—C3A—C21A	127.02 (17)
C2—C3—H3	115.5	C2A—C3A—H3A	116.5
C21—C3—H3	115.5	C21A—C3A—H3A	116.5
O2—C4—C5	108.55 (14)	O2A—C4A—C5A	107.93 (14)
O2—C4—H4A	110.0	O2A—C4A—H4A1	110.1
C5—C4—H4A	110.0	C5A—C4A—H4A1	110.1
O2—C4—H4B	110.0	O2A—C4A—H4A2	110.1
C5—C4—H4B	110.0	C5A—C4A—H4A2	110.1
H4A—C4—H4B	108.4	H4A1—C4A—H4A2	108.4
C4—C5—C6	111.17 (15)	C4A—C5A—C6A	111.76 (15)
C4—C5—H5A	109.4	C4A—C5A—H5A1	109.3
C6—C5—H5A	109.4	C6A—C5A—H5A1	109.3
C4—C5—H5B	109.4	C4A—C5A—H5A2	109.3

C6—C5—H5B	109.4	C6A—C5A—H5A2	109.3
H5A—C5—H5B	108.0	H5A1—C5A—H5A2	107.9
C7—C6—C5	114.12 (15)	C7A—C6A—C5A	113.83 (16)
C7—C6—H6A	108.7	C7A—C6A—H6A1	108.8
C5—C6—H6A	108.7	C5A—C6A—H6A1	108.8
C7—C6—H6B	108.7	C7A—C6A—H6A2	108.8
C5—C6—H6B	108.7	C5A—C6A—H6A2	108.8
H6A—C6—H6B	107.6	H6A1—C6A—H6A2	107.7
C8—C7—C6	113.43 (17)	C6A—C7A—C8A	112.62 (19)
C8—C7—H7A	108.9	C6A—C7A—H7A1	109.1
C6—C7—H7A	108.9	C8A—C7A—H7A1	109.1
C8—C7—H7B	108.9	C6A—C7A—H7A2	109.1
C6—C7—H7B	108.9	C8A—C7A—H7A2	109.1
H7A—C7—H7B	107.7	H7A1—C7A—H7A2	107.8
C7—C8—H8A	109.5	C7A—C8A—H8A1	109.5
C7—C8—H8B	109.5	C7A—C8A—H8A2	109.5
H8A—C8—H8B	109.5	H8A1—C8A—H8A2	109.5
C7—C8—H8C	109.5	C7A—C8A—H8A3	109.5
H8A—C8—H8C	109.5	H8A1—C8A—H8A3	109.5
H8B—C8—H8C	109.5	H8A2—C8A—H8A3	109.5
C16—C11—C12	118.38 (17)	C12A—C11A—C16A	118.70 (18)
C16—C11—C1	118.33 (15)	C12A—C11A—C1A	123.21 (17)
C12—C11—C1	123.28 (16)	C16A—C11A—C1A	118.08 (16)
C13—C12—C11	120.51 (18)	C13A—C12A—C11A	120.43 (18)
C13—C12—H12	119.7	C13A—C12A—H12A	119.8
C11—C12—H12	119.7	C11A—C12A—H12A	119.8
C14—C13—C12	120.10 (18)	C14A—C13A—C12A	120.37 (19)
C14—C13—H13	119.9	C14A—C13A—H13A	119.8
C12—C13—H13	119.9	C12A—C13A—H13A	119.8
C13—C14—C15	120.05 (19)	C13A—C14A—C15A	119.9 (2)
C13—C14—H14	120.0	C13A—C14A—H14A	120.1
C15—C14—H14	120.0	C15A—C14A—H14A	120.1
C16—C15—C14	119.72 (19)	C14A—C15A—C16A	120.21 (19)
C16—C15—H15	120.1	C14A—C15A—H15A	119.9
C14—C15—H15	120.1	C16A—C15A—H15A	119.9
C15—C16—C11	121.22 (18)	C15A—C16A—C11A	120.40 (18)
C15—C16—H16	119.4	C15A—C16A—H16A	119.8
C11—C16—H16	119.4	C11A—C16A—H16A	119.8
C26—C21—C22	117.01 (17)	C26A—C21A—C22A	117.06 (17)
C26—C21—C3	119.03 (16)	C26A—C21A—C3A	120.60 (16)
C22—C21—C3	123.96 (17)	C22A—C21A—C3A	122.34 (16)
C23—C22—C21	121.11 (17)	C23A—C22A—C21A	121.41 (17)
C23—C22—H22	119.4	C23A—C22A—H22A	119.3
C21—C22—H22	119.4	C21A—C22A—H22A	119.3
C22—C23—C24	120.61 (17)	C22A—C23A—C24A	120.38 (17)
C22—C23—H23	119.7	C22A—C23A—H23A	119.8
C24—C23—H23	119.7	C24A—C23A—H23A	119.8
O2—C24—C25	124.47 (17)	O2A—C24A—C23A	115.34 (15)

O2—C24—C23	116.06 (15)	O2A—C24A—C25A	124.88 (16)
C25—C24—C23	119.47 (18)	C23A—C24A—C25A	119.78 (17)
C26—C25—C24	119.22 (17)	C26A—C25A—C24A	118.84 (17)
C26—C25—H25	120.4	C26A—C25A—H25A	120.6
C24—C25—H25	120.4	C24A—C25A—H25A	120.6
C25—C26—C21	122.57 (17)	C25A—C26A—C21A	122.49 (17)
C25—C26—H26	118.7	C25A—C26A—H26A	118.8
C21—C26—H26	118.7	C21A—C26A—H26A	118.8
O1—C1—C2—C3	3.2 (3)	O1A—C1A—C2A—C3A	9.5 (3)
C11—C1—C2—C3	-177.21 (17)	C11A—C1A—C2A—C3A	-170.05 (18)
C1—C2—C3—C21	-179.97 (18)	C1A—C2A—C3A—C21A	179.74 (18)
C24—O2—C4—C5	-178.53 (15)	C24A—O2A—C4A—C5A	-179.11 (16)
O2—C4—C5—C6	-175.23 (15)	O2A—C4A—C5A—C6A	-177.50 (16)
C4—C5—C6—C7	-178.06 (17)	C4A—C5A—C6A—C7A	177.72 (17)
C5—C6—C7—C8	-175.46 (18)	C5A—C6A—C7A—C8A	-176.95 (18)
O1—C1—C11—C16	-12.6 (3)	O1A—C1A—C11A—C12A	170.93 (18)
C2—C1—C11—C12	-13.4 (3)	C2A—C1A—C11A—C12A	-9.5 (3)
C2—C1—C11—C16	167.85 (17)	O1A—C1A—C11A—C16A	-8.0 (3)
O1—C1—C11—C12	166.19 (17)	C2A—C1A—C11A—C16A	171.52 (17)
C16—C11—C12—C13	0.0 (3)	C16A—C11A—C12A—C13A	0.8 (3)
C1—C11—C12—C13	-178.79 (18)	C1A—C11A—C12A—C13A	-178.11 (19)
C11—C12—C13—C14	0.7 (3)	C11A—C12A—C13A—C14A	-1.0 (3)
C12—C13—C14—C15	-1.1 (3)	C12A—C13A—C14A—C15A	0.1 (3)
C13—C14—C15—C16	0.8 (3)	C13A—C14A—C15A—C16A	1.0 (3)
C14—C15—C16—C11	-0.2 (3)	C14A—C15A—C16A—C11A	-1.2 (3)
C12—C11—C16—C15	-0.2 (3)	C12A—C11A—C16A—C15A	0.3 (3)
C1—C11—C16—C15	178.60 (18)	C1A—C11A—C16A—C15A	179.26 (18)
C2—C3—C21—C26	177.7 (2)	C2A—C3A—C21A—C26A	178.60 (19)
C2—C3—C21—C22	-1.8 (3)	C2A—C3A—C21A—C22A	-1.5 (3)
C26—C21—C22—C23	0.3 (3)	C26A—C21A—C22A—C23A	-0.1 (3)
C3—C21—C22—C23	179.79 (19)	C3A—C21A—C22A—C23A	-179.92 (18)
C21—C22—C23—C24	0.0 (3)	C21A—C22A—C23A—C24A	-1.6 (3)
C4—O2—C24—C25	5.0 (3)	C4A—O2A—C24A—C23A	178.06 (17)
C4—O2—C24—C23	-175.43 (17)	C4A—O2A—C24A—C25A	-2.0 (3)
C22—C23—C24—O2	-179.33 (17)	C22A—C23A—C24A—O2A	-178.06 (17)
C22—C23—C24—C25	0.3 (3)	C22A—C23A—C24A—C25A	2.0 (3)
O2—C24—C25—C26	178.69 (18)	O2A—C24A—C25A—C26A	179.28 (17)
C23—C24—C25—C26	-0.9 (3)	C23A—C24A—C25A—C26A	-0.8 (3)
C24—C25—C26—C21	1.2 (3)	C24A—C25A—C26A—C21A	-0.9 (3)
C22—C21—C26—C25	-0.9 (3)	C22A—C21A—C26A—C25A	1.3 (3)
C3—C21—C26—C25	179.55 (18)	C3A—C21A—C26A—C25A	-178.83 (18)