

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(5,6-Dihydrobenzimidazolo[1,2-c]-quinazolin-6-yl)aniline methanol solvate

 Irvin Booyesen,^a Thomas I. A. Gerber,^{a*} Eric Hosten^a and Peter Mayer^b
^aDepartment of Chemistry, Nelson Mandela Metropolitan University, 6031 Port Elizabeth, South Africa, and ^bDepartment of Chemistry, Ludwig-Maximilians University, D-81377 München, Germany

Correspondence e-mail: thomas.gerber@nmmu.ac.za


Received 28 May 2009; accepted 9 June 2009

 Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.048; wR factor = 0.123; data-to-parameter ratio = 12.4.

In the structure of the title compound, $\text{C}_{20}\text{H}_{16}\text{N}_4 \cdot \text{CH}_4\text{O}$, the aniline ring forms dihedral angles of 89.9 (2) and 85.4 (2)° with the benzimidazole and benzene rings, respectively. The orientation of the aniline ring is mainly determined by strong hydrogen bonds between the amino group and the non-fused quinazoline N atom. Intermolecular hydrogen bonds of the $\text{N}-\text{H} \cdots \text{N}-\text{H} \cdots \text{N}$ type along [010] and the $\text{N}-\text{H} \cdots \text{O}-\text{H} \cdots \text{N}$ type along [100] are formed, resulting in $\text{C}_2^2(4)$ and $\text{C}_2^2(10)$ descriptors, respectively, on a binary level of graph-set analysis. There are $\text{C}-\text{H} \cdots \pi$ contacts with $\text{H} \cdots \pi$ distances of 2.44 Å; however, no π -stacking is observed.

Related literature

For the synthesis of quinazolines, see: Kubicova *et al.* (2003); Niementowski (1895). For the conformation, see: Cuny *et al.* (1980); Williamson (1957).



Experimental

Crystal data

 $\text{C}_{20}\text{H}_{16}\text{N}_4 \cdot \text{CH}_4\text{O}$
 $M_r = 344.41$
 Monoclinic, $P2_1/c$
 $a = 9.3703$ (2) Å
 $b = 5.1728$ (1) Å
 $c = 35.5169$ (9) Å

 $\beta = 91.3908$ (14)°
 $V = 1721.02$ (7) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

 $\mu = 0.09$ mm⁻¹
 $T = 200$ K
 $0.23 \times 0.06 \times 0.05$ mm

Data collection

 Nonius KappaCCD diffractometer
 Absorption correction: none
 9302 measured reflections

 3136 independent reflections
 2089 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.123$
 $S = 1.08$
 3136 reflections
 253 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3a} \cdots \text{N4}^{\text{i}}$	0.93 (2)	2.13 (2)	3.058 (3)	174.5 (18)
$\text{N4}-\text{H4a} \cdots \text{O1}^{\text{ii}}$	0.95 (2)	2.00 (3)	2.946 (3)	174.4 (19)
$\text{N4}-\text{H4b} \cdots \text{N3}$	0.86 (2)	2.35 (2)	3.006 (3)	133.8 (19)
$\text{O1}-\text{H1} \cdots \text{N1}$	0.95 (3)	1.88 (3)	2.814 (2)	167 (3)
$\text{C14}-\text{H14} \cdots \text{Cg}^{\text{i}}$	1.00	2.44	3.408 (2)	162

 Symmetry codes: (i) $x, y - 1, z$; (ii) $x + 1, y, z$. Cg is the centroid of the C1–C6 ring.

Data collection: *COLLECT* (Nonius, 2004); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *PARST* (Nardelli, 1995), *pubCIF* (Westrip, 2008) and *WinGX* (Farrugia, 1999).

The authors thank Professor P. Klüfers for generous allocation of diffractometer time.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2264).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Cuny, E., Lichtenthaler, F. W. & Moser, A. (1980). *Tetrahedron Lett.* **21**, 3029–3032.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Kubicova, L., Sustr, M., Kralova, K., Chobnot, V., Vytlačilova, J., Jahodar, L., Vuorela, P., Machacek, M. & Kaustova, J. (2003). *Molecules*, **8**, 756–769.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Niementowski, S. (1895). *J. Prakt. Chem.* **51**, 546–566.
- Nonius (2004). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2008). *pubCIF*. In preparation.
- Williamson, T. A. (1957). *Heterocycl. Compd.* **6**, 331–339.

supporting information

Acta Cryst. (2009). E65, o1565 [doi:10.1107/S1600536809021886]

2-(5,6-Dihydrobenzimidazo[1,2-*c*]quinazolin-6-yl)aniline methanol solvate**Irvin Booyen, Thomas I. A. Gerber, Eric Hosten and Peter Mayer****S1. Comment**

In the present work the structure of 2-(2,3,5,6-tetrahydrobenzimidazo[1,2-*c*]quinazolin-5-yl)benzenamine (Figure 1) has been determined to explore its suitability as a bidentate ligand for various metal ions. In the structure the quinazoline ring adopts a chair conformation: atoms C7, C8, C13, N2 and N3 are coplanar, with atom C14 departing from the plane by 0.2391 Å. The orientation of the aniline ring is mainly determined by a series of hydrogen-bonds between NH₂ and NH groups, and between the phenylamino group and the quinazoline nitrogen (Figure 1 and Table 1). This ring makes dihedral angles of 89.9 (2)° and 85.4 (2)° with the benzimidazole and phenyl rings respectively. The ligand bond distances and angles show that N1—C7 is a localized double bond [1.321 (3) Å], with N2—C7 a single bond at 1.383 (3) Å. The N3—C14 bond length is 1.459 (3) Å, and the N3—C14—N2 bond angle [108.14 (17)°] illustrates the *sp*³ hybridization of C14. All the other bond lengths and angles in the molecule are normal.


Intermolecular hydrogen bonds of the N—H...N—H...N type along [010] and N—H...O—H...N along [100] are formed (Table 1) resulting in a C₂²(4) descriptor and a C₂²(10) descriptor on a binary level of graph set analysis, respectively. C—H... π contacts with H...C_g distances of 2.44 Å are present in the structure (C_g is the centre of gravity of ring C1 – C6); however, no π -stacking is observed.

S2. Experimental

All chemicals used (reagent grade) were commercially available. A mass of 1.22 g (0.010 mol) of 2-aminobenzaldehyde was dissolved in methanol (50 cm³), and 2.09 g (0.010 mol) of 2-(2-aminophenyl)-1-benzimidazole was added with stirring. The mixture was heated under reflux for 2 h, then cooled to room temperature and filtered. The volume of the solution was reduced to ~ 10 cm³, and left to evaporate slowly at room temperature. After 2 days 2.48 g (72%) of colourless crystals, with the formulation C₂₀H₁₆N₄.CH₄O and suitable for X-ray analysis, were collected. *M.p.* 211°C. ¹H NMR (300 MHz, d₆-DMSO): 7.97 (1H, d), 7.62 (1H, d), 7.27 (1H, d), 7.25 (1H, t), 7.13 (2H, t), 6.97–7.02 (2H, m), 6.94 (1H, t), 6.86 (1H, t), 6.78 (1H, d), 6.61 (1H, d), 6.54 (1H, t), 5.45 (2H, s), 3.18 (3H, s).

S3. Refinement

All H atoms bonded to C atoms were calculated in idealized position and refined as riding on their parent atoms with *U*_{iso}(H) values of 1.2 *U*_{eq}(C). All H atoms bonded to N and O atoms were refined freely with individual *U*_{iso}(H) values.

**Figure 1**

The molecular structure of the title compound (anisotropic displacement ellipsoids drawn at the 50% probability level). Hydrogen bonds determining the conformational arrangement of the aniline rings are also shown. Symmetry codes: (i): $x, y+1, z$; (ii) $x+1, y, z$.

2-(5,6-Dihydrobenzimidazo[1,2-c]quinazolin-6-yl)aniline methanol solvate

Crystal data

$C_{20}H_{16}N_4 \cdot CH_4O$

$M_r = 344.41$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.3703(2)\ \text{\AA}$

$b = 5.1728(1)\ \text{\AA}$

$c = 35.5169(9)\ \text{\AA}$

$\beta = 91.3908(14)^\circ$

$V = 1721.02(7)\ \text{\AA}^3$

$Z = 4$

$F(000) = 728$

$D_x = 1.329(1)\ \text{Mg m}^{-3}$

Melting point: 484 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 41046 reflections

$\theta = 3.1\text{--}25.4^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 200\ \text{K}$

Rod, yellow

$0.23 \times 0.06 \times 0.05\ \text{mm}$

Data collection

Nonius KappaCCD
diffractometer

Radiation source: rotating anode

MONTEL, graded multilayered X-ray optics
monochromator

Detector resolution: 9 pixels mm^{-1}

CCD; rotation images, φ and ω scans

9302 measured reflections

3136 independent reflections

2089 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$
 $\theta_{\text{max}} = 25.4^\circ$, $\theta_{\text{min}} = 3.2^\circ$
 $h = -11 \rightarrow 11$

$k = -5 \rightarrow 6$
 $l = -42 \rightarrow 42$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.123$
 $S = 1.08$
 3136 reflections
 253 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0417P)^2 + 0.5855P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.20 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008)
 Extinction coefficient: 0.0138 (19)

Special details

Refinement. N- and O-bonded H: All H-atom parameters refined C-bonded H: H-atom parameters constrained

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.05660 (18)	0.5134 (3)	0.32314 (5)	0.0506 (5)
H1	0.038 (3)	0.482 (5)	0.3324 (8)	0.087 (10)*
N1	0.20470 (18)	0.3599 (3)	0.35685 (5)	0.0349 (5)
N2	0.38507 (17)	0.1319 (3)	0.38325 (5)	0.0321 (5)
N3	0.5138 (2)	-0.1946 (4)	0.35148 (5)	0.0407 (5)
H3a	0.564 (2)	-0.350 (5)	0.3544 (6)	0.043 (7)*
N4	0.6823 (2)	0.2995 (4)	0.35560 (6)	0.0381 (5)
H4a	0.764 (3)	0.364 (4)	0.3436 (6)	0.051 (7)*
H4b	0.635 (2)	0.195 (4)	0.3412 (6)	0.040 (7)*
C1	0.2283 (2)	0.4419 (4)	0.39363 (6)	0.0329 (5)
C2	0.1591 (2)	0.6363 (4)	0.41352 (6)	0.0394 (6)
H2	0.0863	0.7387	0.4019	0.047*
C3	0.2005 (2)	0.6737 (5)	0.45063 (7)	0.0437 (6)
H3	0.1545	0.8037	0.4648	0.052*
C4	0.3083 (3)	0.5254 (5)	0.46794 (6)	0.0443 (6)
H4	0.3332	0.5557	0.4937	0.053*
C5	0.3792 (2)	0.3361 (4)	0.44846 (6)	0.0393 (6)
H5	0.4528	0.2360	0.4601	0.047*
C6	0.3383 (2)	0.2981 (4)	0.41099 (6)	0.0317 (5)
C7	0.2981 (2)	0.1726 (4)	0.35178 (6)	0.0307 (5)
C8	0.3146 (2)	0.0110 (4)	0.31914 (6)	0.0321 (5)
C9	0.2270 (2)	0.0334 (5)	0.28675 (6)	0.0410 (6)
H9	0.1572	0.1664	0.2853	0.049*
C10	0.2408 (3)	-0.1347 (5)	0.25705 (6)	0.0475 (7)
H10	0.1800	-0.1208	0.2354	0.057*
C11	0.3452 (3)	-0.3251 (5)	0.25931 (6)	0.0452 (6)

H11	0.3552	-0.4419	0.2389	0.054*
C12	0.4343 (2)	-0.3483 (4)	0.29040 (6)	0.0393 (6)
H12	0.5060	-0.4781	0.2912	0.047*
C13	0.4193 (2)	-0.1808 (4)	0.32092 (6)	0.0333 (5)
C14	0.4874 (2)	-0.0827 (4)	0.38838 (6)	0.0342 (5)
H14	0.4431	-0.2178	0.4045	0.041*
C15	0.6265 (2)	0.0014 (4)	0.40688 (6)	0.0322 (5)
C16	0.7166 (2)	0.1831 (4)	0.39045 (6)	0.0337 (5)
C17	0.8413 (2)	0.2541 (5)	0.40991 (7)	0.0423 (6)
H17	0.9013	0.3825	0.3996	0.051*
C18	0.8792 (3)	0.1423 (5)	0.44374 (7)	0.0478 (7)
H18	0.9647	0.1948	0.4565	0.057*
C19	0.7947 (3)	-0.0451 (5)	0.45936 (7)	0.0469 (6)
H19	0.8225	-0.1265	0.4824	0.056*
C20	0.6682 (2)	-0.1127 (4)	0.44085 (6)	0.0408 (6)
H20	0.6086	-0.2400	0.4516	0.049*
C21	-0.0816 (3)	0.7792 (5)	0.32160 (8)	0.0677 (8)
H21A	-0.0188	0.8671	0.3400	0.102*
H21B	-0.1814	0.8141	0.3275	0.102*
H21C	-0.0623	0.8430	0.2962	0.102*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0392 (11)	0.0532 (12)	0.0589 (11)	0.0000 (9)	-0.0085 (8)	0.0070 (9)
N1	0.0293 (10)	0.0331 (11)	0.0421 (12)	0.0012 (9)	-0.0009 (8)	-0.0012 (9)
N2	0.0292 (10)	0.0293 (10)	0.0378 (11)	0.0002 (8)	-0.0033 (8)	-0.0017 (8)
N3	0.0379 (12)	0.0347 (12)	0.0491 (13)	0.0077 (10)	-0.0095 (9)	-0.0094 (9)
N4	0.0321 (12)	0.0386 (12)	0.0435 (13)	-0.0024 (10)	0.0001 (10)	0.0054 (10)
C1	0.0268 (12)	0.0324 (13)	0.0395 (13)	-0.0056 (10)	0.0006 (10)	0.0001 (10)
C2	0.0307 (13)	0.0357 (14)	0.0519 (15)	0.0003 (11)	-0.0005 (11)	-0.0032 (11)
C3	0.0395 (14)	0.0424 (15)	0.0494 (15)	-0.0006 (12)	0.0058 (11)	-0.0098 (12)
C4	0.0479 (15)	0.0436 (15)	0.0415 (14)	-0.0049 (13)	0.0014 (11)	-0.0057 (12)
C5	0.0403 (14)	0.0372 (14)	0.0401 (14)	-0.0034 (11)	-0.0028 (11)	-0.0009 (11)
C6	0.0306 (12)	0.0249 (12)	0.0397 (13)	-0.0049 (10)	0.0021 (10)	-0.0008 (10)
C7	0.0241 (12)	0.0301 (13)	0.0379 (13)	-0.0050 (10)	-0.0010 (9)	0.0018 (10)
C8	0.0302 (12)	0.0301 (13)	0.0360 (13)	-0.0063 (10)	0.0022 (10)	0.0023 (10)
C9	0.0403 (14)	0.0437 (15)	0.0388 (14)	0.0016 (11)	-0.0015 (11)	0.0033 (11)
C10	0.0545 (17)	0.0534 (17)	0.0344 (14)	0.0004 (13)	-0.0033 (11)	-0.0003 (12)
C11	0.0562 (16)	0.0419 (15)	0.0379 (14)	-0.0036 (13)	0.0077 (12)	-0.0035 (11)
C12	0.0425 (14)	0.0323 (14)	0.0433 (14)	0.0001 (11)	0.0058 (11)	0.0002 (11)
C13	0.0327 (13)	0.0290 (13)	0.0382 (13)	-0.0061 (10)	0.0016 (10)	0.0028 (10)
C14	0.0314 (13)	0.0280 (12)	0.0432 (13)	0.0016 (10)	-0.0025 (10)	0.0012 (10)
C15	0.0306 (12)	0.0286 (12)	0.0372 (13)	0.0003 (10)	-0.0019 (9)	-0.0004 (10)
C16	0.0302 (12)	0.0287 (13)	0.0421 (13)	0.0040 (10)	-0.0001 (10)	-0.0001 (10)
C17	0.0317 (13)	0.0411 (14)	0.0539 (15)	-0.0051 (11)	-0.0012 (11)	0.0024 (12)
C18	0.0367 (14)	0.0506 (16)	0.0554 (16)	-0.0008 (12)	-0.0142 (12)	-0.0053 (13)
C19	0.0464 (15)	0.0485 (16)	0.0451 (14)	0.0009 (13)	-0.0114 (12)	0.0031 (12)

C20	0.0363 (13)	0.0398 (14)	0.0460 (14)	-0.0013 (11)	-0.0025 (11)	0.0054 (11)
C21	0.0489 (17)	0.055 (2)	0.100 (2)	-0.0015 (14)	0.0074 (15)	0.0046 (16)

Geometric parameters (Å, °)

O1—C21	1.395 (3)	C8—C9	1.402 (3)
O1—H1	0.95 (3)	C9—C10	1.376 (3)
N1—C7	1.320 (3)	C9—H9	0.9500
N1—C1	1.386 (3)	C10—C11	1.389 (3)
N2—C7	1.384 (3)	C10—H10	0.9500
N2—C6	1.387 (3)	C11—C12	1.373 (3)
N2—C14	1.475 (3)	C11—H11	0.9500
N3—C13	1.386 (3)	C12—C13	1.398 (3)
N3—C14	1.459 (3)	C12—H12	0.9500
N3—H3a	0.93 (2)	C14—C15	1.509 (3)
N4—C16	1.406 (3)	C14—H14	1.0000
N4—H4a	0.95 (2)	C15—C20	1.391 (3)
N4—H4b	0.86 (2)	C15—C16	1.400 (3)
C1—C2	1.397 (3)	C16—C17	1.393 (3)
C1—C6	1.402 (3)	C17—C18	1.372 (3)
C2—C3	1.379 (3)	C17—H17	0.9500
C2—H2	0.9500	C18—C19	1.377 (3)
C3—C4	1.399 (3)	C18—H18	0.9500
C3—H3	0.9500	C19—C20	1.386 (3)
C4—C5	1.379 (3)	C19—H19	0.9500
C4—H4	0.9500	C20—H20	0.9500
C5—C6	1.390 (3)	C21—H21A	0.9800
C5—H5	0.9500	C21—H21B	0.9800
C7—C8	1.441 (3)	C21—H21C	0.9800
C8—C13	1.396 (3)		
C21—O1—H1	109.6 (18)	C11—C10—H10	120.6
C7—N1—C1	105.19 (17)	C12—C11—C10	121.5 (2)
C7—N2—C6	106.81 (17)	C12—C11—H11	119.3
C7—N2—C14	125.65 (17)	C10—C11—H11	119.3
C6—N2—C14	126.49 (17)	C11—C12—C13	120.0 (2)
C13—N3—C14	124.33 (19)	C11—C12—H12	120.0
C13—N3—H3a	116.1 (13)	C13—C12—H12	120.0
C14—N3—H3a	109.7 (13)	N3—C13—C8	120.48 (19)
C16—N4—H4a	112.2 (13)	N3—C13—C12	120.1 (2)
C16—N4—H4b	111.0 (15)	C8—C13—C12	119.29 (19)
H4a—N4—H4b	111 (2)	N3—C14—N2	108.15 (16)
N1—C1—C2	129.15 (19)	N3—C14—C15	110.02 (17)
N1—C1—C6	110.53 (18)	N2—C14—C15	112.88 (17)
C2—C1—C6	120.3 (2)	N3—C14—H14	108.6
C3—C2—C1	117.5 (2)	N2—C14—H14	108.6
C3—C2—H2	121.3	C15—C14—H14	108.6
C1—C2—H2	121.3	C20—C15—C16	119.16 (19)

C2—C3—C4	121.8 (2)	C20—C15—C14	118.47 (19)
C2—C3—H3	119.1	C16—C15—C14	122.33 (19)
C4—C3—H3	119.1	C17—C16—C15	118.5 (2)
C5—C4—C3	121.4 (2)	C17—C16—N4	119.7 (2)
C5—C4—H4	119.3	C15—C16—N4	121.82 (19)
C3—C4—H4	119.3	C18—C17—C16	121.3 (2)
C4—C5—C6	117.1 (2)	C18—C17—H17	119.3
C4—C5—H5	121.4	C16—C17—H17	119.3
C6—C5—H5	121.4	C17—C18—C19	120.7 (2)
N2—C6—C5	133.1 (2)	C17—C18—H18	119.7
N2—C6—C1	104.99 (17)	C19—C18—H18	119.7
C5—C6—C1	121.9 (2)	C18—C19—C20	118.7 (2)
N1—C7—N2	112.39 (18)	C18—C19—H19	120.7
N1—C7—C8	128.31 (19)	C20—C19—H19	120.7
N2—C7—C8	119.27 (19)	C19—C20—C15	121.5 (2)
C13—C8—C9	119.5 (2)	C19—C20—H20	119.2
C13—C8—C7	117.73 (18)	C15—C20—H20	119.2
C9—C8—C7	122.7 (2)	O1—C21—H21A	109.5
C10—C9—C8	120.9 (2)	O1—C21—H21B	109.5
C10—C9—H9	119.6	H21A—C21—H21B	109.5
C8—C9—H9	119.6	O1—C21—H21C	109.5
C9—C10—C11	118.9 (2)	H21A—C21—H21C	109.5
C9—C10—H10	120.6	H21B—C21—H21C	109.5
C7—N1—C1—C2	179.8 (2)	C10—C11—C12—C13	1.1 (3)
C7—N1—C1—C6	-0.4 (2)	C14—N3—C13—C8	21.6 (3)
N1—C1—C2—C3	177.9 (2)	C14—N3—C13—C12	-162.73 (19)
C6—C1—C2—C3	-1.9 (3)	C9—C8—C13—N3	175.3 (2)
C1—C2—C3—C4	0.5 (3)	C7—C8—C13—N3	-7.0 (3)
C2—C3—C4—C5	0.6 (4)	C9—C8—C13—C12	-0.4 (3)
C3—C4—C5—C6	-0.4 (3)	C7—C8—C13—C12	177.30 (18)
C7—N2—C6—C5	176.7 (2)	C11—C12—C13—N3	-176.6 (2)
C14—N2—C6—C5	8.0 (4)	C11—C12—C13—C8	-0.8 (3)
C7—N2—C6—C1	-3.1 (2)	C13—N3—C14—N2	-24.8 (3)
C14—N2—C6—C1	-171.81 (18)	C13—N3—C14—C15	-148.5 (2)
C4—C5—C6—N2	179.3 (2)	C7—N2—C14—N3	17.3 (3)
C4—C5—C6—C1	-1.0 (3)	C6—N2—C14—N3	-176.03 (18)
N1—C1—C6—N2	2.2 (2)	C7—N2—C14—C15	139.21 (19)
C2—C1—C6—N2	-178.01 (19)	C6—N2—C14—C15	-54.1 (3)
N1—C1—C6—C5	-177.61 (18)	N3—C14—C15—C20	-118.8 (2)
C2—C1—C6—C5	2.2 (3)	N2—C14—C15—C20	120.3 (2)
C1—N1—C7—N2	-1.7 (2)	N3—C14—C15—C16	59.1 (3)
C1—N1—C7—C8	176.36 (19)	N2—C14—C15—C16	-61.8 (3)
C6—N2—C7—N1	3.1 (2)	C20—C15—C16—C17	-3.7 (3)
C14—N2—C7—N1	171.95 (18)	C14—C15—C16—C17	178.5 (2)
C6—N2—C7—C8	-175.15 (17)	C20—C15—C16—N4	178.1 (2)
C14—N2—C7—C8	-6.3 (3)	C14—C15—C16—N4	0.3 (3)
N1—C7—C8—C13	-178.1 (2)	C15—C16—C17—C18	2.6 (3)

N2—C7—C8—C13	-0.2 (3)	N4—C16—C17—C18	-179.1 (2)
N1—C7—C8—C9	-0.5 (3)	C16—C17—C18—C19	0.3 (4)
N2—C7—C8—C9	177.44 (19)	C17—C18—C19—C20	-2.1 (4)
C13—C8—C9—C10	1.4 (3)	C18—C19—C20—C15	0.9 (4)
C7—C8—C9—C10	-176.1 (2)	C16—C15—C20—C19	2.0 (3)
C8—C9—C10—C11	-1.2 (3)	C14—C15—C20—C19	179.9 (2)
C9—C10—C11—C12	0.0 (4)		

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N3—H3 <i>a</i> ...N4 ⁱ	0.93 (2)	2.13 (2)	3.058 (3)	174.5 (18)
N4—H4 <i>a</i> ...O1 ⁱⁱ	0.95 (2)	2.00 (3)	2.946 (3)	174.4 (19)
N4—H4 <i>b</i> ...N3	0.86 (2)	2.35 (2)	3.006 (3)	133.8 (19)
O1—H1...N1	0.95 (3)	1.88 (3)	2.814 (2)	167 (3)
C14—H14...C <i>g</i> ⁱ	1.00	2.44	3.408 (2)	162

Symmetry codes: (i) *x, y-1, z*; (ii) *x+1, y, z*.