

## Hydroxonium creatininium bis(pyridine-2,6-dicarboxylato- $\kappa^3 O^2, N, O^6$ )nickel-ate(II) trihydrate

Jafar Attar Gharamaleki,<sup>a</sup> Hossein Aghabozorg,<sup>b\*</sup> Zohreh Derikvand<sup>c</sup> and Mohammad Yousefi<sup>d</sup>

<sup>a</sup>Young Researchers Club, Islamic Azad University, North Tehran Branch, Tehran, Iran,

<sup>b</sup>Faculty of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran,

<sup>c</sup>Department of Chemistry, Faculty of Science, Islamic Azad University, Khorramabad Branch, Khorramabad, Iran, and

<sup>d</sup>Department of Chemistry, Islamic Azad University, Shahr-e Rey Branch, Tehran, Iran

Correspondence e-mail: haghabozorg@yahoo.com


Received 19 May 2009; accepted 10 June 2009

Key indicators: single-crystal X-ray study;  $T = 120\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.053;  $wR$  factor = 0.134; data-to-parameter ratio = 16.1.

The title compound,  $(\text{C}_4\text{H}_8\text{N}_3\text{O})(\text{H}_3\text{O})[\text{Ni}(\text{C}_7\text{H}_3\text{NO}_4)_2] \cdot 3\text{H}_2\text{O}$ , exhibits isotypism with its  $\text{Co}^{II}$  analogue. All intramolecular distances and angles are similar for the two structures. This applies also for the intermolecular forces, consisting of  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds and  $\pi-\pi$  interactions [with centroid–centroid distances of 3.428 (2) and 3.579 (2)  $\text{\AA}$ ], that lead to a cohesion of the structure.

### Related literature

For background to proton-transfer agents, see: Aghabozorg, Manteghi *et al.* (2008); Soleimannejad *et al.* (2005); Aghabozorg, Ramezanipour *et al.* (2008). For related structures, see: Moghimi *et al.* (2004, 2005); Aghabozorg, Motyeian *et al.* (2008); Aghabozorg, Derikvand *et al.* (2008). For the isotopic Co compound, see: Aghabozorg *et al.* (2009).



### Experimental

#### Crystal data

$(\text{C}_4\text{H}_8\text{N}_3\text{O})(\text{H}_3\text{O})[\text{Ni}(\text{C}_7\text{H}_3\text{NO}_4)_2] \cdot 3\text{H}_2\text{O}$	$\beta = 89.944 (2)^\circ$
$M_r = 576.12$	$\gamma = 87.564 (3)^\circ$
Triclinic, $P\bar{1}$	$V = 1140.0 (2) \text{ \AA}^3$
$a = 8.1466 (9) \text{ \AA}$	$Z = 2$
$b = 10.7420 (12) \text{ \AA}$	Mo $K\alpha$ radiation
$c = 13.5061 (15) \text{ \AA}$	$\mu = 0.93 \text{ mm}^{-1}$
$\alpha = 74.890 (2)^\circ$	$T = 120 \text{ K}$
	$0.18 \times 0.14 \times 0.12 \text{ mm}$

#### Data collection

Bruker SMART 1000 CCD area-detector diffractometer	11509 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 1998)	5404 independent reflections
$T_{\min} = 0.803$ , $T_{\max} = 0.896$	3827 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.046$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$	335 parameters
$wR(F^2) = 0.134$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.84 \text{ e \AA}^{-3}$
5404 reflections	$\Delta\rho_{\min} = -0.37 \text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3N $\cdots$ O1W	0.85	1.89	2.726 (4)	166
N5—H5NB $\cdots$ O3 <sup>i</sup>	0.93	1.96	2.861 (4)	162
N5—H5NA $\cdots$ O8 <sup>ii</sup>	0.83	1.97	2.777 (4)	165
O1W—H1WA $\cdots$ O3W	0.85	1.84	2.662 (3)	162
O1W—H1WB $\cdots$ O7 <sup>ii</sup>	0.85	1.97	2.804 (3)	166
O2W—H2WA $\cdots$ O6	0.85	1.71	2.529 (3)	161
O2W—H2WB $\cdots$ O4 <sup>iii</sup>	0.85	1.65	2.480 (3)	165
O2W—H2WC $\cdots$ O4W	0.85	1.73	2.521 (3)	155
O3W—H3WA $\cdots$ O9	0.85	2.17	2.926 (4)	148
O3W—H3WB $\cdots$ O2 <sup>iv</sup>	0.85	1.94	2.765 (4)	162
O4W—H4WA $\cdots$ O2 <sup>iv</sup>	0.85	1.85	2.682 (4)	165
O4W—H4WB $\cdots$ O5 <sup>v</sup>	0.85	1.91	2.722 (4)	159

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $x + 1, y + 1, z$ ; (iii)  $x + 1, y, z$ ; (iv)  $x, y + 1, z$ ; (v)  $-x + 1, -y + 1, -z$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT-Plus* (Bruker, 1998); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2161).

### References

- Aghabozorg, H., Derikvand, Z., Attar Gharamaleki, J. & Yousefi, M. (2009). *Acta Cryst. E65*, m826–m827.
- Aghabozorg, H., Derikvand, Z., Olmstead, M. M. & Attar Gharamaleki, J. (2008). *Acta Cryst. E64*, m1234–m1235.
- Aghabozorg, H., Manteghi, F. & Sheshmani, S. (2008). *J. Iran. Chem. Soc.* **5**, 184–227.
- Aghabozorg, H., Motyeian, E., Attar Gharamaleki, J., Soleimannejad, J., Ghadermazi, M. & Spey Sharon, E. (2008). *Acta Cryst. E64*, m144–m145.
- Aghabozorg, H., Ramezanipour, F., Soleimannejad, J., Sharif, M. A., Shokrollahi, A., Shamsipur, M., Moghimi, A., Attar Gharamaleki, J., Lippolis, V. & Blake, A. J. (2008). *Pol. J. Chem.* **82**, 487–507.

- Bruker (1998). *SMART*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Moghimi, A., Sharif, M. A. & Aghabozorg, H. (2004). *Acta Cryst. E* **60**, o1790–o1792.
- Moghimi, A., Sharif, M. A., Shokrollahi, A., Shamsipur, M. & Aghabozorg, H. (2005). *Z. Anorg. Allg. Chem.* **631**, 902–908.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Soleimannejad, J., Sharif, M. A., Sheshmani, S., Alizadeh, R., Moghimi, A. & Aghabozorg, H. (2005). *Anal. Sci.* **21**, x49–x50.

# supporting information

*Acta Cryst.* (2009). E65, m824–m825 [doi:10.1107/S1600536809022053]

## Hydroxonium creatininium bis(pyridine-2,6-dicarboxylato- $\kappa^3O^2,N,O^6$ )nickelate(II) trihydrate

Jafar Attar Gharamaleki, Hossein Aghabozorg, Zohreh Derikvand and Mohammad Yousefi

### S1. Comment

Two proton transfer compounds have been prepared from creatinine as proton acceptor agent by using pyridine-2,6-dicarboxylic acid (pydcH<sub>2</sub>) (Moghimi *et al.*, 2004) and 1,10-phenanthroline-9,13-dicarboxylic acid (phendcH<sub>2</sub>) (Soleimannejad *et al.*, 2005) as proton donor agents. Also three complexes of Bi<sup>III</sup> (Aghabozorg, Ramezanipour *et al.*, 2008), Zn<sup>II</sup> (Moghimi *et al.*, 2005) and Cr<sup>III</sup> (Aghabozorg, Derikvand *et al.*, 2008) have been reported by our research group in which the (creatH)<sup>+</sup> fragment was a part of the crystal structure; more details can be found in our recently published review article (Aghabozorg, Manteghi *et al.*, 2008). We have now synthesized a novel Ni<sup>II</sup> complex, (I), and determined its crystal structure which is presented in this article.

The crystal structure of the title complex, (I), (Fig. 1) contains [Ni(pydc)<sub>2</sub>]<sup>2-</sup>, creatininium (creatH)<sup>+</sup> as counter-ion, hydroxonium cation, (H<sub>3</sub>O)<sup>+</sup> and three uncoordinated water molecules. The Ni<sup>II</sup> atom is coordinated by two tridentate (pydc)<sup>2-</sup> groups and the presence of (creatH)<sup>+</sup> and (H<sub>3</sub>O)<sup>+</sup> ions balance the negative charge. In the structure of (I), the mean Ni—N and Ni—O bond distances are 1.973 (3) and 2.140 (2) Å, respectively, which are consistent with the corresponding distances reported for similar Ni<sup>II</sup> complexes (Aghabozorg, Motyeian *et al.*, 2008). The N-atoms (N1 and N2) of the two (pydc)<sup>2-</sup> fragments occupy the axial positions, while O1, O3, O5 and O7 atoms form the equatorial plane. In the anionic complex, the N1—Ni1—N2 angle [174.74 (11)°] deviates slightly from linearity. Therefore, the coordination around the Ni<sup>II</sup> atom is distorted octahedral. In addition, the O1—Ni1—O3 and O5—Ni1—O7 angles [155.14 (9) and 155.22 (9)°] indicate that the four carboxylate groups of the two dianions are oriented in a flattened tetrahedral arrangement around the Ni<sup>II</sup> atom. The angles O1—Ni1—O5, O1—Ni1—O7, O3—Ni1—O5 and O3—Ni1—O7 are 96.01 (9), 89.16 (9), 89.63 (9) and 95.81 (9)°, respectively. On the other hand, the torsion angles O1—Ni1—O7—C14 and O5—Ni1—O3—C7 are 93.3 (2) and 93.6 (2)°, respectively, indicating that two (pydc)<sup>2-</sup> units are almost perpendicular to each other. The angle between two planes passing aromatic rings of (pydc)<sup>2-</sup> units is 81.13 (15)°.

A remarkable feature of the title compound is the presence of a large number of O—H···O, N—H···O and C—H···O hydrogen bonds, with D···A distances ranging from 2.480 (3) to 3.415 (4) Å between (creatH)<sup>+</sup>, (H<sub>3</sub>O)<sup>+</sup> and [Ni(pydc)<sub>2</sub>]<sup>2-</sup> fragments and uncoordinated water molecules. The O2W—O2WB···O4<sup>iii</sup> hydrogen bond with D···A distance of 2.480 (3) Å, is the strongest (details of the hydrogen bonding geometry have been provided in Table 1). These interactions link the fragments to form a three-dimensional network as shown in Fig. 2. In the structure of (I), the space between two layers of [Ni(pydc)<sub>2</sub>]<sup>2-</sup> anions is filled with layers of (creatH)<sup>+</sup>, (H<sub>3</sub>O)<sup>+</sup> cations and uncoordinated water molecules (Fig. 2).


There is noticeable  $\pi$ – $\pi$  stacking interaction between two aromatic rings of (pydc)<sup>2-</sup> units, with distances 3.428 (2) Å (1 -  $x$ , - $y$ , - $z$  + 1) and 3.579 (2) Å (- $x$ , - $y$ , - $z$ ). Also a considerable centrosymmetric C=O··· $\pi$  stacking interactions between C=O groups of carboxylate fragments with aromatic rings of pyridine-2,6-dicarboxylate with distance of 3.488 (1) Å for C=O2···Cg1 (1 -  $x$ , - $y$ , - $z$ ) [Cg1 is the centroid for N1/C1—C5 ring] are observed in (I) (Fig. 3).

**S2. Experimental**


The reaction of nickel(II) chloride hexahydrate (59 mg, 0.25 mmol), creatinine, creat, (57 mg, 0.5 mmol) and pyridine-2,6-dicarboxylic acid, pydCH<sub>2</sub>, (83 mg, 0.5 mmol) in a 1:2:2 molar ratio in aqueous solution resulted in the formation of green, (H<sub>3</sub>O)(creatH)[Ni(pydc)<sub>2</sub>]. 3H<sub>2</sub>O crystals.

**S3. Refinement**


The H atoms of water molecules, NH and NH<sub>2</sub> groups were located in difference Fourier synthesis. All hydrogen atoms were included in the refinement in isotropic approximation in riding model with distances: N—H (at positions from difference map), C—H = 0.95 (aryl), 0.98 (methyl), 0.99 (methylene) and O—H = 0.85 Å, and the  $U_{\text{iso}}(\text{H})$  parameters equal to 1.2  $U_{\text{eq}}(\text{C})$ , 1.5  $U_{\text{eq}}(\text{O},\text{N})$ , where  $U(\text{C},\text{O},\text{N})$  are respectively the equivalent thermal parameters of the C, O and N atoms to which corresponding H atoms were bonded.

**Figure 1**

The molecular structure of the title compound, thermal ellipsoids are shown at 50% probability level.

**Figure 2**

Unit cell packing diagram of the title compound, hydrogen bonds are shown as dashed lines and H-atoms not involved in H-bonds have been excluded.

**Figure 3**

$\pi-\pi$  Stacking interaction between two aromatic rings of  $(\text{pydc})^{2-}$  units.

### Hydroxonium creatininium bis(pyridine-2,6-dicarboxylato)nickelate(II) trihydrate

#### Crystal data



$M_r = 576.12$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 8.1466 (9)$  Å

$b = 10.7420 (12)$  Å

$c = 13.5061 (15)$  Å

$\alpha = 74.890 (2)^\circ$

$\beta = 89.944 (2)^\circ$

$\gamma = 87.564 (3)^\circ$

$V = 1140.0 (2)$  Å<sup>3</sup>

$Z = 2$

$F(000) = 596$

$D_x = 1.678 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1838 reflections

$\theta = 2.8\text{--}24.2^\circ$

$\mu = 0.93 \text{ mm}^{-1}$

$T = 120$  K

Prism, green

$0.18 \times 0.14 \times 0.12$  mm

*Data collection*

Bruker SMART 1000 CCD area-detector diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\omega$  scans  
 Absorption correction: multi-scan (*SADABS*; Bruker, 1998)  
 $T_{\min} = 0.803$ ,  $T_{\max} = 0.896$

11509 measured reflections  
 5404 independent reflections  
 3827 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$   
 $\theta_{\max} = 28.0^\circ$ ,  $\theta_{\min} = 2.0^\circ$   
 $h = -10 \rightarrow 10$   
 $k = -14 \rightarrow 14$   
 $l = -17 \rightarrow 17$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.053$   
 $wR(F^2) = 0.134$   
 $S = 1.00$   
 5404 reflections  
 335 parameters  
 0 restraints  
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map  
 Hydrogen site location: mixed  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 2.37P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.26883 (5)	0.07342 (4)	0.22939 (3)	0.01905 (13)
O1	0.4241 (3)	-0.0881 (2)	0.23014 (17)	0.0225 (5)
O2	0.4819 (3)	-0.2302 (2)	0.13848 (18)	0.0245 (5)
O3	0.0829 (3)	0.2229 (2)	0.16633 (17)	0.0211 (5)
O4	-0.0572 (3)	0.3100 (2)	0.01983 (17)	0.0254 (5)
O5	0.4573 (3)	0.2152 (2)	0.18788 (17)	0.0220 (5)
O6	0.6101 (3)	0.3404 (2)	0.25748 (18)	0.0271 (5)
O7	0.1100 (3)	-0.0493 (2)	0.33115 (17)	0.0228 (5)
O8	0.0881 (3)	-0.1386 (2)	0.50037 (18)	0.0271 (5)
N1	0.2282 (3)	0.0494 (2)	0.0918 (2)	0.0165 (5)
N2	0.3261 (3)	0.1077 (2)	0.3616 (2)	0.0161 (5)
C1	0.3098 (4)	-0.0458 (3)	0.0640 (2)	0.0175 (6)
C2	0.2950 (4)	-0.0601 (3)	-0.0352 (2)	0.0201 (7)
H2A	0.3515	-0.1291	-0.0547	0.024*
C3	0.1952 (4)	0.0295 (3)	-0.1048 (3)	0.0221 (7)
H3A	0.1842	0.0227	-0.1732	0.027*

C4	0.1118 (4)	0.1285 (3)	-0.0749 (2)	0.0208 (7)
H4A	0.0440	0.1904	-0.1221	0.025*
C5	0.1297 (4)	0.1347 (3)	0.0252 (2)	0.0174 (6)
C6	0.4140 (4)	-0.1286 (3)	0.1508 (2)	0.0189 (7)
C7	0.0456 (4)	0.2315 (3)	0.0738 (2)	0.0203 (7)
C8	0.4375 (4)	0.1947 (3)	0.3651 (2)	0.0175 (6)
C9	0.4809 (4)	0.2192 (3)	0.4569 (2)	0.0207 (7)
H9A	0.5607	0.2803	0.4591	0.025*
C10	0.4039 (4)	0.1512 (3)	0.5470 (2)	0.0212 (7)
H10A	0.4291	0.1671	0.6112	0.025*
C11	0.2910 (4)	0.0610 (3)	0.5404 (2)	0.0220 (7)
H11A	0.2383	0.0134	0.6003	0.026*
C12	0.2554 (4)	0.0409 (3)	0.4455 (2)	0.0168 (6)
C13	0.5092 (4)	0.2559 (3)	0.2620 (2)	0.0202 (7)
C14	0.1407 (4)	-0.0569 (3)	0.4255 (2)	0.0193 (7)
O9	0.7005 (3)	0.4732 (2)	0.46109 (18)	0.0300 (6)
N3	0.8635 (3)	0.5880 (3)	0.5417 (2)	0.0186 (6)
H3N	0.9098	0.6328	0.4886	0.028*
N4	0.7837 (3)	0.5199 (3)	0.7029 (2)	0.0202 (6)
N5	0.9931 (3)	0.6668 (3)	0.6669 (2)	0.0213 (6)
H5NB	0.9862	0.6905	0.7286	0.032*
H5NA	1.0388	0.7217	0.6217	0.032*
C15	0.7458 (4)	0.5019 (3)	0.5378 (3)	0.0217 (7)
C16	0.8858 (4)	0.5953 (3)	0.6403 (2)	0.0185 (6)
C17	0.6828 (4)	0.4533 (3)	0.6458 (3)	0.0226 (7)
H17A	0.5647	0.4771	0.6499	0.027*
H17B	0.6996	0.3585	0.6710	0.027*
C18	0.7939 (5)	0.4855 (4)	0.8144 (3)	0.0288 (8)
H18A	0.8137	0.5627	0.8376	0.043*
H18B	0.8844	0.4214	0.8376	0.043*
H18C	0.6904	0.4491	0.8429	0.043*
O1W	0.9969 (3)	0.7014 (2)	0.35667 (18)	0.0257 (5)
H1WA	0.9086	0.6978	0.3243	0.038*
H1WB	1.0206	0.7804	0.3398	0.038*
O2W	0.7807 (3)	0.4512 (2)	0.10606 (19)	0.0310 (6)
H2WA	0.7267	0.3996	0.1517	0.046*
H2WB	0.8208	0.4006	0.0723	0.046*
H2WC	0.7360	0.5102	0.0580	0.046*
O3W	0.7459 (3)	0.6405 (3)	0.2550 (2)	0.0385 (7)
H3WA	0.7188	0.5734	0.3000	0.058*
H3WB	0.6564	0.6811	0.2316	0.058*
O4W	0.6023 (4)	0.6403 (3)	0.0074 (2)	0.0454 (8)
H4WA	0.5592	0.6919	0.0393	0.068*
H4WB	0.5964	0.6694	-0.0574	0.068*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ni1	0.0227 (2)	0.0187 (2)	0.0156 (2)	-0.00178 (16)	-0.00045 (16)	-0.00419 (16)
O1	0.0295 (13)	0.0209 (12)	0.0176 (11)	0.0012 (10)	0.0011 (10)	-0.0064 (9)
O2	0.0315 (13)	0.0208 (12)	0.0220 (12)	0.0086 (10)	-0.0026 (10)	-0.0087 (10)
O3	0.0254 (12)	0.0215 (12)	0.0168 (11)	0.0008 (10)	0.0001 (9)	-0.0062 (9)
O4	0.0287 (13)	0.0262 (13)	0.0193 (12)	0.0074 (10)	-0.0034 (10)	-0.0040 (10)
O5	0.0249 (12)	0.0263 (12)	0.0155 (11)	-0.0045 (10)	0.0021 (9)	-0.0060 (9)
O6	0.0306 (14)	0.0279 (13)	0.0220 (12)	-0.0110 (11)	0.0015 (10)	-0.0038 (10)
O7	0.0294 (13)	0.0204 (12)	0.0186 (12)	-0.0068 (10)	0.0023 (10)	-0.0042 (9)
O8	0.0349 (14)	0.0231 (12)	0.0222 (12)	-0.0078 (11)	0.0060 (10)	-0.0027 (10)
N1	0.0151 (13)	0.0140 (12)	0.0186 (13)	-0.0019 (10)	0.0003 (10)	-0.0009 (10)
N2	0.0170 (13)	0.0132 (12)	0.0167 (13)	0.0031 (10)	0.0000 (10)	-0.0023 (10)
C1	0.0153 (15)	0.0184 (15)	0.0193 (16)	-0.0025 (12)	0.0014 (12)	-0.0058 (12)
C2	0.0194 (16)	0.0195 (16)	0.0226 (17)	-0.0040 (13)	0.0014 (13)	-0.0071 (13)
C3	0.0268 (18)	0.0234 (17)	0.0170 (16)	-0.0025 (14)	-0.0039 (13)	-0.0065 (13)
C4	0.0216 (17)	0.0204 (16)	0.0178 (16)	-0.0010 (13)	-0.0030 (13)	-0.0002 (12)
C5	0.0180 (16)	0.0158 (15)	0.0178 (15)	-0.0031 (12)	0.0026 (12)	-0.0031 (12)
C6	0.0182 (16)	0.0195 (16)	0.0187 (16)	-0.0032 (13)	0.0012 (12)	-0.0039 (13)
C7	0.0229 (17)	0.0175 (15)	0.0188 (16)	0.0005 (13)	-0.0011 (13)	-0.0019 (12)
C8	0.0146 (15)	0.0129 (14)	0.0233 (16)	0.0017 (12)	0.0007 (12)	-0.0021 (12)
C9	0.0204 (16)	0.0193 (16)	0.0216 (16)	0.0027 (13)	-0.0028 (13)	-0.0046 (13)
C10	0.0263 (18)	0.0221 (16)	0.0157 (15)	0.0054 (14)	-0.0040 (13)	-0.0066 (13)
C11	0.0242 (17)	0.0226 (17)	0.0164 (16)	0.0071 (14)	0.0006 (13)	-0.0012 (13)
C12	0.0148 (15)	0.0151 (15)	0.0193 (16)	0.0040 (12)	0.0008 (12)	-0.0032 (12)
C13	0.0204 (16)	0.0196 (16)	0.0195 (16)	0.0003 (13)	0.0004 (13)	-0.0030 (13)
C14	0.0180 (16)	0.0168 (15)	0.0232 (17)	0.0015 (12)	0.0016 (13)	-0.0055 (13)
O9	0.0341 (14)	0.0351 (14)	0.0227 (13)	-0.0115 (12)	-0.0032 (11)	-0.0094 (11)
N3	0.0211 (14)	0.0209 (14)	0.0132 (13)	-0.0024 (11)	0.0022 (10)	-0.0035 (10)
N4	0.0246 (15)	0.0185 (13)	0.0165 (13)	-0.0040 (11)	0.0016 (11)	-0.0021 (11)
N5	0.0267 (15)	0.0225 (14)	0.0140 (13)	-0.0018 (12)	-0.0001 (11)	-0.0034 (11)
C15	0.0211 (16)	0.0188 (16)	0.0241 (17)	-0.0007 (13)	-0.0012 (13)	-0.0037 (13)
C16	0.0199 (16)	0.0165 (15)	0.0179 (16)	0.0050 (12)	0.0009 (12)	-0.0032 (12)
C17	0.0220 (17)	0.0231 (17)	0.0220 (17)	-0.0050 (14)	-0.0002 (13)	-0.0043 (13)
C18	0.040 (2)	0.0268 (18)	0.0181 (17)	-0.0089 (16)	0.0017 (15)	-0.0015 (14)
O1W	0.0261 (13)	0.0248 (13)	0.0248 (13)	-0.0066 (10)	-0.0014 (10)	-0.0034 (10)
O2W	0.0373 (15)	0.0255 (13)	0.0266 (13)	0.0023 (11)	0.0072 (11)	-0.0013 (11)
O3W	0.0363 (16)	0.0365 (15)	0.0345 (15)	-0.0107 (12)	-0.0101 (12)	0.0067 (12)
O4W	0.082 (2)	0.0346 (15)	0.0169 (13)	0.0270 (15)	-0.0012 (13)	-0.0065 (11)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Ni1—N1	1.972 (3)	C10—C11	1.384 (5)
Ni1—N2	1.974 (3)	C10—H10A	0.9500
Ni1—O1	2.101 (2)	C11—C12	1.387 (4)
Ni1—O7	2.121 (2)	C11—H11A	0.9500
Ni1—O3	2.163 (2)	C12—C14	1.511 (4)

Ni1—O5	2.178 (2)	O9—C15	1.218 (4)
O1—C6	1.262 (4)	N3—C16	1.368 (4)
O2—C6	1.252 (4)	N3—C15	1.370 (4)
O3—C7	1.265 (4)	N3—H3N	0.8511
O4—C7	1.251 (4)	N4—C16	1.327 (4)
O5—C13	1.272 (4)	N4—C18	1.455 (4)
O6—C13	1.240 (4)	N4—C17	1.457 (4)
O7—C14	1.280 (4)	N5—C16	1.298 (4)
O8—C14	1.243 (4)	N5—H5NB	0.9334
N1—C1	1.330 (4)	N5—H5NA	0.8300
N1—C5	1.342 (4)	C15—C17	1.512 (5)
N2—C12	1.320 (4)	C17—H17A	0.9900
N2—C8	1.340 (4)	C17—H17B	0.9900
C1—C2	1.394 (4)	C18—H18A	0.9800
C1—C6	1.509 (4)	C18—H18B	0.9800
C2—C3	1.391 (5)	C18—H18C	0.9800
C2—H2A	0.9500	O1W—H1WA	0.8500
C3—C4	1.385 (5)	O1W—H1WB	0.8500
C3—H3A	0.9500	O2W—H2WA	0.8499
C4—C5	1.379 (4)	O2W—H2WB	0.8500
C4—H4A	0.9500	O2W—H2WC	0.8500
C5—C7	1.509 (4)	O3W—H3WA	0.8500
C8—C9	1.382 (5)	O3W—H3WB	0.8500
C8—C13	1.505 (4)	O4W—H4WA	0.8500
C9—C10	1.408 (5)	O4W—H4WB	0.8500
C9—H9A	0.9500		
N1—Ni1—N2	174.74 (11)	C8—C9—H9A	120.8
N1—Ni1—O1	78.05 (10)	C10—C9—H9A	120.8
N2—Ni1—O1	101.80 (10)	C11—C10—C9	118.8 (3)
N1—Ni1—O7	106.74 (10)	C11—C10—H10A	120.6
N2—Ni1—O7	78.49 (10)	C9—C10—H10A	120.6
O1—Ni1—O7	89.16 (9)	C10—C11—C12	119.3 (3)
N1—Ni1—O3	77.19 (9)	C10—C11—H11A	120.3
N2—Ni1—O3	103.06 (9)	C12—C11—H11A	120.3
O1—Ni1—O3	155.14 (9)	N2—C12—C11	121.0 (3)
O7—Ni1—O3	95.81 (9)	N2—C12—C14	112.9 (3)
N1—Ni1—O5	98.03 (9)	C11—C12—C14	126.1 (3)
N2—Ni1—O5	76.74 (10)	O6—C13—O5	126.9 (3)
O1—Ni1—O5	96.01 (9)	O6—C13—C8	117.6 (3)
O7—Ni1—O5	155.22 (9)	O5—C13—C8	115.5 (3)
O3—Ni1—O5	89.63 (9)	O8—C14—O7	125.9 (3)
C6—O1—Ni1	114.6 (2)	O8—C14—C12	118.2 (3)
C7—O3—Ni1	113.8 (2)	O7—C14—C12	115.9 (3)
C13—O5—Ni1	114.4 (2)	C16—N3—C15	110.3 (3)
C14—O7—Ni1	112.9 (2)	C16—N3—H3N	126.5
C1—N1—C5	121.1 (3)	C15—N3—H3N	123.1
C1—N1—Ni1	118.9 (2)	C16—N4—C18	125.2 (3)

C5—N1—Ni1	119.8 (2)	C16—N4—C17	110.3 (3)
C12—N2—C8	121.4 (3)	C18—N4—C17	123.3 (3)
C12—N2—Ni1	118.5 (2)	C16—N5—H5NB	122.1
C8—N2—Ni1	120.1 (2)	C16—N5—H5NA	119.2
N1—C1—C2	120.9 (3)	H5NB—N5—H5NA	111.9
N1—C1—C6	111.8 (3)	O9—C15—N3	125.8 (3)
C2—C1—C6	127.3 (3)	O9—C15—C17	127.5 (3)
C3—C2—C1	118.1 (3)	N3—C15—C17	106.7 (3)
C3—C2—H2A	120.9	N5—C16—N4	125.9 (3)
C1—C2—H2A	120.9	N5—C16—N3	123.6 (3)
C4—C3—C2	120.3 (3)	N4—C16—N3	110.6 (3)
C4—C3—H3A	119.8	N4—C17—C15	102.1 (3)
C2—C3—H3A	119.8	N4—C17—H17A	111.3
C5—C4—C3	118.3 (3)	C15—C17—H17A	111.3
C5—C4—H4A	120.9	N4—C17—H17B	111.3
C3—C4—H4A	120.9	C15—C17—H17B	111.3
N1—C5—C4	121.3 (3)	H17A—C17—H17B	109.2
N1—C5—C7	112.1 (3)	N4—C18—H18A	109.5
C4—C5—C7	126.6 (3)	N4—C18—H18B	109.5
O2—C6—O1	125.8 (3)	H18A—C18—H18B	109.5
O2—C6—C1	118.1 (3)	N4—C18—H18C	109.5
O1—C6—C1	116.1 (3)	H18A—C18—H18C	109.5
O4—C7—O3	126.4 (3)	H18B—C18—H18C	109.5
O4—C7—C5	117.0 (3)	H1WA—O1W—H1WB	105.5
O3—C7—C5	116.6 (3)	H2WA—O2W—H2WB	101.4
N2—C8—C9	121.1 (3)	H2WA—O2W—H2WC	123.5
N2—C8—C13	113.2 (3)	H2WB—O2W—H2WC	100.7
C9—C8—C13	125.7 (3)	H3WA—O3W—H3WB	106.0
C8—C9—C10	118.4 (3)	H4WA—O4W—H4WB	113.1
N1—Ni1—O1—C6	-3.8 (2)	Ni1—O1—C6—O2	-172.4 (3)
N2—Ni1—O1—C6	-178.4 (2)	Ni1—O1—C6—C1	7.4 (3)
O7—Ni1—O1—C6	103.5 (2)	N1—C1—C6—O2	171.8 (3)
O3—Ni1—O1—C6	1.4 (4)	C2—C1—C6—O2	-8.7 (5)
O5—Ni1—O1—C6	-100.8 (2)	N1—C1—C6—O1	-8.1 (4)
N1—Ni1—O3—C7	-4.7 (2)	C2—C1—C6—O1	171.5 (3)
N2—Ni1—O3—C7	169.9 (2)	Ni1—O3—C7—O4	-179.7 (3)
O1—Ni1—O3—C7	-10.0 (4)	Ni1—O3—C7—C5	2.4 (3)
O7—Ni1—O3—C7	-110.6 (2)	N1—C5—C7—O4	-175.4 (3)
O5—Ni1—O3—C7	93.6 (2)	C4—C5—C7—O4	4.0 (5)
N1—Ni1—O5—C13	178.0 (2)	N1—C5—C7—O3	2.7 (4)
N2—Ni1—O5—C13	-2.5 (2)	C4—C5—C7—O3	-177.9 (3)
O1—Ni1—O5—C13	-103.3 (2)	C12—N2—C8—C9	-1.0 (4)
O7—Ni1—O5—C13	-2.2 (4)	Ni1—N2—C8—C9	-179.2 (2)
O3—Ni1—O5—C13	101.0 (2)	C12—N2—C8—C13	177.3 (3)
N1—Ni1—O7—C14	170.5 (2)	Ni1—N2—C8—C13	-1.0 (3)
N2—Ni1—O7—C14	-8.9 (2)	N2—C8—C9—C10	-0.6 (5)
O1—Ni1—O7—C14	93.3 (2)	C13—C8—C9—C10	-178.7 (3)

O3—Ni1—O7—C14	−111.1 (2)	C8—C9—C10—C11	1.4 (5)
O5—Ni1—O7—C14	−9.3 (4)	C9—C10—C11—C12	−0.7 (5)
O1—Ni1—N1—C1	−1.0 (2)	C8—N2—C12—C11	1.8 (4)
O7—Ni1—N1—C1	−86.4 (2)	Ni1—N2—C12—C11	−179.9 (2)
O3—Ni1—N1—C1	−178.7 (2)	C8—N2—C12—C14	−176.9 (3)
O5—Ni1—N1—C1	93.5 (2)	Ni1—N2—C12—C14	1.4 (3)
O1—Ni1—N1—C5	−175.6 (2)	C10—C11—C12—N2	−0.9 (5)
O7—Ni1—N1—C5	98.9 (2)	C10—C11—C12—C14	177.5 (3)
O3—Ni1—N1—C5	6.6 (2)	Ni1—O5—C13—O6	−176.5 (3)
O5—Ni1—N1—C5	−81.1 (2)	Ni1—O5—C13—C8	2.8 (3)
O1—Ni1—N2—C12	−83.0 (2)	N2—C8—C13—O6	178.0 (3)
O7—Ni1—N2—C12	3.6 (2)	C9—C8—C13—O6	−3.9 (5)
O3—Ni1—N2—C12	97.0 (2)	N2—C8—C13—O5	−1.3 (4)
O5—Ni1—N2—C12	−176.5 (2)	C9—C8—C13—O5	176.8 (3)
O1—Ni1—N2—C8	95.3 (2)	Ni1—O7—C14—O8	−165.9 (3)
O7—Ni1—N2—C8	−178.0 (2)	Ni1—O7—C14—C12	12.2 (3)
O3—Ni1—N2—C8	−84.7 (2)	N2—C12—C14—O8	168.8 (3)
O5—Ni1—N2—C8	1.8 (2)	C11—C12—C14—O8	−9.8 (5)
C5—N1—C1—C2	−0.3 (4)	N2—C12—C14—O7	−9.5 (4)
Ni1—N1—C1—C2	−174.8 (2)	C11—C12—C14—O7	171.9 (3)
C5—N1—C1—C6	179.3 (3)	C16—N3—C15—O9	−178.9 (3)
Ni1—N1—C1—C6	4.8 (3)	C16—N3—C15—C17	2.2 (4)
N1—C1—C2—C3	1.3 (5)	C18—N4—C16—N5	−11.9 (5)
C6—C1—C2—C3	−178.2 (3)	C17—N4—C16—N5	−179.3 (3)
C1—C2—C3—C4	−0.9 (5)	C18—N4—C16—N3	167.9 (3)
C2—C3—C4—C5	−0.5 (5)	C17—N4—C16—N3	0.5 (4)
C1—N1—C5—C4	−1.2 (5)	C15—N3—C16—N5	178.1 (3)
Ni1—N1—C5—C4	173.3 (2)	C15—N3—C16—N4	−1.8 (4)
C1—N1—C5—C7	178.2 (3)	C16—N4—C17—C15	0.8 (3)
Ni1—N1—C5—C7	−7.3 (3)	C18—N4—C17—C15	−166.9 (3)
C3—C4—C5—N1	1.6 (5)	O9—C15—C17—N4	179.4 (3)
C3—C4—C5—C7	−177.8 (3)	N3—C15—C17—N4	−1.8 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N3—H3N···O1W	0.85	1.89	2.726 (4)	166
N5—H5NB···O3 <sup>i</sup>	0.93	1.96	2.861 (4)	162
N5—H5NA···O8 <sup>ii</sup>	0.83	1.97	2.777 (4)	165
O1W—H1WA···O3W	0.85	1.84	2.662 (3)	162
O1W—H1WB···O7 <sup>ii</sup>	0.85	1.97	2.804 (3)	166
O2W—H2WA···O6	0.85	1.71	2.529 (3)	161
O2W—H2WB···O4 <sup>iii</sup>	0.85	1.65	2.480 (3)	165
O2W—H2WC···O4W	0.85	1.73	2.521 (3)	155
O3W—H3WA···O9	0.85	2.17	2.926 (4)	148
O3W—H3WB···O2 <sup>iv</sup>	0.85	1.94	2.765 (4)	162

---

O4W—H4WA···O2 <sup>iv</sup>	0.85	1.85	2.682 (4)	165
O4W—H4WB···O5 <sup>v</sup>	0.85	1.91	2.722 (4)	159

---

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x+1, y+1, z$ ; (iii)  $x+1, y, z$ ; (iv)  $x, y+1, z$ ; (v)  $-x+1, -y+1, -z$ .