

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 2-hydroxy-5-oxo-4-phenyl-2,3,4,5-tetrahydropyrano[3,2-c]chromene-2-carboxylate

 Wei Zhang,^a Guangcun Zhang,^a Bailin Li^{a,b} and Yifeng Wang^{a*}

^aState Key Laboratory Breeding Base of Green Chemistry-Synthesis Technology, Zhejiang University of Technology, Hangzhou 310014, People's Republic of China, and ^bDepartment of Pharmaceutical and Chemical Engineering, Taizhou College, Linhai, Zhejiang 317000, People's Republic of China
Correspondence e-mail: yifengwang108@gmail.com


Received 9 July 2009; accepted 17 July 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.035; wR factor = 0.080; data-to-parameter ratio = 13.9.

The main structural unit of the title compound, $\text{C}_{21}\text{H}_{18}\text{O}_6$, is a fused three-ring group consisting of coumarin and tetrahydropyrane ring systems. Two C atoms of the tetrahydropyrane ring are displaced by 0.295 (3) and -0.360 (2) Å from the mean plane of coumarin ring. The dihedral angle between the phenyl and coumarin rings is 73.94 (3)°. Intermolecular O—H···O hydrogen bonds are present in the crystal structure.

Related literature

For the synthesis of (*E*)-ethyl 2-oxo-4-phenylbut-3-enoate, see: Vaijyanthi & Chadha (2007).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{18}\text{O}_6$
 $M_r = 366.37$
 Monoclinic, $P2_1/n$
 $a = 5.4988$ (2) Å
 $b = 14.9975$ (5) Å
 $c = 21.342$ (1) Å
 $\beta = 98.5487$ (13)°
 $V = 1740.48$ (12) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 $0.41 \times 0.39 \times 0.14$ mm

Data collection

Rigaku R-Axis RAPID diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 2005)
 $T_{\min} = 0.954$, $T_{\max} = 0.986$
 16523 measured reflections
 3422 independent reflections
 2568 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.080$
 $S = 1.00$
 3422 reflections
 246 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.21$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H101···O2 ⁱ	0.82	2.30	2.9198 (15)	132
O1—H101···O5	0.82	2.17	2.6628 (17)	119

 Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *PROCESS-AUTO* (Rigaku, 2006); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku, 2007); program(s) used to solve structure: *SHELXL97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

We are grateful for the help of Professor Jian-Ming Gu of Zhejiang University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2181).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Higashi, T. (2005). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (2006). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (2007). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vaijyanthi, T. & Chadha, A. (2007). *Tetrahedron Asymmetry*, **18**, 1077–1084.

supporting information

Acta Cryst. (2009). E65, o1969 [doi:10.1107/S1600536809028311]

Ethyl 2-hydroxy-5-oxo-4-phenyl-2,3,4,5-tetrahydropyrano[3,2-*c*]chromene-2-carboxylate

Wei Zhang, Guangcun Zhang, Bailin Li and Yifeng Wang

S1. Comment


Coumarin derivatives are widely distributed in nature and are used as versatile intermediates in organic and natural product synthesis. Moreover, this class of compound possess a wide range of biological activities, including anticoagulant and HIV protease inhibition properties. The title compound, which is readily synthesized from commercially available 4-hydroxycoumarin and (*E*)-ethyl 2-oxo-4-phenylbut-3-enoate, can act as an intermediate in organic and natural product synthesis. In this article, the crystal structure of the title compound, ethyl 2-hydroxy-5-oxo-4-phenyl-2,3,4,5-tetrahydropyrano[3,2-*c*]chromene-2-carboxylate was described (Fig. 1). The main structural unit is a three-ring group consisting of a coumarin ring and a tetrahydropyrane. Two carbon atoms of the tetrahydropyrane structure are not coplanar with the coumarin backbone: one carbon atom lies 0.295 (3) Å from the mean plane of coumarin ring and the other lies 0.360 (2) Å from the plane in opposite direction. The dihedral angle between benzene and coumarin rings is 73.94 (3) °. The distance from O1 of the hydroxyl group to coumarin plane is 1.664 (2) Å. In addition, intermolecular O—H···O hydrogen bonds in the crystal are observed (Fig. 2).

S2. Experimental


The title compound was synthesized by treating (*E*)-ethyl 2-oxo-4-phenylbut-3-enoate (2.04 g, 10 mmol) with 4-hydroxycoumarin (1.62 g, 10 mmol) in the presence of triethylamine as a catalyst in dichloromethane (30 ml) under stirring at room temperature for 24 h. The solvent was distilled under vacuum, and the residue was purified by flash column chromatography (silica gel, Hex/AcOEt, *v/v*, 3:1) giving the title compound (3.3 g, 90%). The compound (*E*)-ethyl 2-oxo-4-phenylbut-3-enoate was obtained from commercially available benzaldehyde by condensation with pyruvic acid and subsequent esterification with ethanol. Suitable crystals of the title compound were obtained by slow evaporation of dichloromethane solution at room temperature.

S3. Refinement

H atoms were placed in calculated position with C—H=0.98 Å(*sp*), C—H=0.97 Å(*sp*²), C—H=0.96 Å(*sp*³), C—H=0.93 Å(aromatic). All H atoms included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}$ of the carrier atoms.

**Figure 1**

The asymmetric unit of the structure of the title compound, with the atomic labeling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The molecular packing of the title compound showing H-bond interactions. Symmetry code (i) is 0.5-x, 0.5+y, 0.5-z.

Ethyl 2-hydroxy-5-oxo-4-phenyl-2,3,4,5- tetrahydropyrano[3,2-c]chromene-2-carboxylate

Crystal data

$C_{21}H_{18}O_6$	$F(000) = 768.00$
$M_r = 366.37$	$D_x = 1.398 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71075 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 11365 reflections
$a = 5.4988 (2) \text{ \AA}$	$\theta = 3.2\text{--}27.4^\circ$
$b = 14.9975 (5) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 21.342 (1) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 98.5487 (13)^\circ$	Platelet, colorless
$V = 1740.48 (12) \text{ \AA}^3$	$0.41 \times 0.39 \times 0.14 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-AXIS RAPID diffractometer	3422 independent reflections
Detector resolution: 10.00 pixels mm^{-1}	2568 reflections with $F^2 > 2\sigma(F^2)$
ω scans	$R_{\text{int}} = 0.026$
Absorption correction: multi-scan (ABSCOR; Higashi, 2005)	$\theta_{\text{max}} = 27.4^\circ$
$T_{\text{min}} = 0.954$, $T_{\text{max}} = 0.986$	$h = -7 \rightarrow 7$
16523 measured reflections	$k = -19 \rightarrow 16$
	$l = -27 \rightarrow 27$

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.013P)^2 + P]$
$R[F^2 > 2\sigma(F^2)] = 0.035$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.080$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
3422 reflections	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
246 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008)
0 restraints	Extinction coefficient: 0.0057 (4)
H-atom parameters constrained	

Special details

Geometry. The tetrahydropyrane structure in the crystal displays an envelope configuration, with atom C2 at the flap position, displaced by 0.603 (2) Å from the mean plane of the other atoms.

Refinement. Refinement using reflections with $F^2 > 2.0 \sigma(F^2)$. The weighted R -factor (wR), goodness of fit (S) and R -factor (gt) are based on F , with F set to zero for negative F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.3270 (2)	0.47260 (6)	0.24846 (5)	0.0400 (2)
O2	0.3755 (2)	0.12605 (6)	0.19275 (5)	0.0442 (3)
O3	0.2981 (2)	0.39483 (6)	0.15453 (5)	0.0369 (2)
O4	0.7169 (2)	0.14798 (8)	0.25944 (6)	0.0548 (3)
O5	0.2708 (2)	0.60253 (9)	0.16249 (6)	0.0562 (3)
O6	0.5597 (2)	0.54329 (8)	0.11100 (6)	0.0483 (3)
C1	0.4400 (2)	0.46019 (10)	0.19571 (6)	0.0320 (3)
C2	0.7020 (2)	0.42864 (10)	0.21432 (8)	0.0344 (3)
C3	0.7153 (2)	0.33839 (10)	0.24884 (6)	0.0326 (3)

C4	0.5223 (2)	0.27767 (10)	0.21402 (6)	0.0322 (3)
C5	0.5514 (3)	0.18281 (11)	0.22491 (8)	0.0383 (3)
C6	0.1792 (3)	0.15776 (11)	0.15092 (6)	0.0390 (3)
C7	0.0084 (3)	0.09575 (12)	0.12389 (8)	0.0504 (4)
C8	-0.1910 (3)	0.12559 (12)	0.08244 (9)	0.0548 (5)
C9	-0.2214 (3)	0.21535 (12)	0.06798 (8)	0.0490 (4)
C10	-0.0505 (2)	0.27653 (11)	0.09506 (6)	0.0394 (3)
C11	0.1532 (2)	0.24823 (10)	0.13760 (6)	0.0335 (3)
C12	0.3355 (2)	0.30795 (10)	0.17075 (6)	0.0312 (3)
C13	0.7091 (2)	0.34673 (10)	0.31968 (6)	0.0334 (3)
C14	0.9040 (3)	0.38961 (12)	0.35684 (8)	0.0429 (4)
C15	0.9097 (3)	0.39905 (12)	0.42137 (9)	0.0526 (4)
C16	0.7198 (3)	0.36628 (13)	0.45022 (9)	0.0554 (5)
C17	0.5260 (3)	0.32322 (13)	0.41407 (9)	0.0526 (4)
C18	0.5209 (3)	0.31326 (12)	0.34917 (8)	0.0427 (4)
C19	0.4127 (2)	0.54477 (11)	0.15467 (8)	0.0368 (3)
C20	0.5607 (4)	0.62325 (13)	0.07224 (10)	0.0632 (5)
C21	0.7368 (4)	0.60702 (17)	0.02723 (11)	0.0822 (7)
H3	0.8750	0.3122	0.2443	0.039*
H7	0.0279	0.0354	0.1335	0.060*
H8	-0.3073	0.0848	0.0638	0.066*
H9	-0.3574	0.2343	0.0400	0.059*
H10	-0.0706	0.3367	0.0851	0.047*
H11	0.7910	0.4727	0.2421	0.041*
H12	0.7781	0.4226	0.1763	0.041*
H14	1.0325	0.4123	0.3378	0.051*
H15	1.0420	0.4276	0.4455	0.063*
H16	0.7223	0.3731	0.4936	0.066*
H17	0.3979	0.3007	0.4333	0.063*
H18	0.3898	0.2838	0.3253	0.051*
H101	0.2350	0.5159	0.2417	0.048*
H201	0.6127	0.6743	0.0988	0.076*
H202	0.3976	0.6341	0.0492	0.076*
H211	0.8950	0.5924	0.0504	0.099*
H212	0.7504	0.6597	0.0025	0.099*
H213	0.6791	0.5585	-0.0003	0.099*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0426 (6)	0.0374 (6)	0.0417 (6)	0.0076 (5)	0.0117 (5)	0.0020 (5)
O2	0.0565 (7)	0.0292 (5)	0.0443 (6)	-0.0051 (5)	-0.0012 (5)	0.0017 (5)
O3	0.0413 (6)	0.0266 (5)	0.0394 (6)	-0.0019 (4)	-0.0047 (4)	0.0001 (4)
O4	0.0603 (8)	0.0355 (6)	0.0628 (8)	0.0082 (6)	-0.0107 (6)	0.0050 (6)
O5	0.0556 (7)	0.0415 (7)	0.0723 (9)	0.0130 (6)	0.0120 (6)	0.0145 (6)
O6	0.0652 (8)	0.0371 (6)	0.0445 (7)	-0.0020 (5)	0.0149 (6)	0.0096 (5)
C1	0.0357 (8)	0.0282 (7)	0.0322 (8)	-0.0027 (6)	0.0054 (6)	-0.0020 (6)
C2	0.0326 (7)	0.0315 (8)	0.0393 (8)	-0.0032 (6)	0.0056 (6)	0.0008 (6)

C3	0.0293 (7)	0.0318 (8)	0.0364 (8)	0.0025 (6)	0.0041 (6)	-0.0009 (6)
C4	0.0358 (8)	0.0281 (7)	0.0329 (8)	-0.0006 (6)	0.0059 (6)	-0.0011 (6)
C5	0.0457 (9)	0.0324 (8)	0.0365 (8)	-0.0014 (7)	0.0053 (7)	-0.0002 (7)
C6	0.0492 (9)	0.0373 (9)	0.0303 (8)	-0.0064 (7)	0.0049 (7)	-0.0005 (6)
C7	0.0719 (12)	0.0364 (9)	0.0404 (10)	-0.0173 (9)	0.0005 (9)	0.0006 (7)
C8	0.0683 (12)	0.0529 (11)	0.0398 (10)	-0.0267 (10)	-0.0031 (9)	-0.0026 (8)
C9	0.0507 (10)	0.0559 (11)	0.0376 (9)	-0.0111 (9)	-0.0025 (7)	-0.0013 (8)
C10	0.0457 (9)	0.0386 (9)	0.0333 (8)	-0.0048 (7)	0.0040 (7)	-0.0014 (7)
C11	0.0401 (8)	0.0331 (8)	0.0276 (7)	-0.0054 (7)	0.0064 (6)	-0.0032 (6)
C12	0.0370 (8)	0.0273 (7)	0.0304 (7)	-0.0023 (6)	0.0089 (6)	-0.0010 (6)
C13	0.0335 (7)	0.0303 (8)	0.0355 (8)	0.0065 (6)	0.0016 (6)	0.0027 (6)
C14	0.0405 (9)	0.0452 (10)	0.0408 (9)	0.0015 (7)	-0.0014 (7)	-0.0003 (7)
C15	0.0515 (10)	0.0561 (11)	0.0445 (10)	0.0083 (9)	-0.0117 (8)	-0.0060 (8)
C16	0.0647 (12)	0.0645 (12)	0.0351 (9)	0.0220 (10)	0.0014 (9)	-0.0008 (9)
C17	0.0525 (10)	0.0642 (12)	0.0433 (10)	0.0116 (9)	0.0146 (8)	0.0081 (9)
C18	0.0394 (9)	0.0465 (10)	0.0416 (9)	0.0015 (7)	0.0042 (7)	0.0010 (7)
C19	0.0398 (8)	0.0317 (8)	0.0372 (9)	-0.0057 (7)	0.0005 (7)	0.0013 (6)
C20	0.0818 (14)	0.0490 (11)	0.0601 (13)	-0.0083 (10)	0.0147 (11)	0.0222 (9)
C21	0.1051 (19)	0.0793 (16)	0.0674 (15)	-0.0195 (14)	0.0301 (14)	0.0151 (13)

Geometric parameters (Å, °)

O1—C1	1.3772 (19)	C13—C18	1.383 (2)
O2—C5	1.3906 (19)	C14—C15	1.380 (2)
O2—C6	1.3791 (18)	C15—C16	1.379 (2)
O3—C1	1.4623 (17)	C16—C17	1.379 (2)
O3—C12	1.3560 (18)	C17—C18	1.389 (2)
O4—C5	1.2019 (19)	C20—C21	1.482 (3)
O5—C19	1.194 (2)	O1—H101	0.822
O6—C19	1.321 (2)	C2—H11	0.970
O6—C20	1.457 (2)	C2—H12	0.970
C1—C2	1.512 (2)	C3—H3	0.980
C1—C19	1.536 (2)	C7—H7	0.930
C2—C3	1.538 (2)	C8—H8	0.930
C3—C4	1.508 (2)	C9—H9	0.930
C3—C13	1.522 (2)	C10—H10	0.930
C4—C5	1.447 (2)	C14—H14	0.930
C4—C12	1.353 (2)	C15—H15	0.930
C6—C7	1.385 (2)	C16—H16	0.930
C6—C11	1.389 (2)	C17—H17	0.930
C7—C8	1.377 (2)	C18—H18	0.930
C8—C9	1.386 (2)	C20—H201	0.970
C9—C10	1.378 (2)	C20—H202	0.970
C10—C11	1.399 (2)	C21—H211	0.960
C11—C12	1.448 (2)	C21—H212	0.960
C13—C14	1.392 (2)	C21—H213	0.960
C5—O2—C6	121.90 (12)	O5—C19—C1	122.00 (16)

C1—O3—C12	116.34 (10)	O6—C19—C1	111.72 (13)
C19—O6—C20	116.16 (14)	O6—C20—C21	106.91 (17)
O1—C1—O3	108.45 (12)	C1—O1—H101	107.7
O1—C1—C2	110.84 (12)	C1—C2—H11	108.8
O1—C1—C19	109.68 (12)	C1—C2—H12	108.8
O3—C1—C2	110.52 (12)	C3—C2—H11	108.8
O3—C1—C19	102.28 (11)	C3—C2—H12	108.8
C2—C1—C19	114.63 (13)	H11—C2—H12	109.5
C1—C2—C3	112.15 (12)	C2—C3—H3	106.5
C2—C3—C4	108.40 (11)	C4—C3—H3	106.5
C2—C3—C13	113.39 (12)	C13—C3—H3	106.5
C4—C3—C13	114.91 (12)	C6—C7—H7	120.8
C3—C4—C5	117.50 (12)	C8—C7—H7	120.8
C3—C4—C12	122.74 (13)	C7—C8—H8	119.4
C5—C4—C12	119.62 (13)	C9—C8—H8	119.4
O2—C5—O4	116.37 (14)	C8—C9—H9	119.9
O2—C5—C4	118.06 (13)	C10—C9—H9	119.9
O4—C5—C4	125.57 (14)	C9—C10—H10	120.0
O2—C6—C7	117.07 (14)	C11—C10—H10	120.0
O2—C6—C11	121.07 (13)	C13—C14—H14	119.4
C7—C6—C11	121.85 (14)	C15—C14—H14	119.4
C6—C7—C8	118.37 (16)	C14—C15—H15	119.9
C7—C8—C9	121.12 (17)	C16—C15—H15	119.9
C8—C9—C10	120.10 (15)	C15—C16—H16	120.3
C9—C10—C11	120.05 (15)	C17—C16—H16	120.3
C6—C11—C10	118.50 (13)	C16—C17—H17	119.8
C6—C11—C12	117.48 (12)	C18—C17—H17	119.8
C10—C11—C12	123.97 (13)	C13—C18—H18	119.7
O3—C12—C4	124.57 (13)	C17—C18—H18	119.7
O3—C12—C11	113.70 (12)	O6—C20—H201	110.1
C4—C12—C11	121.72 (13)	O6—C20—H202	110.1
C3—C13—C14	118.36 (14)	C21—C20—H201	110.1
C3—C13—C18	123.41 (13)	C21—C20—H202	110.1
C14—C13—C18	118.23 (14)	H201—C20—H202	109.5
C13—C14—C15	121.16 (16)	C20—C21—H211	109.5
C14—C15—C16	120.15 (16)	C20—C21—H212	109.5
C15—C16—C17	119.36 (17)	C20—C21—H213	109.5
C16—C17—C18	120.49 (18)	H211—C21—H212	109.5
C13—C18—C17	120.61 (15)	H211—C21—H213	109.5
O5—C19—O6	126.28 (16)	H212—C21—H213	109.5
C5—O2—C6—C7	-176.87 (15)	C3—C4—C5—O2	-179.57 (14)
C5—O2—C6—C11	2.1 (2)	C3—C4—C5—O4	-0.0 (2)
C6—O2—C5—O4	-178.82 (15)	C3—C4—C12—O3	-1.0 (2)
C6—O2—C5—C4	0.8 (2)	C3—C4—C12—C11	179.57 (14)
C1—O3—C12—C4	-12.7 (2)	C5—C4—C12—O3	-176.60 (15)
C1—O3—C12—C11	166.76 (13)	C5—C4—C12—C11	4.0 (2)
C12—O3—C1—O1	-79.87 (15)	C12—C4—C5—O2	-3.8 (2)

C12—O3—C1—C2	41.82 (17)	C12—C4—C5—O4	175.79 (17)
C12—O3—C1—C19	164.28 (13)	O2—C6—C7—C8	179.32 (16)
C19—O6—C20—C21	179.82 (15)	O2—C6—C11—C10	-179.62 (14)
C20—O6—C19—O5	5.1 (2)	O2—C6—C11—C12	-1.9 (2)
C20—O6—C19—C1	-175.90 (13)	C7—C6—C11—C10	-0.7 (2)
O1—C1—C2—C3	61.60 (16)	C7—C6—C11—C12	176.99 (16)
O1—C1—C19—O5	-14.1 (2)	C11—C6—C7—C8	0.4 (2)
O1—C1—C19—O6	166.88 (12)	C6—C7—C8—C9	-0.1 (2)
O3—C1—C2—C3	-58.67 (16)	C7—C8—C9—C10	0.1 (2)
O3—C1—C19—O5	100.86 (17)	C8—C9—C10—C11	-0.5 (2)
O3—C1—C19—O6	-78.16 (14)	C9—C10—C11—C6	0.8 (2)
C2—C1—C19—O5	-139.51 (16)	C9—C10—C11—C12	-176.80 (16)
C2—C1—C19—O6	41.46 (18)	C6—C11—C12—O3	179.34 (14)
C19—C1—C2—C3	-173.59 (12)	C6—C11—C12—C4	-1.2 (2)
C1—C2—C3—C4	43.95 (17)	C10—C11—C12—O3	-3.1 (2)
C1—C2—C3—C13	-84.91 (15)	C10—C11—C12—C4	176.41 (16)
C2—C3—C4—C5	160.47 (14)	C3—C13—C14—C15	-179.77 (15)
C2—C3—C4—C12	-15.2 (2)	C3—C13—C18—C17	-179.88 (15)
C2—C3—C13—C14	-64.76 (18)	C14—C13—C18—C17	0.7 (2)
C2—C3—C13—C18	115.79 (16)	C18—C13—C14—C15	-0.3 (2)
C4—C3—C13—C14	169.80 (14)	C13—C14—C15—C16	-0.4 (2)
C4—C3—C13—C18	-9.7 (2)	C14—C15—C16—C17	0.7 (2)
C13—C3—C4—C5	-71.53 (18)	C15—C16—C17—C18	-0.4 (3)
C13—C3—C4—C12	112.80 (17)	C16—C17—C18—C13	-0.4 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H101...O2 ⁱ	0.82	2.30	2.9198 (15)	132
O1—H101...O5	0.82	2.17	2.6628 (17)	119

Symmetry code: (i) $-x+1/2, y+1/2, -z+1/2$.