

1,3-Dimethyl-5-(2-methylbenzylidene)-pyrimidine-2,4,6(1H,3H,5H)-trione

R. Panchatcharam,^a V. Dhayalan,^b
A. K. Mohanakrishnan,^b G. Chakkavarthi^c and
V. Manivannan^{a*}

^aCentre for Research and Development, PRIST University, Vallam, Thanjavur 613 403, Tamil Nadu, India, ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, and ^cDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India
Correspondence e-mail: manivan_1999@yahoo.com


Received 2 September 2009; accepted 2 September 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.072; wR factor = 0.238; data-to-parameter ratio = 21.9.

In the title compound, $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_3$, the dihedral angle between the pyrimidine and benzene rings is $14.9(1)^\circ$. The molecular structure is stabilized by weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ interactions and the crystal structure exhibits a weak intermolecular $\pi-\pi$ interaction [centroid–centroid distance = $3.575(3)\text{ \AA}$].

Related literature

For the biological activity of pyrimidine derivatives, see: Cody *et al.* (1997); Li *et al.* (1995). For related structures, see: Da Silva *et al.* (2005); Rezende *et al.* (2005). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_3$

$M_r = 258.27$

Monoclinic, $P2_1/n$
 $a = 8.182(5)\text{ \AA}$
 $b = 8.334(4)\text{ \AA}$
 $c = 18.202(5)\text{ \AA}$
 $\beta = 94.267(5)^\circ$
 $V = 1237.7(10)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.30 \times 0.28 \times 0.18\text{ mm}$

Data collection

Bruker Kappa APEXII
diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.971$, $T_{\max} = 0.982$

16198 measured reflections
3837 independent reflections
2517 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.238$
 $S = 1.04$
3837 reflections

175 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.52\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.34\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots\text{O}2$	0.93	2.26	2.893 (3)	125

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge SAIF, IIT, Madras, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5051).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cody, V., Galitsky, N., Luft, J. R., Pangborn, W., Gangjee, A., Devraj, R., Queener, S. F. & Blakley, R. L. (1997). *Acta Cryst. D* **53**, 638–649.
- Li, J. J., Anderson, D., Burton, E. G. & Cogburn, J. N. (1995). *J. Med. Chem.* **38**, 4570–4578.
- Rezende, M. C., Dominguez, M., Wardell, J. L., Skakle, J. M. S., Low, J. N. & Glidewell, C. (2005). *Acta Cryst. C* **61**, o306–o311.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Silva, E. T. da, Ribiero, R. S., Lima, E. L. S., Wardell, J. L., Skakle, J. M. S., Low, J. N. & Glidewell, C. (2005). *Acta Cryst. C* **61**, o15–o20.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2009). E65, o2394 [doi:10.1107/S1600536809035521]

1,3-Dimethyl-5-(2-methylbenzylidene)pyrimidine-2,4,6(1*H*,3*H*,5*H*)-trione

R. Panchatcharam, V. Dhayalan, A. K. Mohanakrishnan, G. Chakkavarthi and V. Manivannan

S1. Comment

Pyrimidine derivatives show biological activities such as antitumor, antibacterial, insulin releasing and anti-inflammatory activities (Cody *et al.*, 1997; Li *et al.*, 1995). The geometric parameters in (I) (Fig. 1) agree with the reported values of similar structures (Da Silva *et al.*, 2005; Rezende *et al.*, 2005).

The dihedral angle between the pyrimidine ring (N1/C11/N2/C10/C9/C12) and benzene ring (C1—C6) is 14.9 (1) $^{\circ}$. The molecular structure is stabilized by weak intramolecular C—H···O interactions. The crystal structure exhibits an intermolecular weak π – π interaction [$Cg1\cdots Cg2 = 3.575$ (3) Å; symmetry code: $-x, -y, 1 - z$; $Cg1$ and $Cg2$ are the centroids of N1/C11/N2/C10/C9/C12 and C1—C6 rings, respectively].


The intramolecular C8—H8···O1 and C13—H13B···O3 interactions generate five-membered rings, each with graph-set motif S(5) and C2—H2···O2 interaction generates a seven-membered ring, with graph-set motif S(7) (Bernstein *et al.*, 1995).

S2. Experimental

To a solution of *o*-tolualdehyde (4.0 g, 33.29 mmol) in dry benzene (80 ml), *N,N*-dimethylbarbituric acid (5.72 g, 36.63 mmol), piperidine (0.6 ml) and acetic acid (0.3 ml) were added and refluxed in a RB flask fitted with Dean-Stark apparatus for 12 h. Removal of solvent followed by recrystallization from CDCl₃ afforded the compound.

S3. Refinement

H atoms were positioned geometrically and refined using riding model with C—H = 0.93 Å and $U_{iso}(\text{H}) = 1.2U_{eq}(\text{C})$ for C—H and C—H = 0.96 Å and $U_{iso}(\text{H}) = 1.5U_{eq}(\text{C})$ for CH₃.

**Figure 1**

The molecular structure of (I), with atom labels and 30% probability displacement ellipsoids for non-H atoms.

1,3-Dimethyl-5-(2-methylbenzylidene)pyrimidine- 2,4,6(1H,3H,5H)-trione

Crystal data

$C_{14}H_{14}N_2O_3$
 $M_r = 258.27$
Monoclinic, $P2_1/n$
Hall symbol: -P 2yn
 $a = 8.182 (5)$ Å
 $b = 8.334 (4)$ Å
 $c = 18.202 (5)$ Å
 $\beta = 94.267 (5)$ °
 $V = 1237.7 (10)$ Å³
 $Z = 4$

$F(000) = 544$
 $D_x = 1.386$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6022 reflections
 $\theta = 2.2\text{--}29.8$ °
 $\mu = 0.10$ mm⁻¹
 $T = 295$ K
Block, colourless
 $0.30 \times 0.28 \times 0.18$ mm

Data collection

Bruker Kappa APEXII
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω and φ scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.971$, $T_{\max} = 0.982$

16198 measured reflections
3837 independent reflections
2517 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
 $\theta_{\max} = 30.7$ °, $\theta_{\min} = 2.2$ °
 $h = -11 \rightarrow 11$
 $k = -8 \rightarrow 11$
 $l = -26 \rightarrow 26$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.238$
 $S = 1.04$
3837 reflections
175 parameters

0 restraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1214P)^2 + 0.5641P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.52 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0138 (2)	0.2553 (2)	0.49118 (10)	0.0391 (4)
C2	-0.1183 (3)	0.2104 (3)	0.44177 (12)	0.0547 (6)
H2	-0.1974	0.1406	0.4573	0.066*
C3	-0.1326 (3)	0.2681 (4)	0.37060 (12)	0.0589 (6)
H3	-0.2205	0.2370	0.3384	0.071*
C4	-0.0168 (3)	0.3713 (3)	0.34746 (12)	0.0548 (6)
H4	-0.0285	0.4147	0.3003	0.066*
C5	0.1166 (3)	0.4106 (3)	0.39422 (12)	0.0519 (5)
H5	0.1961	0.4785	0.3775	0.062*
C6	0.1363 (2)	0.3521 (3)	0.46527 (11)	0.0424 (5)
C7	0.2861 (3)	0.4007 (4)	0.51198 (15)	0.0714 (8)
H7A	0.3586	0.3105	0.5188	0.107*
H7B	0.2552	0.4370	0.5590	0.107*
H7C	0.3408	0.4859	0.4881	0.107*
C8	0.0343 (3)	0.2024 (3)	0.56714 (11)	0.0447 (5)
H8	0.1412	0.2161	0.5873	0.054*
C9	-0.0641 (2)	0.1379 (2)	0.61602 (10)	0.0390 (4)
C10	0.0217 (3)	0.0920 (3)	0.68754 (10)	0.0429 (5)
C11	-0.2160 (2)	-0.0676 (2)	0.71455 (10)	0.0396 (4)
C12	-0.2395 (3)	0.1068 (3)	0.60570 (11)	0.0449 (5)
C13	0.0281 (3)	-0.0647 (3)	0.79980 (12)	0.0584 (6)
H13A	0.1247	-0.1223	0.7885	0.088*
H13B	-0.0416	-0.1340	0.8257	0.088*
H13C	0.0589	0.0263	0.8300	0.088*
C14	-0.4706 (3)	-0.0581 (4)	0.63701 (16)	0.0715 (8)
H14A	-0.4822	-0.1675	0.6524	0.107*
H14B	-0.5004	-0.0496	0.5851	0.107*
H14C	-0.5409	0.0093	0.6636	0.107*
N1	-0.3007 (2)	-0.0073 (2)	0.65217 (9)	0.0433 (4)
N2	-0.0600 (2)	-0.0100 (2)	0.73120 (8)	0.0397 (4)
O1	0.1595 (2)	0.1369 (3)	0.70726 (9)	0.0682 (6)
O2	-0.3317 (2)	0.1744 (3)	0.56044 (11)	0.0748 (6)
O3	-0.2774 (2)	-0.1642 (2)	0.75393 (10)	0.0587 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0383 (9)	0.0465 (10)	0.0326 (8)	0.0012 (8)	0.0033 (7)	0.0046 (7)
C2	0.0453 (11)	0.0775 (16)	0.0407 (10)	-0.0112 (11)	-0.0004 (9)	0.0047 (11)
C3	0.0505 (12)	0.0883 (18)	0.0367 (10)	0.0003 (12)	-0.0041 (9)	0.0010 (11)
C4	0.0654 (14)	0.0672 (15)	0.0323 (9)	0.0100 (11)	0.0064 (9)	0.0082 (9)

C5	0.0614 (13)	0.0556 (13)	0.0403 (10)	-0.0084 (10)	0.0144 (9)	0.0035 (9)
C6	0.0407 (10)	0.0498 (11)	0.0371 (9)	-0.0012 (8)	0.0061 (7)	-0.0031 (8)
C7	0.0539 (14)	0.104 (2)	0.0549 (14)	-0.0309 (15)	-0.0017 (11)	0.0038 (14)
C8	0.0417 (10)	0.0534 (12)	0.0382 (9)	-0.0060 (9)	-0.0016 (8)	0.0071 (9)
C9	0.0413 (10)	0.0433 (10)	0.0320 (8)	-0.0013 (8)	-0.0009 (7)	0.0049 (7)
C10	0.0415 (10)	0.0535 (11)	0.0330 (9)	-0.0047 (9)	-0.0017 (7)	0.0061 (8)
C11	0.0460 (10)	0.0390 (10)	0.0350 (9)	-0.0009 (8)	0.0100 (7)	-0.0014 (7)
C12	0.0391 (10)	0.0608 (13)	0.0345 (9)	0.0025 (9)	0.0008 (7)	0.0053 (9)
C13	0.0598 (14)	0.0761 (16)	0.0384 (10)	0.0067 (12)	-0.0022 (9)	0.0186 (11)
C14	0.0449 (13)	0.108 (2)	0.0611 (15)	-0.0255 (14)	0.0006 (11)	-0.0008 (15)
N1	0.0378 (8)	0.0532 (10)	0.0387 (8)	-0.0071 (7)	0.0025 (6)	-0.0027 (7)
N2	0.0408 (8)	0.0466 (9)	0.0315 (7)	0.0021 (7)	0.0020 (6)	0.0070 (6)
O1	0.0512 (9)	0.1035 (15)	0.0473 (9)	-0.0257 (9)	-0.0139 (7)	0.0199 (9)
O2	0.0443 (9)	0.1221 (17)	0.0570 (10)	0.0125 (10)	-0.0022 (8)	0.0328 (11)
O3	0.0673 (11)	0.0574 (10)	0.0536 (9)	-0.0123 (8)	0.0186 (8)	0.0089 (7)

Geometric parameters (\AA , $^{\circ}$)

C1—C6	1.396 (3)	C9—C12	1.457 (3)
C1—C2	1.404 (3)	C9—C10	1.482 (3)
C1—C8	1.449 (3)	C10—O1	1.217 (3)
C2—C3	1.379 (3)	C10—N2	1.371 (3)
C2—H2	0.9300	C11—O3	1.212 (2)
C3—C4	1.369 (4)	C11—N2	1.376 (3)
C3—H3	0.9300	C11—N1	1.380 (3)
C4—C5	1.373 (4)	C12—O2	1.214 (3)
C4—H4	0.9300	C12—N1	1.391 (3)
C5—C6	1.380 (3)	C13—N2	1.467 (3)
C5—H5	0.9300	C13—H13A	0.9600
C6—C7	1.495 (3)	C13—H13B	0.9600
C7—H7A	0.9600	C13—H13C	0.9600
C7—H7B	0.9600	C14—N1	1.460 (3)
C7—H7C	0.9600	C14—H14A	0.9600
C8—C9	1.355 (3)	C14—H14B	0.9600
C8—H8	0.9300	C14—H14C	0.9600
C6—C1—C2	118.38 (18)	C12—C9—C10	117.69 (17)
C6—C1—C8	117.61 (18)	O1—C10—N2	120.04 (18)
C2—C1—C8	123.97 (19)	O1—C10—C9	123.19 (19)
C3—C2—C1	121.1 (2)	N2—C10—C9	116.74 (18)
C3—C2—H2	119.5	O3—C11—N2	121.34 (19)
C1—C2—H2	119.5	O3—C11—N1	121.6 (2)
C4—C3—C2	119.7 (2)	N2—C11—N1	117.08 (17)
C4—C3—H3	120.1	O2—C12—N1	119.8 (2)
C2—C3—H3	120.1	O2—C12—C9	124.2 (2)
C3—C4—C5	119.7 (2)	N1—C12—C9	116.06 (17)
C3—C4—H4	120.1	N2—C13—H13A	109.5
C5—C4—H4	120.1	N2—C13—H13B	109.5

C4—C5—C6	121.9 (2)	H13A—C13—H13B	109.5
C4—C5—H5	119.0	N2—C13—H13C	109.5
C6—C5—H5	119.0	H13A—C13—H13C	109.5
C5—C6—C1	118.95 (19)	H13B—C13—H13C	109.5
C5—C6—C7	118.1 (2)	N1—C14—H14A	109.5
C1—C6—C7	122.9 (2)	N1—C14—H14B	109.5
C6—C7—H7A	109.5	H14A—C14—H14B	109.5
C6—C7—H7B	109.5	N1—C14—H14C	109.5
H7A—C7—H7B	109.5	H14A—C14—H14C	109.5
C6—C7—H7C	109.5	H14B—C14—H14C	109.5
H7A—C7—H7C	109.5	C11—N1—C12	124.58 (17)
H7B—C7—H7C	109.5	C11—N1—C14	117.58 (19)
C9—C8—C1	135.66 (19)	C12—N1—C14	117.7 (2)
C9—C8—H8	112.2	C10—N2—C11	124.92 (16)
C1—C8—H8	112.2	C10—N2—C13	117.15 (18)
C8—C9—C12	127.86 (18)	C11—N2—C13	117.91 (18)
C8—C9—C10	114.43 (18)		
C6—C1—C2—C3	-4.0 (4)	C10—C9—C12—O2	159.2 (2)
C8—C1—C2—C3	178.4 (2)	C8—C9—C12—N1	158.5 (2)
C1—C2—C3—C4	-0.2 (4)	C10—C9—C12—N1	-20.2 (3)
C2—C3—C4—C5	3.1 (4)	O3—C11—N1—C12	178.2 (2)
C3—C4—C5—C6	-1.7 (4)	N2—C11—N1—C12	-0.2 (3)
C4—C5—C6—C1	-2.5 (4)	O3—C11—N1—C14	2.9 (3)
C4—C5—C6—C7	179.6 (3)	N2—C11—N1—C14	-175.4 (2)
C2—C1—C6—C5	5.2 (3)	O2—C12—N1—C11	-166.3 (2)
C8—C1—C6—C5	-177.0 (2)	C9—C12—N1—C11	13.1 (3)
C2—C1—C6—C7	-177.0 (3)	O2—C12—N1—C14	8.9 (4)
C8—C1—C6—C7	0.7 (3)	C9—C12—N1—C14	-171.7 (2)
C6—C1—C8—C9	164.5 (3)	O1—C10—N2—C11	179.7 (2)
C2—C1—C8—C9	-17.9 (4)	C9—C10—N2—C11	-2.3 (3)
C1—C8—C9—C12	-3.4 (4)	O1—C10—N2—C13	-2.0 (3)
C1—C8—C9—C10	175.4 (2)	C9—C10—N2—C13	176.01 (19)
C8—C9—C10—O1	14.5 (3)	O3—C11—N2—C10	176.1 (2)
C12—C9—C10—O1	-166.7 (2)	N1—C11—N2—C10	-5.6 (3)
C8—C9—C10—N2	-163.5 (2)	O3—C11—N2—C13	-2.2 (3)
C12—C9—C10—N2	15.4 (3)	N1—C11—N2—C13	176.12 (18)
C8—C9—C12—O2	-22.1 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C2—H2 \cdots O2	0.93	2.26	2.893 (3)	125
C8—H8 \cdots O1	0.93	2.28	2.732 (3)	110
C13—H13B \cdots O3	0.96	2.26	2.706 (4)	107