

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(3-Bromobenzyl)-1H-isochromen-1-one

 Farukh Iftakhar Ali,^a Tariq Mahmood Babar,^a
 Nasim Hasan Rama^{a*} and Peter G. Jones^b
^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan, and ^bInstitut für Anorganische und Analytische Chemie, Technische Universität Braunschweig, Postfach 3329, 38023 Braunschweig, Germany
 Correspondence e-mail: nhrama@qau.edu.pk


Received 4 September 2009; accepted 14 September 2009

 Key indicators: single-crystal X-ray study; $T = 103$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.020; wR factor = 0.045; data-to-parameter ratio = 20.1.

In the title compound, $\text{C}_{16}\text{H}_{11}\text{BrO}_2$, the isocoumarin ring system is planar (r.m.s. deviation = 0.015 Å) and subtends a dihedral angle of 88.90 (2)° with the bromobenzene ring. In the crystal, molecules are linked, forming a three-dimensional packing pattern involving C—H···O interactions, Br···O contacts [3.4734 (10) Å] and π – π stacking interactions with centroid–centroid distances ranging from 3.667 (2) to 3.765 (2) Å.

Related literature

For the properties and applications of isocoumarins and 3,4-dihydroisocoumarins, see: Chinworrungsee *et al.* (2002); Devienne *et al.* (2002); Mali & Babu (1998); Rama *et al.* (1998); Waters & Kozlowski (2001). For related structures, see: Abid *et al.* (2008); Babar *et al.* (2008).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{11}\text{BrO}_2$	$b = 8.1824$ (6) Å
$M_r = 315.16$	$c = 11.3663$ (8) Å
Triclinic, $P\bar{1}$	$\alpha = 90.130$ (6)°
$a = 7.4508$ (5) Å	$\beta = 98.392$ (7)°

$\gamma = 113.844$ (8)°	$\mu = 3.28$ mm ⁻¹
$V = 625.58$ (8) Å ³	$T = 103$ K
$Z = 2$	$0.25 \times 0.25 \times 0.20$ mm
Mo $K\alpha$ radiation	

Data collection

Oxford Xcalibur E diffractometer	16250 measured reflections
Absorption correction: multi-scan (<i>CrysAlisPro</i> ; Oxford Diffraction 2009)	3449 independent reflections
$T_{\min} = 0.922$, $T_{\max} = 1.000$	2899 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.020$	172 parameters
$wR(F^2) = 0.045$	H-atom parameters constrained
$S = 0.97$	$\Delta\rho_{\text{max}} = 0.43$ e Å ⁻³
3449 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{O1}^{\text{i}}$	0.95	2.58	3.4666 (16)	155
$\text{C10}-\text{H10A}\cdots\text{O2}^{\text{ii}}$	0.99	2.50	3.4685 (17)	166

 Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, -y, -z + 1$.

Data collection: *CrysAlisPro* (Oxford Diffraction, 2009); cell refinement: *CrysAlisPro*; data reduction: *CrysAlisPro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Siemens, 1994); software used to prepare material for publication: *SHELXL97*.

TMB is grateful to the Higher Education Commission of Pakistan for financial support for a PhD program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2357).

References

- Abid, O.-U.-R., Qadeer, G., Rama, N. H., Ruzicka, A. & Padelkova, Z. (2008). *Acta Cryst.* **E64**, o2018.
- Babar, T. M., Qadeer, G., Abid, O.-R., Rama, N. H. & Ruzicka, A. (2008). *Acta Cryst.* **E64**, o2266.
- Chinworrungsee, M., Kittakoop, P., Isaka, M., Chanphen, R., Tanticharoen, M. & Thebtaranonth, Y. (2002). *J. Chem. Soc. Perkin Trans. 1*, pp. 2473–2476.
- Devienne, K. F., Raddi, M. S. G., Varanda, E. A. & Vilegas, W. (2002). *Z. Naturforsch. Teil C*, **57**, 85–88.
- Mali, R. S. & Babu, K. N. (1998). *J. Org. Chem.* **63**, 2288–2492.
- Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, England.
- Rama, N. H., Iqbal, R. & Zamani, K. (1998). *J. Chem. Soc. Pak.* **62**, 18–21.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1994). *XP*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Waters, S. P. & Kozlowski, M. C. (2001). *Tetrahedron Lett.* **42**, 3567–3570.

supporting information

Acta Cryst. (2009). E65, o2511 [doi:10.1107/S1600536809037246]

3-(3-Bromobenzyl)-1*H*-isochromen-1-one

Farukh Iftakhar Ali, Tariq Mahmood Babar, Nasim Hasan Rama and Peter G. Jones

S1. Comment

In recent years, there has been increasing interest in the synthesis of natural products, since they are an excellent and reliable source for the development of new drugs. Isocoumarins and 3,4-dihydroisocoumarins are a class of natural products that often occur as microbial metabolites and that have been found to exhibit interesting biological properties (Mali & Babu, 1998), including anti-fungal, anti-inflammatory, anti-allergic, antiangiogenic, anti-malaria (Chinworrungsee *et al.*, 2002), anti-bacterial (Rama *et al.*, 1998), anti-cancer, anti-virus (Waters & Kozlowski, 2001) and anti-microbial activities (Davienne *et al.*, 2002). In view of the importance of this class of compounds, the title compound, an isocoumarine derivative containing a 3-bromobenzyl substituent, has been synthesized and its crystal structure is reported here. We have previously reported the structures of the analogous fluorine and chlorine derivatives (Babar *et al.*, 2008; Abid *et al.*, 2008), which crystallize with two and three molecules respectively in the asymmetric unit.

The molecule of the title compound is shown in Fig. 1. The structure is not isotopic to either of the analogous derivatives. Bond lengths and angles may be regarded as normal by comparison with the earlier structures (Babar *et al.*, 2008; Abid *et al.*, 2008), although in each structure several bond angles are appreciably different from ideal values [*e. g.* in the current structure O2—C1—C9 126.19 (12), C3—C2—C10 128.78 (12), O1—C2—C10 109.56 (11), C2—C10—C11 112.98 (11)°]. The isocoumarin ring system and the bromobenzene ring are both planar within r.m.s. deviations of 0.015 Å and subtend a dihedral angle of 88.90 (2)°.

The packing diagram (Fig. 2) shows the molecules to be linked by two C—H···O hydrogen bonds (Table 1) and by π – π stacking interactions between the coumarin units and between the bromobenzene rings, with centroid-to-centroid distances ranging from 3.667 (2) to 3.765 (2) Å. A marginal Br···O interaction [Br···O2ⁱ 3.4734 (10) Å; symmetry code: (i) -1 + x, y, -1 + z] is also observed.

S2. Experimental

A mixture of 2-(3-bromophenyl)acetic acid (5 g, 0.023 mol) and oxalyl chloride (2 ml, 0.024 mol) was stirred overnight. Completion of the reaction was indicated by cessation of gas evolution. Excess oxalyl chloride was removed under reduced pressure to afford 2-(3-bromophenyl)acetyl chloride. Homophthalic acid (1.0 g, 0.006 mol) was added and the solution was heated at 473 K for 4 h. The reaction mixture was dissolved in ethyl acetate and aqueous solution of sodium carbonate was added in order to remove the unreacted homophthalic acid. The organic layer was separated, concentrated and chromatographed on silica gel using pet ether as eluent to afford title compound (yield 65%,; m.p. 97–98°C) as a colourless solid. Crystals suitable for X-ray analysis were obtained by slow evaporation of an ethanol solution.

S3. Refinement

H atoms were placed in calculated positions and refined using a riding model with C—H = 0.95–0.99 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

Molecular structure of the title compound showing the atom labelling scheme and displacement ellipsoids at the 50% probability level.




Figure 2

Packing diagram of the title compound viewed perpendicular to the bc plane. C—H...O hydrogen interactions are indicated by dashed lines. H atoms not involved in hydrogen bonding are omitted.




Figure 3

Reaction scheme.

3-(3-Bromobenzyl)-1*H*-isochromen-1-one

Crystal data

$C_{16}H_{11}BrO_2$

$M_r = 315.16$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.4508$ (5) Å

$b = 8.1824$ (6) Å

$c = 11.3663$ (8) Å

$\alpha = 90.130$ (6)°

$\beta = 98.392$ (7)°

$\gamma = 113.844$ (8)°

$V = 625.58$ (8) Å³

$Z = 2$

$F(000) = 316$
 $D_x = 1.673 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 9750 reflections
 $\theta = 2.7\text{--}30.7^\circ$

$\mu = 3.28 \text{ mm}^{-1}$
 $T = 103 \text{ K}$
 Block, colourless
 $0.25 \times 0.25 \times 0.20 \text{ mm}$

Data collection

Oxford Xcalibur E
 diffractometer
 Radiation source: Enhance (Mo) X-ray Source
 Graphite monochromator
 Detector resolution: 16.1419 pixels mm^{-1}
 ω scan
 Absorption correction: multi-scan
 (CrysAlis PRO; Oxford Diffraction 2009)
 $T_{\min} = 0.922$, $T_{\max} = 1.000$

16250 measured reflections
 3449 independent reflections
 2899 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 29.6^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -10 \rightarrow 10$
 $k = -11 \rightarrow 11$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.020$
 $wR(F^2) = 0.045$
 $S = 0.97$
 3449 reflections
 172 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0233P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.43 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Non-bonded contact:

3.4734 (0.0010) Br - O2_5 157.19 (0.04) C13 - Br - O2_5 144.68 (0.08) Br - O2_5 - C1_5 Operator for generating equivalent atoms: 5 $x - 1, y, z - 1$

Least-squares planes (x, y, z in crystal coordinates) and deviations from them (* indicates atom used to define plane)
 $-1.8912 (0.0020) x + 7.3803 (0.0014) y + 4.4053 (0.0037) z = 0.7728 (0.0010)$

* 0.0088 (0.0008) C10 * -0.0063 (0.0011) C11 * 0.0058 (0.0010) C12 * 0.0167 (0.0011) C13 * 0.0215 (0.0010) C14 *
 $-0.0036 (0.0009) C15 * -0.0224 (0.0011) C16 * -0.0204 (0.0006) \text{ Br}$

Rms deviation of fitted atoms = 0.0151

7.0134 (0.0011) $x - 0.6456 (0.0018) y - 2.6242 (0.0032) z = 3.6495 (0.0018)$

Angle to previous plane (with approximate e.s.d.) = 88.90 (0.02)

* -0.0183 (0.0009) C10 * -0.0095 (0.0009) O1 * 0.0027 (0.0011) C1 * -0.0081 (0.0008) O2 * 0.0273 (0.0011) C3 *
 $0.0129 (0.0012) C4 * -0.0052 (0.0011) C5 * -0.0241 (0.0011) C6 * -0.0078 (0.0012) C7 * 0.0144 (0.0012) C8 * 0.0159$
 $(0.0012) C9$

Rms deviation of fitted atoms = 0.0152

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.13942 (2)	0.205557 (19)	-0.113746 (13)	0.02023 (5)
O1	0.70881 (14)	0.19099 (12)	0.46026 (8)	0.0146 (2)
O2	0.78538 (15)	0.22208 (13)	0.65672 (8)	0.0195 (2)
C1	0.75969 (19)	0.29281 (17)	0.56653 (12)	0.0141 (3)
C2	0.67678 (19)	0.25762 (17)	0.35091 (11)	0.0128 (3)
C3	0.69140 (19)	0.42394 (17)	0.34239 (11)	0.0136 (3)
H3	0.6699	0.4662	0.2661	0.016*
C4	0.73964 (18)	0.54146 (17)	0.44789 (11)	0.0124 (3)
C5	0.7521 (2)	0.71742 (18)	0.44484 (12)	0.0170 (3)
H5	0.7288	0.7643	0.3706	0.020*
C6	0.7982 (2)	0.82214 (18)	0.54931 (13)	0.0182 (3)
H6	0.8045	0.9404	0.5465	0.022*
C7	0.8354 (2)	0.75669 (19)	0.65887 (13)	0.0185 (3)
H7	0.8683	0.8306	0.7301	0.022*
C8	0.8248 (2)	0.58509 (18)	0.66416 (12)	0.0162 (3)
H8	0.8507	0.5406	0.7389	0.019*
C9	0.77565 (19)	0.47639 (17)	0.55899 (11)	0.0126 (3)
C10	0.6231 (2)	0.11432 (17)	0.25339 (11)	0.0161 (3)
H10A	0.4972	0.0138	0.2639	0.019*
H10B	0.7280	0.0682	0.2602	0.019*
C11	0.5989 (2)	0.17984 (17)	0.12982 (11)	0.0144 (3)
C12	0.4136 (2)	0.16677 (17)	0.07491 (11)	0.0146 (3)
H12	0.3011	0.1147	0.1140	0.018*
C13	0.39414 (19)	0.23009 (17)	-0.03706 (12)	0.0143 (3)
C14	0.5553 (2)	0.30743 (18)	-0.09635 (12)	0.0155 (3)
H14	0.5401	0.3518	-0.1726	0.019*
C15	0.7391 (2)	0.31864 (17)	-0.04193 (12)	0.0165 (3)
H15	0.8510	0.3704	-0.0815	0.020*
C16	0.7613 (2)	0.25498 (17)	0.06999 (12)	0.0157 (3)
H16	0.8880	0.2627	0.1061	0.019*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.01563 (7)	0.02239 (8)	0.02281 (8)	0.00845 (6)	0.00152 (5)	0.00243 (6)
O1	0.0220 (5)	0.0115 (5)	0.0106 (5)	0.0075 (4)	0.0015 (4)	0.0015 (4)
O2	0.0273 (6)	0.0183 (5)	0.0130 (5)	0.0098 (5)	0.0022 (4)	0.0048 (4)
C1	0.0134 (6)	0.0152 (7)	0.0135 (7)	0.0051 (6)	0.0033 (5)	0.0009 (5)
C2	0.0130 (6)	0.0147 (6)	0.0106 (6)	0.0051 (5)	0.0027 (5)	0.0033 (5)
C3	0.0162 (7)	0.0141 (7)	0.0107 (6)	0.0065 (6)	0.0016 (5)	0.0022 (5)
C4	0.0109 (6)	0.0123 (6)	0.0145 (7)	0.0046 (5)	0.0033 (5)	0.0014 (5)
C5	0.0191 (7)	0.0163 (7)	0.0170 (7)	0.0087 (6)	0.0031 (5)	0.0034 (6)
C6	0.0188 (7)	0.0124 (7)	0.0246 (8)	0.0069 (6)	0.0053 (6)	-0.0005 (6)
C7	0.0181 (7)	0.0183 (7)	0.0183 (7)	0.0062 (6)	0.0045 (6)	-0.0049 (6)
C8	0.0164 (7)	0.0187 (7)	0.0123 (7)	0.0054 (6)	0.0039 (5)	0.0009 (5)

C9	0.0111 (6)	0.0130 (6)	0.0138 (7)	0.0045 (5)	0.0034 (5)	0.0013 (5)
C10	0.0218 (7)	0.0117 (6)	0.0141 (7)	0.0060 (6)	0.0028 (5)	0.0010 (5)
C11	0.0208 (7)	0.0089 (6)	0.0124 (7)	0.0053 (6)	0.0014 (5)	-0.0030 (5)
C12	0.0173 (7)	0.0119 (6)	0.0140 (7)	0.0046 (6)	0.0052 (5)	-0.0014 (5)
C13	0.0147 (6)	0.0121 (6)	0.0155 (7)	0.0058 (5)	0.0001 (5)	-0.0032 (5)
C14	0.0196 (7)	0.0137 (7)	0.0131 (7)	0.0068 (6)	0.0021 (5)	0.0004 (5)
C15	0.0170 (7)	0.0149 (7)	0.0170 (7)	0.0051 (6)	0.0055 (5)	0.0019 (6)
C16	0.0159 (7)	0.0145 (7)	0.0154 (7)	0.0057 (6)	0.0002 (5)	-0.0012 (5)

Geometric parameters (Å, °)

Br—C13	1.9001 (13)	C11—C16	1.3947 (19)
O1—C1	1.3796 (15)	C12—C13	1.3854 (18)
O1—C2	1.3859 (15)	C13—C14	1.3859 (19)
O2—C1	1.2073 (15)	C14—C15	1.3842 (18)
C1—C9	1.4618 (18)	C15—C16	1.3887 (18)
C2—C3	1.3256 (18)	C3—H3	0.9500
C2—C10	1.4996 (18)	C5—H5	0.9500
C3—C4	1.4426 (18)	C6—H6	0.9500
C4—C9	1.4035 (18)	C7—H7	0.9500
C4—C5	1.4056 (18)	C8—H8	0.9500
C5—C6	1.3787 (19)	C10—H10A	0.9900
C6—C7	1.392 (2)	C10—H10B	0.9900
C7—C8	1.3759 (19)	C12—H12	0.9500
C8—C9	1.3994 (18)	C14—H14	0.9500
C10—C11	1.5166 (18)	C15—H15	0.9500
C11—C12	1.3911 (18)	C16—H16	0.9500
C1—O1—C2	122.56 (10)	C15—C14—C13	118.52 (13)
O2—C1—O1	117.22 (12)	C14—C15—C16	120.63 (13)
O2—C1—C9	126.19 (12)	C15—C16—C11	120.46 (13)
O1—C1—C9	116.59 (11)	C2—C3—H3	119.7
C3—C2—O1	121.65 (12)	C4—C3—H3	119.7
C3—C2—C10	128.78 (12)	C6—C5—H5	119.9
O1—C2—C10	109.56 (11)	C4—C5—H5	119.9
C2—C3—C4	120.62 (12)	C5—C6—H6	119.6
C9—C4—C5	118.50 (12)	C7—C6—H6	119.6
C9—C4—C3	118.22 (12)	C8—C7—H7	119.9
C5—C4—C3	123.28 (12)	C6—C7—H7	119.9
C6—C5—C4	120.19 (13)	C7—C8—H8	120.1
C5—C6—C7	120.77 (13)	C9—C8—H8	120.1
C8—C7—C6	120.14 (13)	C2—C10—H10A	109.0
C7—C8—C9	119.74 (13)	C11—C10—H10A	109.0
C8—C9—C4	120.65 (12)	C2—C10—H10B	109.0
C8—C9—C1	119.01 (12)	C11—C10—H10B	109.0
C4—C9—C1	120.34 (12)	H10A—C10—H10B	107.8
C2—C10—C11	112.98 (11)	C13—C12—H12	120.2
C12—C11—C16	119.06 (12)	C11—C12—H12	120.2

C12—C11—C10	120.17 (12)	C15—C14—H14	120.7
C16—C11—C10	120.77 (12)	C13—C14—H14	120.7
C13—C12—C11	119.64 (12)	C14—C15—H15	119.7
C12—C13—C14	121.67 (13)	C16—C15—H15	119.7
C12—C13—Br	119.50 (10)	C15—C16—H16	119.8
C14—C13—Br	118.81 (10)	C11—C16—H16	119.8
C2—O1—C1—O2	-179.57 (11)	O2—C1—C9—C8	-0.1 (2)
C2—O1—C1—C9	0.95 (17)	O1—C1—C9—C8	179.34 (11)
C1—O1—C2—C3	-0.72 (19)	O2—C1—C9—C4	-179.30 (13)
C1—O1—C2—C10	-179.69 (11)	O1—C1—C9—C4	0.14 (18)
O1—C2—C3—C4	-0.6 (2)	C3—C2—C10—C11	5.0 (2)
C10—C2—C3—C4	178.12 (13)	O1—C2—C10—C11	-176.15 (11)
C2—C3—C4—C9	1.66 (19)	C2—C10—C11—C12	-90.92 (15)
C2—C3—C4—C5	-178.28 (13)	C2—C10—C11—C16	88.60 (15)
C9—C4—C5—C6	-0.22 (19)	C16—C11—C12—C13	-0.72 (19)
C3—C4—C5—C6	179.72 (13)	C10—C11—C12—C13	178.81 (11)
C4—C5—C6—C7	0.9 (2)	C11—C12—C13—C14	-0.27 (19)
C5—C6—C7—C8	-0.6 (2)	C11—C12—C13—Br	178.14 (9)
C6—C7—C8—C9	-0.2 (2)	C12—C13—C14—C15	0.9 (2)
C7—C8—C9—C4	0.9 (2)	Br—C13—C14—C15	-177.54 (10)
C7—C8—C9—C1	-178.34 (13)	C13—C14—C15—C16	-0.50 (19)
C5—C4—C9—C8	-0.63 (19)	C14—C15—C16—C11	-0.5 (2)
C3—C4—C9—C8	179.43 (12)	C12—C11—C16—C15	1.09 (19)
C5—C4—C9—C1	178.56 (12)	C10—C11—C16—C15	-178.44 (12)
C3—C4—C9—C1	-1.38 (18)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...O1 ⁱ	0.95	2.58	3.4666 (16)	155
C10—H10 <i>A</i> ...O2 ⁱⁱ	0.99	2.50	3.4685 (17)	166

Symmetry codes: (i) *x*, *y*+1, *z*; (ii) -*x*+1, -*y*, -*z*+1.