

(5Z,7Z)-6,8-Dimethyl-9H-tetrazolo-[1,5-b][1,2,4]triazepine

Chun-Lin He, Zhi-Ming Du,* Zheng-Qiang Tang, Xiao-Min Cong and Ling-Qiao Meng

State Key Laboratory of Explosion Science and Technology, Beijing Institute of Technology, Beijing 100081, People's Republic of China
Correspondence e-mail: duzhiming430@sohu.com


Received 12 October 2009; accepted 23 October 2009

Key indicators: single-crystal X-ray study; $T = 93\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.036; wR factor = 0.082; data-to-parameter ratio = 13.8.

The molecule of the title compound, $\text{C}_6\text{H}_8\text{N}_6$, is approximately planar, with a maximum deviation from planarity of $0.099(1)\text{ \AA}$. In the crystal, molecules are linked to each other via pairs of $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonding, forming inversion dimers. The crystal structure is further stabilized by $\pi-\pi$ stacking interactions, with a centroid–centroid distance of $3.419(1)\text{ \AA}$.

Related literature

For the preparation of the title compound, see: Gaponnik & Karavai (1984). For applications of fused tetrazole ring compounds, see: Taha (2007); Zbigniew *et al.* (2007); Galvez-Ruiz *et al.* (2005); Klapötke & Sabaté (2008). For related structures, see: Taha (2005); He *et al.* (2009a,b).



Experimental

Crystal data

$\text{C}_6\text{H}_8\text{N}_6$
 $M_r = 164.18$
Monoclinic, $P2_1/c$
 $a = 3.9184(8)\text{ \AA}$

$b = 13.584(3)\text{ \AA}$
 $c = 13.767(3)\text{ \AA}$
 $\beta = 96.274(3)^\circ$
 $V = 728.4(3)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$

$T = 93\text{ K}$
 $0.47 \times 0.33 \times 0.13\text{ mm}$

Data collection

Rigaku Saturn 724+ diffractometer
Absorption correction: none
5029 measured reflections

1647 independent reflections
1445 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.082$
 $S = 1.00$
1647 reflections
119 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.26\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.22\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N5—H5N \cdots N4 ⁱ	0.920 (19)	1.999 (19)	2.9156 (17)	173.5 (15)

Symmetry code: (i) $-x + 1, -y, -z + 1$.

Data collection: *CrystalClear* (Rigaku, 2008); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported financially by the State Key Laboratory of Explosion Science and Technology of the Beijing Institute of Technology, China (No. ZDKT08-01).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2354).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Galvez-Ruiz, J. C., Holl, G., Karaghiosoff, K., Klapötke, T. M., Lohnwitz, K., Mayer, P., Noth, H., Polborn, K., Rohrbogner, C. J., Suter, M. & Weigand, J. J. (2005). *Inorg. Chem.* **44**, 4237–4253.
- Gaponnik, P. N. & Karavai, V. P. (1984). *Khim. Geterotsikl. Soedin.* **12**, 1683–1686.
- He, C.-L., Du, Z.-M., Tang, Z.-Q., Cong, X.-M. & Meng, L.-Q. (2009a). *Acta Cryst. E65*, o1760.
- He, C.-L., Du, Z.-M., Tang, Z.-Q., Cong, X.-M. & Meng, L.-Q. (2009b). *Acta Cryst. E65*, o1902.
- Klapötke, T. M. & Sabaté, C. M. (2008). *Chem. Mater.* **20**, 1750–1763.
- Rigaku (2008). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Taha, M. A. M. (2005). *J. Indian Chem. Soc.* **82**, 172–174.
- Taha, M. A. M. (2007). *Monatsh. Chem.* **138**, 505–509.
- Zbigniew, K., Mariusz, M. & Andrzej, R. (2007). *J. Mol. Struct.* **829**, 22–28.

supporting information

Acta Cryst. (2009). E65, o2894 [https://doi.org/10.1107/S160053680904392X]

(5Z,7Z)-6,8-Dimethyl-9H-tetrazolo[1,5-*b*][1,2,4]triazepine

Chun-Lin He, Zhi-Ming Du, Zheng-Qiang Tang, Xiao-Min Cong and Ling-Qiao Meng

S1. Comment

Compounds based on the tetrazole ring have generated much interest (Taha 2005). On the one hand, fused tetrazole derivatives play an important role in many biological activities (Taha 2007; Zbigniew *et al.*, 2007); on the other hand, nitrogen-rich compounds, in particular those containing the tetrazole ring, have great potential for energetic applications (Klapötke & Sabaté 2008); Galvez-Ruiz *et al.* 2005). The title compound was first prepared by Gaponnik & Karavai (1984), but its crystal structure has hitherto not been reported. Here, we present the crystal structure of the title compound.

The molecular structure is shown in Fig. 1. The molecule of the title compound assumes an approximately planar structure. The maximum deviation from planarity is 0.099 (1) Å for atom C3. Bond distances and angles are similar to the corresponding distances and angles reported for related compounds (He *et al.* 2009a; He *et al.* 2009b).

In the crystal structure, molecules are linked to each other *via* N—H···N hydrogen bonding (Table 1), forming a dimer structure. Intermolecular π – π interactions with a centroid···centroid distance of 3.419 (1) Å (Table 2), further help to stabilize the crystal structure. The crystal packing of the title compound is shown in Fig. 2, viewed down the *a* axis.

S2. Experimental

The title compound was obtained according to the literature method (Gaponnik & Karavai, 1984). The purity of the compound was checked by determining its melting point, 477–479 K. Single crystals suitable for X-ray crystal structure determination were obtained by slow evaporation of an acetone solution at room temperature over two days.

S3. Refinement

The H atom directly attached to the triazepine ring and that bonded to N were located in difference Fourier maps and refined freely. Methyl H atoms were placed in calculated positions, with C—H = 0.98 Å and refined as riding; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.





Figure 1

The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as spheres of arbitrary radius.

**Figure 2**

The packing of the title compound, viewed along the a -axis, showing the formation of dimers *via* N—H···N hydrogen bonds (dashed lines).

(5 Z ,7 Z)-6,8-Dimethyl-9*H*-tetrazolo[1,5-*b*][1,2,4]triazepine

Crystal data

$C_6H_8N_6$
 $M_r = 164.18$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 3.9184 (8) \text{ \AA}$
 $b = 13.584 (3) \text{ \AA}$
 $c = 13.767 (3) \text{ \AA}$
 $\beta = 96.274 (3)^\circ$
 $V = 728.4 (3) \text{ \AA}^3$
 $Z = 4$

$F(000) = 344$
 $D_x = 1.497 \text{ Mg m}^{-3}$
Melting point: 477 K
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 2263 reflections
 $\theta = 3.3\text{--}27.5^\circ$
 $\mu = 0.11 \text{ mm}^{-1}$
 $T = 93 \text{ K}$
Chunk, red
 $0.47 \times 0.33 \times 0.13 \text{ mm}$

Data collection

Rigaku Saturn 724+
diffractometer
Radiation source: Rotating Anode
Graphite monochromator
Detector resolution: 28.5714 pixels mm⁻¹
multi-scan
5029 measured reflections

1647 independent reflections
1445 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.3^\circ$
 $h = -4 \rightarrow 5$
 $k = -17 \rightarrow 17$
 $l = -11 \rightarrow 17$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.082$
 $S = 1.00$
1647 reflections
119 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.031P)^2 + 0.36P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.26 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.22 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.4467 (3)	0.24707 (7)	0.45448 (7)	0.0165 (2)
N2	0.5870 (3)	0.24983 (8)	0.36816 (7)	0.0208 (2)
N3	0.6547 (3)	0.16030 (8)	0.34635 (8)	0.0223 (3)
N4	0.5650 (3)	0.09674 (8)	0.41698 (7)	0.0195 (2)
N5	0.3238 (3)	0.11503 (8)	0.56546 (8)	0.0207 (3)
N6	0.3868 (3)	0.34046 (7)	0.49433 (7)	0.0176 (2)
C1	0.4389 (3)	0.15238 (9)	0.48318 (8)	0.0163 (3)
C2	0.1660 (3)	0.16746 (9)	0.63408 (8)	0.0165 (3)
C3	0.1175 (3)	0.26544 (9)	0.63462 (9)	0.0180 (3)
C4	0.2395 (3)	0.34313 (9)	0.57359 (9)	0.0165 (3)
C5	0.0598 (3)	0.10241 (9)	0.71313 (9)	0.0202 (3)
H5A	0.2595	0.0819	0.7548	0.024*
H5B	-0.0569	0.0456	0.6844	0.024*
H5C	-0.0914	0.1380	0.7509	0.024*
C6	0.1945 (3)	0.44529 (9)	0.61191 (9)	0.0206 (3)
H6A	0.2920	0.4920	0.5705	0.025*
H6B	0.3084	0.4503	0.6771	0.025*

H6C	-0.0457	0.4588	0.6126	0.025*
H3	0.006 (4)	0.2905 (11)	0.6876 (11)	0.025 (4)*
H5N	0.342 (4)	0.0478 (14)	0.5724 (12)	0.036 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0214 (5)	0.0138 (5)	0.0148 (5)	0.0006 (4)	0.0043 (4)	0.0002 (4)
N2	0.0285 (6)	0.0182 (5)	0.0167 (5)	0.0010 (4)	0.0074 (4)	0.0004 (4)
N3	0.0309 (6)	0.0180 (5)	0.0191 (5)	0.0013 (5)	0.0078 (4)	0.0014 (4)
N4	0.0261 (6)	0.0167 (5)	0.0166 (5)	0.0007 (4)	0.0062 (4)	0.0005 (4)
N5	0.0315 (6)	0.0130 (5)	0.0193 (5)	0.0016 (4)	0.0106 (5)	0.0019 (4)
N6	0.0230 (5)	0.0110 (5)	0.0189 (5)	0.0009 (4)	0.0030 (4)	-0.0018 (4)
C1	0.0177 (6)	0.0151 (6)	0.0161 (6)	0.0000 (5)	0.0013 (4)	0.0005 (4)
C2	0.0174 (6)	0.0168 (6)	0.0153 (5)	-0.0009 (5)	0.0021 (4)	-0.0007 (4)
C3	0.0196 (6)	0.0181 (6)	0.0168 (6)	0.0000 (5)	0.0050 (5)	-0.0007 (5)
C4	0.0163 (6)	0.0147 (6)	0.0181 (6)	0.0004 (4)	-0.0002 (5)	-0.0002 (4)
C5	0.0240 (6)	0.0174 (6)	0.0202 (6)	-0.0003 (5)	0.0072 (5)	0.0024 (5)
C6	0.0246 (6)	0.0155 (6)	0.0224 (6)	0.0011 (5)	0.0055 (5)	-0.0020 (5)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.3469 (15)	C2—C3	1.3446 (17)
N1—N2	1.3635 (14)	C2—C5	1.4958 (16)
N1—N6	1.4119 (14)	C3—C4	1.4615 (17)
N2—N3	1.2874 (15)	C3—H3	0.953 (15)
N3—N4	1.3745 (14)	C4—C6	1.5018 (16)
N4—C1	1.3207 (15)	C5—H5A	0.9600
N5—C1	1.3623 (15)	C5—H5B	0.9600
N5—C2	1.3820 (15)	C5—H5C	0.9600
N5—H5N	0.920 (18)	C6—H6A	0.9600
N6—C4	1.2897 (16)	C6—H6B	0.9600
C1—N1—N2	107.83 (10)	C2—C3—H3	115.8 (9)
C1—N1—N6	137.23 (10)	C4—C3—H3	112.8 (9)
N2—N1—N6	114.45 (9)	N6—C4—C3	132.13 (11)
N3—N2—N1	106.88 (10)	N6—C4—C6	113.82 (10)
N2—N3—N4	110.69 (10)	C3—C4—C6	114.02 (11)
C1—N4—N3	105.82 (10)	C2—C5—H5A	109.5
C1—N5—C2	126.17 (11)	C2—C5—H5B	109.5
C1—N5—H5N	115.3 (10)	H5A—C5—H5B	109.5
C2—N5—H5N	118.4 (10)	C2—C5—H5C	109.5
C4—N6—N1	117.58 (10)	H5A—C5—H5C	109.5
N4—C1—N1	108.78 (10)	H5B—C5—H5C	109.5
N4—C1—N5	122.92 (11)	C4—C6—H6A	109.5
N1—C1—N5	128.29 (11)	C4—C6—H6B	109.5
C3—C2—N5	126.01 (11)	H6A—C6—H6B	109.5
C3—C2—C5	122.01 (11)	C4—C6—H6C	109.5

N5—C2—C5	111.95 (11)	H6A—C6—H6C	109.5
C2—C3—C4	131.03 (12)	H6B—C6—H6C	109.5
C1—N1—N2—N3	0.80 (13)	N6—N1—C1—N5	8.2 (2)
N6—N1—N2—N3	174.17 (10)	C2—N5—C1—N4	-174.90 (12)
N1—N2—N3—N4	-0.47 (14)	C2—N5—C1—N1	5.0 (2)
N2—N3—N4—C1	-0.04 (14)	C1—N5—C2—C3	-4.3 (2)
C1—N1—N6—C4	-12.5 (2)	C1—N5—C2—C5	177.67 (12)
N2—N1—N6—C4	176.81 (10)	N5—C2—C3—C4	-7.1 (2)
N3—N4—C1—N1	0.55 (13)	C5—C2—C3—C4	170.75 (12)
N3—N4—C1—N5	-179.54 (11)	N1—N6—C4—C3	0.64 (19)
N2—N1—C1—N4	-0.84 (14)	N1—N6—C4—C6	178.52 (10)
N6—N1—C1—N4	-171.93 (13)	C2—C3—C4—N6	10.9 (2)
N2—N1—C1—N5	179.24 (12)	C2—C3—C4—C6	-167.02 (13)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N5—H5N···N4 ⁱ	0.920 (19)	1.999 (19)	2.9156 (17)	173.5 (15)

Symmetry code: (i) $-x+1, -y, -z+1$.