

Hydronium 4-oxo-1,4-dihydropyridine-3-sulfonate dihydrate

Zhi-Biao Zhu,^a Shan Gao^a and Seik Weng Ng^{b*}

^aCollege of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 5 October 2009; accepted 6 October 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.049; wR factor = 0.170; data-to-parameter ratio = 14.7.

2-Hydroxypyridine when treated with concentrated sulfuric acid is sulfonated at the 3-position to yield the title hydrated salt, $\text{H}_3\text{O}^+\cdot\text{C}_5\text{H}_4\text{NO}_3\text{S}^-\cdot2\text{H}_2\text{O}$. In the crystal structure, the cations, anions and uncoordinated water molecules are linked by extensive $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into a three-dimensional network. The crystal studied is a non-merohedral twin with a twin component of 36%.

Related literature

For the treatment of non-merohedral twins, see: Spek (2003). For the cobalt salt of the anion, see: Zhu *et al.* (2007).



Experimental

Crystal data

$\text{H}_3\text{O}^+\cdot\text{C}_5\text{H}_4\text{NO}_3\text{S}^-\cdot2\text{H}_2\text{O}$
 $M_r = 229.21$

Triclinic, $P\bar{1}$
 $a = 6.6610 (8)\text{ \AA}$
 $b = 7.4179 (10)\text{ \AA}$
 $c = 10.0514 (11)\text{ \AA}$
 $\alpha = 90.922 (4)^\circ$
 $\beta = 94.543 (3)^\circ$

$\gamma = 96.226 (4)^\circ$
 $V = 492.01 (10)\text{ \AA}^3$
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.34\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.25 \times 0.22 \times 0.20\text{ mm}$

Data collection

Rigaku R-AXIS RAPID IP diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.856$, $T_{\max} = 0.932$

4711 measured reflections
2234 independent reflections
1719 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.170$
 $S = 1.06$
2234 reflections
152 parameters
13 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.65\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.59\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1w—H11···O1	0.85 (1)	1.90 (1)	2.745 (4)	175 (5)
O1w—H12···O2 ⁱ	0.85 (1)	2.50 (4)	2.983 (4)	116 (3)
O2w—H21···O1w	0.85 (1)	1.89 (1)	2.741 (4)	175 (6)
O2w—H22···O4 ⁱⁱ	0.85 (1)	2.07 (2)	2.910 (4)	168 (6)
O3w—H31···O2 ⁱ	0.86 (1)	2.12 (2)	2.951 (4)	165 (5)
O3w—H32···O2 ⁱⁱ	0.85 (1)	2.07 (2)	2.898 (4)	162 (5)
O3w—H33···O3 ⁱⁱⁱ	0.85 (1)	2.18 (2)	2.936 (4)	147 (4)
N1—H1···O3 ^{iv}	0.86 (1)	2.05 (2)	2.857 (4)	156 (5)

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x, y, z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x + 1, y, z$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Natural Science Foundation of Heilongjiang Province (No. B200501), Heilongjiang University, China, and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2628).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Westrip, S. P. (2009). *publCIF*. In preparation.
- Zhu, Z.-B., Gao, S., Huo, L.-H. & Zhao, H. (2007). *Acta Cryst. E* **63**, m3126–m3127.

supporting information

Acta Cryst. (2009). E65, o2687 [https://doi.org/10.1107/S1600536809040641]

Hydronium 4-oxo-1,4-dihydropyridine-3-sulfonate dihydrate

Zhi-Biao Zhu, Shan Gao and Seik Weng Ng

S1. Experimental

A 1:3 molar mixture of 4-hydroxypyridine and oleum (20%) was heated at 483 K for 3 h. Barium carbonate was added to the cool mixture until no more carbon dioxide evolved. Colorless crystals were isolated from the filtered solution after several days. CH&N elemental analysis. Calc. for $C_5H_{10}NO_7S$: C 26.20, H 4.84, N 6.11%; found: C 26.36, H 4.48, N 6.11%.

S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. The water, oxonium and amino H-atoms were located in a difference Fourier map and refined with a distance restraint of O—H = N—H 0.85 ± 0.01 Å. The temperature factors of some of the atoms could not be refined; as such, all temperature factors were tied.

The crystal is a non-merohedral twin as shown by *PLATON* (Spek, 2003).




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $[H_3O][C_5H_4NO_3S] \cdot 2H_2O$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Hydronium 4-oxo-1,4-dihydropyridine-3-sulfonate dihydrate*Crystal data*

$M_r = 229.21$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.6610 (8)$ Å

$b = 7.4179 (10)$ Å

$c = 10.0514 (11)$ Å

$\alpha = 90.922 (4)^\circ$

$\beta = 94.543 (3)^\circ$

$\gamma = 96.226 (4)^\circ$

$V = 492.01 (10)$ Å³

$Z = 2$

$F(000) = 240$

$D_x = 1.547 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3881 reflections

$\theta = 3.1\text{--}27.5^\circ$

$\mu = 0.34 \text{ mm}^{-1}$

$T = 293$ K

Block, colorless

$0.25 \times 0.22 \times 0.20$ mm

Data collection

Rigaku R-AXIS RAPID IP
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.856$, $T_{\max} = 0.932$

4711 measured reflections

2234 independent reflections

1719 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -8 \rightarrow 8$

$k = -9 \rightarrow 9$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.049$

$wR(F^2) = 0.170$

$S = 1.06$

2234 reflections

152 parameters

13 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0803P)^2 + 0.6366P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.65 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.59 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.68686 (13)	0.38798 (11)	0.16672 (8)	0.0280 (2)
O1	0.6502 (4)	0.2383 (4)	0.4423 (2)	0.0363 (6)
O2	0.5597 (4)	0.2165 (3)	0.1408 (2)	0.0365 (6)
O3	0.5787 (4)	0.5226 (3)	0.2300 (2)	0.0350 (6)
O4	0.7845 (4)	0.4562 (4)	0.0509 (2)	0.0402 (6)
O1W	0.5750 (5)	0.0846 (4)	0.6825 (3)	0.0479 (7)

H11	0.605 (9)	0.135 (5)	0.610 (3)	0.072*
H12	0.523 (8)	-0.024 (3)	0.664 (4)	0.072*
O2W	0.8700 (5)	0.1491 (5)	0.8890 (3)	0.0582 (8)
H21	0.783 (7)	0.132 (7)	0.822 (4)	0.087*
H22	0.855 (9)	0.249 (5)	0.928 (5)	0.087*
O3W	0.2908 (5)	0.1387 (4)	0.9010 (3)	0.0540 (8)
H31	0.310 (8)	0.028 (2)	0.887 (4)	0.081*
H32	0.350 (7)	0.176 (5)	0.976 (2)	0.081*
H33	0.338 (7)	0.204 (5)	0.839 (3)	0.081*
N1	1.2314 (5)	0.3608 (5)	0.3510 (3)	0.0403 (7)
H1	1.351 (3)	0.385 (7)	0.327 (5)	0.060*
C1	1.1944 (6)	0.2958 (6)	0.4722 (4)	0.0405 (9)
H1A	1.3022	0.2809	0.5342	0.049*
C2	1.0013 (5)	0.2520 (5)	0.5047 (4)	0.0346 (8)
H2	0.9795	0.2068	0.5887	0.042*
C3	0.8321 (5)	0.2734 (4)	0.4136 (3)	0.0278 (6)
C4	0.8828 (5)	0.3435 (4)	0.2858 (3)	0.0264 (7)
C5	1.0802 (6)	0.3834 (5)	0.2586 (4)	0.0334 (7)
H5	1.1100	0.4267	0.1752	0.040*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0305 (4)	0.0281 (4)	0.0249 (4)	0.0000 (3)	0.0025 (3)	0.0033 (3)
O1	0.0272 (12)	0.0504 (15)	0.0308 (12)	-0.0019 (11)	0.0066 (10)	0.0093 (11)
O2	0.0385 (13)	0.0313 (13)	0.0367 (13)	-0.0032 (11)	-0.0062 (11)	0.0012 (10)
O3	0.0367 (13)	0.0310 (12)	0.0389 (13)	0.0080 (11)	0.0072 (11)	0.0049 (10)
O4	0.0465 (16)	0.0461 (15)	0.0281 (12)	-0.0003 (13)	0.0082 (11)	0.0093 (11)
O1W	0.0621 (19)	0.0438 (16)	0.0346 (14)	-0.0083 (15)	0.0032 (14)	0.0057 (11)
O2W	0.0456 (17)	0.066 (2)	0.062 (2)	0.0103 (17)	-0.0055 (16)	-0.0101 (16)
O3W	0.063 (2)	0.0457 (17)	0.0525 (18)	0.0037 (15)	0.0025 (16)	0.0059 (13)
N1	0.0244 (14)	0.0515 (19)	0.0452 (18)	0.0016 (14)	0.0100 (14)	-0.0040 (14)
C1	0.0287 (18)	0.050 (2)	0.043 (2)	0.0081 (16)	-0.0009 (16)	-0.0059 (17)
C2	0.0309 (18)	0.043 (2)	0.0300 (16)	0.0057 (15)	0.0013 (14)	0.0019 (14)
C3	0.0269 (15)	0.0281 (15)	0.0274 (15)	-0.0018 (13)	0.0031 (13)	0.0003 (12)
C4	0.0233 (14)	0.0282 (16)	0.0275 (15)	0.0017 (12)	0.0030 (12)	-0.0005 (12)
C5	0.0325 (17)	0.0332 (17)	0.0347 (17)	0.0002 (15)	0.0087 (15)	-0.0017 (13)

Geometric parameters (\AA , $^\circ$)

S1—O4	1.449 (2)	O3W—H33	0.854 (10)
S1—O2	1.457 (2)	N1—C5	1.341 (5)
S1—O3	1.459 (3)	N1—C1	1.348 (5)
S1—C4	1.762 (3)	N1—H1	0.855 (10)
O1—C3	1.268 (4)	C1—C2	1.358 (5)
O1W—H11	0.850 (10)	C1—H1A	0.9300
O1W—H12	0.854 (10)	C2—C3	1.419 (5)
O2W—H21	0.854 (10)	C2—H2	0.9300

O2W—H22	0.851 (10)	C3—C4	1.445 (4)
O3W—H31	0.855 (10)	C4—C5	1.367 (5)
O3W—H32	0.853 (10)	C5—H5	0.9300
O4—S1—O2	113.46 (15)	N1—C1—H1A	119.7
O4—S1—O3	112.83 (16)	C2—C1—H1A	119.7
O2—S1—O3	111.76 (16)	C1—C2—C3	121.7 (3)
O4—S1—C4	106.31 (16)	C1—C2—H2	119.2
O2—S1—C4	106.22 (15)	C3—C2—H2	119.2
O3—S1—C4	105.54 (15)	O1—C3—C2	123.1 (3)
H11—O1W—H12	108.9 (17)	O1—C3—C4	122.2 (3)
H21—O2W—H22	108.9 (17)	C2—C3—C4	114.7 (3)
H31—O3W—H32	109.7 (17)	C5—C4—C3	120.9 (3)
H31—O3W—H33	109.4 (17)	C5—C4—S1	119.7 (3)
H32—O3W—H33	109.9 (17)	C3—C4—S1	119.3 (2)
C5—N1—C1	121.5 (3)	N1—C5—C4	120.5 (3)
C5—N1—H1	116 (3)	N1—C5—H5	119.7
C1—N1—H1	122 (3)	C4—C5—H5	119.7
N1—C1—C2	120.7 (4)		
C5—N1—C1—C2	-0.4 (6)	O2—S1—C4—C5	124.1 (3)
N1—C1—C2—C3	-0.3 (6)	O3—S1—C4—C5	-117.1 (3)
C1—C2—C3—O1	-178.0 (4)	O4—S1—C4—C3	179.9 (3)
C1—C2—C3—C4	0.4 (5)	O2—S1—C4—C3	-58.9 (3)
O1—C3—C4—C5	178.7 (3)	O3—S1—C4—C3	59.8 (3)
C2—C3—C4—C5	0.3 (5)	C1—N1—C5—C4	1.0 (6)
O1—C3—C4—S1	1.8 (5)	C3—C4—C5—N1	-1.0 (5)
C2—C3—C4—S1	-176.6 (2)	S1—C4—C5—N1	175.9 (3)
O4—S1—C4—C5	2.9 (3)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1w—H11···O1	0.85 (1)	1.90 (1)	2.745 (4)	175 (5)
O1w—H12···O2 ⁱ	0.85 (1)	2.50 (4)	2.983 (4)	116 (3)
O2w—H21···O1w	0.85 (1)	1.89 (1)	2.741 (4)	175 (6)
O2w—H22···O4 ⁱⁱ	0.85 (1)	2.07 (2)	2.910 (4)	168 (6)
O3w—H31···O2 ⁱ	0.86 (1)	2.12 (2)	2.951 (4)	165 (5)
O3w—H32···O2 ⁱⁱ	0.85 (1)	2.07 (2)	2.898 (4)	162 (5)
O3w—H33···O3 ⁱⁱⁱ	0.85 (1)	2.18 (2)	2.936 (4)	147 (4)
N1—H1···O3 ^{iv}	0.86 (1)	2.05 (2)	2.857 (4)	156 (5)

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x, y, z+1$; (iii) $-x+1, -y+1, -z+1$; (iv) $x+1, y, z$.