

Tetraaquabis(pyridine- κ N)cobalt(II) bis[4-amino-*N*-(6-chloropyridazin-3-yl)benzenesulfonamidate]

Nan Li, Hong-Li Zou, Xiao-Yan Song, Yan-Cheng Liu and Zhen-Feng Chen*

Key Laboratory for the Chemistry and Molecular Engineering of Medicinal Resources (Ministry Education of China), School of Chemistry & Chemical Engineering, Guangxi Normal University, Guilin 541004, People's Republic of China
Correspondence e-mail: chenzfgxnu@yahoo.com


Received 16 November 2009; accepted 19 November 2009

Key indicators: single-crystal X-ray study; $T = 193$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.054; wR factor = 0.108; data-to-parameter ratio = 16.4.

The structure of the title compound, $[\text{Co}(\text{C}_5\text{H}_5\text{N})_2(\text{H}_2\text{O})_4]-(\text{C}_{10}\text{H}_8\text{ClN}_4\text{O}_2\text{S})_2$, consists of a discrete tetraaquabis(pyridine- κ N)cobalt(II) cation and two 4-amino-*N*-(6-chloropyridazin-3-yl)benzenesulfonamidate anions. In the cation, the Co^{II} ion sits on an inversion centre and is octahedrally coordinated by two pyridine N atoms and four O atoms. A two-dimensional network parallel to (010) is formed *via* intermolecular O—H...O, O—H...N, N—H...N and N—H...O hydrogen bonds.

Related literature

For the structure of sulfachloropyridazine, see: Tan *et al.* (2005). For a sulfachloropyridazine–metal complex, see: Fogg *et al.* (1995). For an aquapyridine–cobalt(II) complex, see: Clegg *et al.* (2006).



Experimental

Crystal data

$[\text{Co}(\text{C}_5\text{H}_5\text{N})_2(\text{H}_2\text{O})_4]-(\text{C}_{10}\text{H}_8\text{ClN}_4\text{O}_2\text{S})_2$
 $M_r = 856.62$
Monoclinic, $P2_1/c$

$a = 8.5897$ (12) Å
 $b = 25.807$ (3) Å
 $c = 8.5338$ (12) Å
 $\beta = 101.694$ (3)°

$V = 1852.5$ (4) Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.78$ mm⁻¹
 $T = 193$ K
 $0.21 \times 0.15 \times 0.12$ mm

Data collection

Rigaku Mercury CCD diffractometer
Absorption correction: multi-scan (*REQAB*; Jacobson, 1998)
 $T_{\text{min}} = 0.853$, $T_{\text{max}} = 0.912$

20359 measured reflections
4226 independent reflections
3331 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.054$
 $wR(F^2) = 0.108$
 $S = 1.14$
4226 reflections
258 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.66$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3A...N5 ⁱ	0.88	2.52	3.135 (4)	127
N3—H3B...O3 ⁱⁱⁱ	0.88	2.23	3.001 (3)	147
O1—H1A...O3	0.81 (4)	2.07 (4)	2.874 (3)	174 (4)
O1—H1B...N4 ⁱⁱⁱ	0.83 (4)	2.01 (4)	2.838 (3)	178 (3)
O2—H2A...O4	0.82 (3)	1.93 (4)	2.726 (3)	165 (4)
O2—H2B...N2 ^{iv}	0.82 (4)	1.96 (4)	2.768 (3)	170 (4)
O2—H2B...O4 ^{iv}	0.82 (4)	2.62 (4)	3.142 (3)	123 (3)

Symmetry codes: (i) $x + 1, y, z + 1$; (ii) $x, y, z + 1$; (iii) $x + 1, y, z$; (iv) $-x + 1, -y + 1, -z + 1$.

Data collection: *CrystalClear* (Rigaku, 1999); cell refinement: *CrystalClear*; data reduction: *CrystalStructure* (Rigaku/MSC and Rigaku, 2000); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (No. 20861002), the 973 Plan of China (2009CB526503) the Natural Science Foundation of Guangxi, China (Nos. 0991003, 0991012Z) and the Open Foundation of the Key Laboratory for the Chemistry and Molecular Engineering of Medicinal Resources (Ministry of Education of China) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2211).

References

- Clegg, J. K., Hayter, M. J., Jolliffe, K. A. & Lindoy, L. F. (2006). *Acta Cryst.* **E62**, m873–m874.
Fogg, A. G., Yusoff, A. R. H. M. & Ahmad, R. (1995). *Anal. Proc.* **32**, 337–340.
Jacobson, R. (1998). *REQAB*. Private communication to the Rigaku Corporation, Tokyo, Japan.
Rigaku (1999). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MSC and Rigaku (2000). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Tan, Y.-S., Chen, Z.-F., Liang, H. & Zhang, Y. (2005). *Acta Cryst.* **E61**, o1842–o1844.

supporting information

Acta Cryst. (2009). E65, m1666 [doi:10.1107/S1600536809049599]

Tetraaquabis(pyridine- κ N)cobalt(II) bis[4-amino-*N*-(6-chloropyridazin-3-yl)benzenesulfonamidate]

Nan Li, Hong-Li Zou, Xiao-Yan Song, Yan-Cheng Liu and Zhen-Feng Chen

S1. Comment


Sulfachloropyridazine [4-amino-*N*-(6-chloro-3-pyridazinyl)-benzenesulfonamide], is a synthetic sulfanilamide antibacterial drug whose crystal structure is known (Tan *et al.* 2005). However, as far as we are aware, no crystal structure of a metal complex containing sulfachloropyridazine has been published, although the electrochemistry of its copper(I) complex has been described by Fogg *et al.* (1995). The compound consists of a $[\text{Co}(\text{H}_2\text{O})_4(\text{py})_2]^{2+}$ cation and two sulfachloropyridazine anions. Similar to tetraaquabis(pyridine- κ N)cobalt(II) diacetate (Clegg *et al.*, 2006), the coordination geometry for the Co^{II} cation (Fig. 1) is close to an ideal octahedron (N1—Co1—N1^{#1} 180.000 (1)°, O1—Co1—O2 92.37 (9)°, O1—Co1—O2^{#1} 87.63 (9)°, #1: $-x + 2, -y + 1, -z + 1$), with the O atoms of the coordinated water molecules occupying the equatorial positions and with the axial sites occupied by coordinated pyridine ligands. The sulfachloropyridazine is deprotonated at sulfamide N2 to generate anions, whose geometric parameters are comparable to sulfachloropyridazine (Tan *et al.*, 2005). A two-dimensional network is formed *via* intermolecular hydrogen bonds of type O—H \cdots O, O—H \cdots N and N—H \cdots O (Table 1).

S2. Experimental


Samples of sulfachloropyridazine (0.2 mmol) and $\text{Co}(\text{Ac})_2 \cdot 4\text{H}_2\text{O}$ (0.1 mmol) were placed in a thick-walled Pyrex tube (*ca* 20 cm long). After addition of ethanol (2.2 ml), H_2O (0.2 ml) and pyridine (0.1 ml), the tube was frozen with liquid nitrogen, evacuated under vacuum and sealed with a torch. The tube was heated at 80°C for 3 days and then was slowly cooled down to room temperature, and orange-yellow block-shaped crystals were obtained. Yield: 80%.

S3. Refinement

The H atoms bonded to C atoms were positioned geometrically and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ (C—H = 0.95 Å). The H atoms attached to amino N were placed in the calculated positions, with N—H distance of 0.88 Å. The $U_{\text{iso}}(\text{H})$ values were constrained to be $1.2U_{\text{eq}}$ of the carrier atom for amino H atom. H atom attached to O was located in an electron-density difference map and refined isotropically.

**Figure 1**

The molecular structure showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

A packing diagram of title compound, viewing down the [100] direction. Hydrogen bonds are shown as dashed lines.

Tetraaquabis(pyridine- κ N)cobalt(II) bis[4-amino-*N*-(6-chloropyridazin-3-yl)benzenesulfonamidate]*Crystal data*[Co(C₅H₅N)₂(H₂O)₄](C₁₀H₈ClN₄O₂S)₂ $M_r = 856.62$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 8.5897$ (12) Å $b = 25.807$ (3) Å $c = 8.5338$ (12) Å $\beta = 101.694$ (3)° $V = 1852.5$ (4) Å³ $Z = 2$ $F(000) = 882$ $D_x = 1.536$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71070$ Å

Cell parameters from 6394 reflections

 $\theta = 3.1$ – 27.5 ° $\mu = 0.78$ mm⁻¹ $T = 193$ K

Block, orange-yellow

 $0.21 \times 0.15 \times 0.12$ mm*Data collection*

Rigaku Mercury CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 7.31 pixels mm⁻¹ ω scans

Absorption correction: multi-scan

(REQAB; Jacobson, 1998)

 $T_{\min} = 0.853$, $T_{\max} = 0.912$

20359 measured reflections

4226 independent reflections

3331 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.050$ $\theta_{\max} = 27.5$ °, $\theta_{\min} = 3.2$ ° $h = -11 \rightarrow 9$ $k = -32 \rightarrow 33$ $l = -10 \rightarrow 11$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.054$ $wR(F^2) = 0.108$ $S = 1.14$

4226 reflections

258 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0291P)^2 + 1.7941P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.66$ e Å⁻³ $\Delta\rho_{\min} = -0.45$ e Å⁻³*Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	1.0000	0.5000	0.5000	0.02572 (15)
Cl1	0.07781 (10)	0.20506 (3)	0.22874 (12)	0.0489 (2)
S1	0.54834 (8)	0.39936 (3)	0.61366 (9)	0.02737 (18)

O1	0.9392 (3)	0.42160 (8)	0.4313 (3)	0.0348 (5)
O2	0.7795 (2)	0.51495 (9)	0.5400 (3)	0.0337 (5)
O3	0.6620 (2)	0.37878 (8)	0.5233 (2)	0.0330 (5)
O4	0.5622 (2)	0.45443 (7)	0.6462 (2)	0.0331 (5)
N1	0.9177 (3)	0.52499 (9)	0.2536 (3)	0.0315 (6)
N2	0.3692 (3)	0.39029 (9)	0.5297 (3)	0.0299 (6)
N3	0.7205 (3)	0.30830 (10)	1.2615 (3)	0.0394 (6)
H3A	0.7967	0.2850	1.2827	0.047*
H3B	0.6729	0.3190	1.3377	0.047*
N4	0.1620 (3)	0.34401 (9)	0.3915 (3)	0.0309 (6)
N5	0.0921 (3)	0.30070 (9)	0.3205 (3)	0.0323 (6)
C1	0.9858 (4)	0.56332 (13)	0.1875 (4)	0.0405 (8)
H1C	1.0705	0.5817	0.2530	0.049*
C2	0.9403 (4)	0.57775 (15)	0.0292 (4)	0.0479 (9)
H2C	0.9928	0.6054	-0.0125	0.057*
C3	0.8187 (4)	0.55177 (14)	-0.0666 (4)	0.0457 (9)
H3	0.7851	0.5609	-0.1760	0.055*
C4	0.7458 (4)	0.51210 (14)	-0.0018 (4)	0.0469 (9)
H4	0.6609	0.4933	-0.0657	0.056*
C5	0.7977 (4)	0.49995 (12)	0.1575 (4)	0.0378 (7)
H5	0.7463	0.4726	0.2015	0.045*
C6	0.5891 (3)	0.36813 (10)	0.8007 (3)	0.0262 (6)
C7	0.7082 (4)	0.33130 (12)	0.8356 (4)	0.0339 (7)
H7	0.7613	0.3199	0.7545	0.041*
C8	0.7499 (4)	0.31125 (12)	0.9877 (4)	0.0355 (7)
H8	0.8307	0.2856	1.0101	0.043*
C9	0.6757 (3)	0.32790 (11)	1.1094 (3)	0.0299 (6)
C10	0.5524 (4)	0.36397 (12)	1.0716 (4)	0.0349 (7)
H10	0.4976	0.3750	1.1519	0.042*
C11	0.5097 (3)	0.38357 (11)	0.9191 (4)	0.0334 (7)
H11	0.4252	0.4079	0.8946	0.040*
C12	0.3157 (3)	0.34323 (11)	0.4672 (3)	0.0275 (6)
C13	0.1763 (3)	0.25836 (11)	0.3265 (4)	0.0315 (7)
C14	0.3340 (4)	0.25379 (11)	0.4036 (4)	0.0351 (7)
H14	0.3898	0.2219	0.4054	0.042*
C15	0.4049 (3)	0.29672 (11)	0.4761 (4)	0.0328 (7)
H15	0.5124	0.2957	0.5321	0.039*
H1A	0.859 (5)	0.4089 (15)	0.451 (5)	0.060 (13)*
H1B	1.005 (4)	0.3990 (13)	0.422 (4)	0.038 (10)*
H2A	0.717 (4)	0.4932 (12)	0.559 (5)	0.070 (14)*
H2B	0.734 (4)	0.5420 (15)	0.508 (5)	0.058 (12)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0215 (3)	0.0216 (3)	0.0339 (3)	0.0004 (2)	0.0054 (2)	-0.0016 (2)
Cl1	0.0384 (5)	0.0396 (5)	0.0696 (6)	-0.0099 (3)	0.0126 (4)	-0.0209 (4)
S1	0.0224 (4)	0.0263 (4)	0.0332 (4)	-0.0016 (3)	0.0051 (3)	0.0001 (3)

O1	0.0275 (12)	0.0230 (11)	0.0555 (15)	-0.0012 (10)	0.0120 (11)	-0.0052 (10)
O2	0.0246 (11)	0.0258 (12)	0.0520 (14)	0.0042 (9)	0.0105 (10)	0.0054 (10)
O3	0.0261 (11)	0.0372 (11)	0.0375 (12)	-0.0036 (9)	0.0105 (9)	-0.0032 (9)
O4	0.0326 (11)	0.0247 (10)	0.0413 (12)	-0.0052 (8)	0.0061 (10)	0.0010 (9)
N1	0.0284 (13)	0.0287 (13)	0.0373 (14)	0.0042 (10)	0.0063 (11)	-0.0024 (11)
N2	0.0241 (13)	0.0237 (12)	0.0401 (14)	0.0002 (9)	0.0026 (11)	0.0004 (11)
N3	0.0413 (16)	0.0428 (15)	0.0343 (14)	0.0065 (12)	0.0082 (12)	0.0040 (12)
N4	0.0235 (12)	0.0295 (13)	0.0377 (14)	0.0019 (10)	0.0017 (11)	-0.0023 (11)
N5	0.0261 (13)	0.0296 (13)	0.0399 (15)	-0.0022 (10)	0.0034 (11)	-0.0043 (11)
C1	0.0365 (18)	0.0438 (19)	0.0411 (19)	-0.0041 (14)	0.0075 (15)	0.0011 (15)
C2	0.049 (2)	0.055 (2)	0.041 (2)	0.0012 (17)	0.0155 (17)	0.0091 (17)
C3	0.050 (2)	0.056 (2)	0.0301 (17)	0.0174 (18)	0.0056 (16)	-0.0023 (16)
C4	0.046 (2)	0.047 (2)	0.042 (2)	0.0085 (16)	-0.0042 (17)	-0.0120 (16)
C5	0.0353 (17)	0.0331 (16)	0.0422 (18)	-0.0005 (13)	0.0014 (15)	-0.0027 (14)
C6	0.0198 (14)	0.0253 (14)	0.0334 (16)	-0.0004 (11)	0.0049 (12)	0.0000 (12)
C7	0.0295 (16)	0.0374 (17)	0.0360 (17)	0.0054 (13)	0.0093 (14)	-0.0033 (14)
C8	0.0316 (17)	0.0368 (17)	0.0379 (18)	0.0097 (13)	0.0063 (14)	0.0014 (14)
C9	0.0294 (16)	0.0269 (15)	0.0325 (16)	-0.0039 (12)	0.0046 (13)	-0.0014 (13)
C10	0.0341 (17)	0.0353 (17)	0.0390 (17)	0.0016 (13)	0.0159 (14)	-0.0013 (14)
C11	0.0284 (16)	0.0299 (16)	0.0442 (18)	0.0059 (12)	0.0132 (14)	-0.0003 (14)
C12	0.0245 (14)	0.0260 (14)	0.0318 (16)	0.0021 (11)	0.0053 (12)	0.0017 (12)
C13	0.0270 (15)	0.0292 (16)	0.0394 (17)	-0.0045 (12)	0.0095 (13)	-0.0053 (13)
C14	0.0293 (16)	0.0250 (15)	0.0505 (19)	0.0052 (12)	0.0070 (14)	-0.0020 (14)
C15	0.0211 (15)	0.0292 (15)	0.0463 (18)	0.0015 (11)	0.0025 (13)	0.0002 (14)

Geometric parameters (Å, °)

Co1—O2 ⁱ	2.028 (2)	C1—C2	1.379 (5)
Co1—O2	2.028 (2)	C1—H1C	0.9500
Co1—O1	2.142 (2)	C2—C3	1.365 (5)
Co1—O1 ⁱ	2.142 (2)	C2—H2C	0.9500
Co1—N1 ⁱ	2.176 (2)	C3—C4	1.374 (5)
Co1—N1	2.176 (2)	C3—H3	0.9500
C11—C13	1.737 (3)	C4—C5	1.378 (5)
S1—O4	1.448 (2)	C4—H4	0.9500
S1—O3	1.461 (2)	C5—H5	0.9500
S1—N2	1.577 (2)	C6—C7	1.384 (4)
S1—C6	1.759 (3)	C6—C11	1.387 (4)
O1—H1A	0.81 (4)	C7—C8	1.376 (4)
O1—H1B	0.83 (4)	C7—H7	0.9500
O2—H2A	0.82 (3)	C8—C9	1.392 (4)
O2—H2B	0.82 (4)	C8—H8	0.9500
N1—C1	1.331 (4)	C9—C10	1.398 (4)
N1—C5	1.346 (4)	C10—C11	1.375 (4)
N2—C12	1.368 (3)	C10—H10	0.9500
N3—C9	1.374 (4)	C11—H11	0.9500
N3—H3A	0.8800	C12—C15	1.418 (4)
N3—H3B	0.8800	C13—C14	1.386 (4)

N4—C12	1.347 (4)	C14—C15	1.352 (4)
N4—N5	1.352 (3)	C14—H14	0.9500
N5—C13	1.306 (4)	C15—H15	0.9500
O2 ⁱ —Co1—O2	180.0	C1—C2—H2C	120.5
O2 ⁱ —Co1—O1	87.63 (9)	C2—C3—C4	118.8 (3)
O2—Co1—O1	92.37 (9)	C2—C3—H3	120.6
O2 ⁱ —Co1—O1 ⁱ	92.37 (9)	C4—C3—H3	120.6
O2—Co1—O1 ⁱ	87.63 (9)	C3—C4—C5	119.0 (3)
O1—Co1—O1 ⁱ	180.00 (4)	C3—C4—H4	120.5
O2 ⁱ —Co1—N1 ⁱ	88.56 (9)	C5—C4—H4	120.5
O2—Co1—N1 ⁱ	91.44 (9)	N1—C5—C4	123.1 (3)
O1—Co1—N1 ⁱ	89.87 (9)	N1—C5—H5	118.4
O1 ⁱ —Co1—N1 ⁱ	90.13 (9)	C4—C5—H5	118.4
O2 ⁱ —Co1—N1	91.44 (9)	C7—C6—C11	119.4 (3)
O2—Co1—N1	88.56 (9)	C7—C6—S1	120.8 (2)
O1—Co1—N1	90.13 (9)	C11—C6—S1	119.5 (2)
O1 ⁱ —Co1—N1	89.87 (9)	C8—C7—C6	120.1 (3)
N1 ⁱ —Co1—N1	180.000 (1)	C8—C7—H7	120.0
O4—S1—O3	114.83 (12)	C6—C7—H7	120.0
O4—S1—N2	105.51 (12)	C7—C8—C9	121.1 (3)
O3—S1—N2	113.63 (13)	C7—C8—H8	119.5
O4—S1—C6	106.39 (13)	C9—C8—H8	119.5
O3—S1—C6	106.34 (13)	N3—C9—C8	120.5 (3)
N2—S1—C6	109.90 (13)	N3—C9—C10	121.1 (3)
Co1—O1—H1A	120 (3)	C8—C9—C10	118.3 (3)
Co1—O1—H1B	124 (2)	C11—C10—C9	120.5 (3)
H1A—O1—H1B	111 (3)	C11—C10—H10	119.8
Co1—O2—H2A	125 (3)	C9—C10—H10	119.8
Co1—O2—H2B	120 (3)	C10—C11—C6	120.5 (3)
H2A—O2—H2B	111 (4)	C10—C11—H11	119.7
C1—N1—C5	116.5 (3)	C6—C11—H11	119.7
C1—N1—Co1	123.0 (2)	N4—C12—N2	113.2 (2)
C5—N1—Co1	120.4 (2)	N4—C12—C15	120.3 (3)
C12—N2—S1	121.93 (19)	N2—C12—C15	126.5 (3)
C9—N3—H3A	120.0	N5—C13—C14	124.8 (3)
C9—N3—H3B	120.0	N5—C13—C11	115.6 (2)
H3A—N3—H3B	120.0	C14—C13—C11	119.6 (2)
C12—N4—N5	120.4 (2)	C15—C14—C13	117.0 (3)
C13—N5—N4	118.8 (2)	C15—C14—H14	121.5
N1—C1—C2	123.7 (3)	C13—C14—H14	121.5
N1—C1—H1C	118.2	C14—C15—C12	118.7 (3)
C2—C1—H1C	118.2	C14—C15—H15	120.6
C3—C2—C1	119.0 (3)	C12—C15—H15	120.6
C3—C2—H2C	120.5		
O2 ⁱ —Co1—N1—C1	-57.6 (2)	O3—S1—C6—C11	173.7 (2)
O2—Co1—N1—C1	122.4 (2)	N2—S1—C6—C11	-62.9 (3)

O1—Co1—N1—C1	-145.2 (2)	C11—C6—C7—C8	-1.6 (4)
O1 ⁱ —Co1—N1—C1	34.8 (2)	S1—C6—C7—C8	173.1 (2)
O2 ⁱ —Co1—N1—C5	119.1 (2)	C6—C7—C8—C9	-1.0 (5)
O2—Co1—N1—C5	-60.9 (2)	C7—C8—C9—N3	-178.7 (3)
O1—Co1—N1—C5	31.5 (2)	C7—C8—C9—C10	2.8 (5)
O1 ⁱ —Co1—N1—C5	-148.5 (2)	N3—C9—C10—C11	179.5 (3)
O4—S1—N2—C12	173.4 (2)	C8—C9—C10—C11	-2.1 (4)
O3—S1—N2—C12	46.8 (3)	C9—C10—C11—C6	-0.5 (5)
C6—S1—N2—C12	-72.3 (3)	C7—C6—C11—C10	2.3 (4)
C12—N4—N5—C13	-0.2 (4)	S1—C6—C11—C10	-172.5 (2)
C5—N1—C1—C2	-0.3 (5)	N5—N4—C12—N2	178.8 (2)
Co1—N1—C1—C2	176.4 (3)	N5—N4—C12—C15	-1.3 (4)
N1—C1—C2—C3	0.1 (5)	S1—N2—C12—N4	-176.0 (2)
C1—C2—C3—C4	0.0 (5)	S1—N2—C12—C15	4.2 (4)
C2—C3—C4—C5	0.1 (5)	N4—N5—C13—C14	1.2 (5)
C1—N1—C5—C4	0.4 (5)	N4—N5—C13—C11	-178.8 (2)
Co1—N1—C5—C4	-176.4 (2)	N5—C13—C14—C15	-0.7 (5)
C3—C4—C5—N1	-0.3 (5)	C11—C13—C14—C15	179.3 (2)
O4—S1—C6—C7	-123.9 (2)	C13—C14—C15—C12	-0.9 (5)
O3—S1—C6—C7	-1.0 (3)	N4—C12—C15—C14	1.8 (5)
N2—S1—C6—C7	122.4 (2)	N2—C12—C15—C14	-178.3 (3)
O4—S1—C6—C11	50.8 (3)		

Symmetry code: (i) $-x+2, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3—H3 <i>A</i> \cdots N5 ⁱⁱ	0.88	2.52	3.135 (4)	127
N3—H3 <i>B</i> \cdots O3 ⁱⁱⁱ	0.88	2.23	3.001 (3)	147
O1—H1 <i>A</i> \cdots O3	0.81 (4)	2.07 (4)	2.874 (3)	174 (4)
O1—H1 <i>B</i> \cdots N4 ^{iv}	0.83 (4)	2.01 (4)	2.838 (3)	178 (3)
O2—H2 <i>A</i> \cdots O4	0.82 (3)	1.93 (4)	2.726 (3)	165 (4)
O2—H2 <i>B</i> \cdots N2 ^v	0.82 (4)	1.96 (4)	2.768 (3)	170 (4)
O2—H2 <i>B</i> \cdots O4 ^v	0.82 (4)	2.62 (4)	3.142 (3)	123 (3)

Symmetry codes: (ii) $x+1, y, z+1$; (iii) $x, y, z+1$; (iv) $x+1, y, z$; (v) $-x+1, -y+1, -z+1$.