

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-4-(2-Chlorobenzylideneamino)-3-(2-chlorophenyl)-1*H*-1,2,4-triazole-5(4*H*)-thione–(E)-1,5-bis(2-chlorobenzylidene)-thiocarbohydrazide–methanol (1/1/1)

Qingliang Guo

Department of Chemistry and Environmental Science, Taishan University, 271021 Taian, Shandong, People's Republic of China

Correspondence e-mail: jingtixuebao@163.com


Received 7 December 2009; accepted 10 December 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}–\text{C}) = 0.005$ Å; R factor = 0.046; wR factor = 0.111; data-to-parameter ratio = 14.5.

In the title compound, $\text{C}_{15}\text{H}_{12}\text{Cl}_2\text{N}_4\text{S} \cdot \text{C}_{15}\text{H}_{10}\text{Cl}_2\text{N}_4\text{S} \cdot \text{C}_2\text{H}_6\text{O}$, the two chlorophenyl rings of the triazole derivative form dihedral angles of 65.7 (2) and 44.2 (2)° with the triazole ring. In the thiocarbohydrazide derivative, the dihedral angle between the two chlorophenyl rings is 5.4 (2)°. In the crystal, the triazole, thiocarbohydrazide and methanol molecules are linked by $\text{N}–\text{H} \cdots \text{O}$, $\text{N}–\text{H} \cdots \text{S}$ and $\text{O}–\text{H} \cdots \text{S}$ hydrogen bonds, forming a hexameric unit.

Related literature

For general background to Schiff bases, see: Ren *et al.* (1999); Yang *et al.* (2005); Sen *et al.* (1998); Xia *et al.* (2007). For the biological activity of Schiff bases, see: Liang (2003); Bacci *et al.* (2005).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{10}\text{Cl}_2\text{N}_4\text{S} \cdot \text{C}_{15}\text{H}_{12}\text{Cl}_2\text{N}_4\text{S} \cdot \text{C}_2\text{H}_6\text{O}$	$\beta = 92.083$ (1)°
$M_r = 746.54$	$\gamma = 99.851$ (1)°
Triclinic, $P\bar{1}$	$V = 1744.8$ (2) Å ³
$a = 7.7086$ (6) Å	$Z = 2$
$b = 10.9999$ (9) Å	Mo $K\alpha$ radiation
$c = 20.9827$ (16) Å	$\mu = 0.50$ mm ⁻¹
$\alpha = 94.678$ (1)°	$T = 293$ K
	$0.35 \times 0.26 \times 0.23$ mm

Data collection

Bruker SMART APEX diffractometer	9324 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	6154 independent reflections
$T_{\min} = 0.845$, $T_{\max} = 0.894$	3724 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	425 parameters
$wR(F^2) = 0.111$	H-atom parameters constrained
$S = 1.01$	$\Delta\rho_{\text{max}} = 0.28$ e Å ⁻³
6154 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D–H \cdots A$	$D–H$	$H \cdots A$	$D \cdots A$	$D–H \cdots A$
$\text{N2}–\text{H2} \cdots \text{O1}$	0.86	2.20	2.978 (4)	150
$\text{N6}–\text{H6} \cdots \text{S1}^{\text{i}}$	0.86	2.41	3.261 (3)	170
$\text{N3}–\text{H3} \cdots \text{S2}^{\text{ii}}$	0.86	2.56	3.410 (2)	169
$\text{O1}–\text{H1} \cdots \text{S1}^{\text{iii}}$	0.82	2.65	3.413 (3)	156

 Symmetry codes: (i) $x, y - 1, z$; (ii) $x, y + 1, z$; (iii) $-x, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

The author thanks the Postgraduate Foundation of Taishan University for financial support (grant No. Y06-2-10).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2984).

References

- Bacci, A., Carcelli, M., Pelagatti, P., Pelizzi, G., Rodriguez-Arguelles, M. C., Rogolino, D., Solinas, C. & Zani, F. (2005). *J. Inorg. Biochem.* **99**, 397–408.
- Bruker (2005). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Liang, F.-Z. (2003). *J. Shandong Normal Univ. (Nat. Sci.)*, **18**, 50–51.
- Ren, Y. P., Dai, R. B., Wang, L. F. & Wu, J. G. (1999). *Synth. Commun.* **29**, 613–617.
- Sen, A. K., Singh, R. N., Handa, R. N., Dubey, S. N. & Squattrito, P. J. (1998). *J. Mol. Struct.* **470**, 61–69.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xia, H.-T., Liu, Y.-F., Yang, S.-P. & Wang, D.-Q. (2007). *Acta Cryst.* **E63**, o40–o41.
- Yang, J. G. & Pan, F. Y. (2005). *Chin. J. Struct. Chem.* **24**, 1403–1407.

supporting information

Acta Cryst. (2010). E66, o141 [doi:10.1107/S1600536809053203]

(E)-4-(2-Chlorobenzylideneamino)-3-(2-chlorophenyl)-1H-1,2,4-triazole-5(4H)-thione–(E)-1,5-bis(2-chlorobenzylidene)thiocarbonylhydrazide–methanol (1/1/1)

Qingliang Guo

S1. Comment

The synthesis and structural investigation of Schiff base compounds have attracted much attention due to their interesting structures and potential applications. Some of them have biological activities (Liang, 2003; Bacci *et al.*, 2005). They also play an important role in the development of coordination chemistry as well as inorganic biochemistry, catalysis and optical materials (Ren *et al.*, 1999; Yang *et al.* 2005; Sen *et al.* 1998).

The triazole ring forms dihedral angles of 65.7 (2) and 44.2 (2)°, respectively, with the C16-C21 and C25-C30 rings. The dihedral angle between the C1-C6 and C10-C15 rings is 5.4 (2)°. Two triazole, two thiocarbonylhydrazide and two methanol molecules are linked by N—H···O, N—H···S and O—H···S hydrogen bonds to form a hexamer.

S2. Experimental

The Schiff base compound was synthesized according to the modified method of Xia *et al.* (2007). A mixture of (2-chlorophenyl)methanamine and thiourea in methanol (30 ml) was refluxed for 3 h and filtered. The filtrate was placed for several days yielding colourless block-shaped crystals. (yield 79%). Elemental analysis: Calculated for C₃₂H₂₈Cl₄N₈OS₂: C 51.48, H 3.78, N 15.01; found: C 51.51, H 3.49, N 15.13.

S3. Refinement

The H atoms were found in a difference map, then placed in idealized positions (C-H = 0.93–0.97 Å, N-H = 0.86 Å and O-H = 0.82 Å), and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$ and $1.5U_{\text{eq}}(\text{O}, \text{C}_{\text{methyl}})$.




Figure 1

The asymmetric unit of the title compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms. H atoms have been omitted for clarity.

(*E*)-4-(2-Chlorobenzylideneamino)-3-(2-chlorophenyl)-1*H*-1,2,4-triazole-5(4*H*)-thione-(*E*)-1,5-bis(2-chlorobenzylidene)thiocarbonylhydrazide-methanol (1/1/1)

Crystal data

$C_{15}H_{10}Cl_2N_4S \cdot C_{15}H_{12}Cl_2N_4S \cdot C_2H_6O$

$M_r = 746.54$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.7086$ (6) Å

$b = 10.9999$ (9) Å

$c = 20.9827$ (16) Å

$\alpha = 94.678$ (1)°

$\beta = 92.083$ (1)°

$\gamma = 99.851$ (1)°

$V = 1744.8$ (2) Å³

$Z = 2$

$F(000) = 768$

$D_x = 1.421$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1667 reflections

$\theta = 2.2$ – 20.5 °

$\mu = 0.50$ mm⁻¹

$T = 293$ K

Block, colourless

$0.35 \times 0.26 \times 0.23$ mm

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.845$, $T_{\max} = 0.894$

9324 measured reflections

6154 independent reflections

3724 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 25.1$ °, $\theta_{\min} = 1.0$ °

$h = -7 \rightarrow 9$

$k = -12 \rightarrow 13$

$l = -24 \rightarrow 24$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.111$

$S = 1.01$

6154 reflections

425 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0365P)^2 + 0.4564P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.25610 (13)	1.26698 (8)	0.43362 (4)	0.0574 (3)
S2	0.40630 (12)	0.14705 (9)	0.24096 (4)	0.0614 (3)
Cl1	0.15231 (15)	0.94451 (10)	0.71997 (4)	0.0836 (3)
Cl2	0.48468 (13)	0.77336 (9)	0.21104 (4)	0.0738 (3)
Cl3	0.21316 (12)	0.52305 (8)	0.10925 (5)	0.0689 (3)
Cl4	0.22413 (15)	-0.15631 (9)	0.07317 (5)	0.0861 (3)
O1	0.1682 (4)	0.7954 (3)	0.52080 (14)	0.0837 (8)
H1	0.0608	0.7901	0.5211	0.126*
N1	0.1820 (3)	1.1229 (2)	0.54344 (12)	0.0487 (7)
N2	0.2441 (3)	1.0597 (2)	0.49247 (11)	0.0472 (7)
H2	0.2608	0.9847	0.4944	0.057*
N3	0.3341 (3)	1.0539 (2)	0.39024 (11)	0.0506 (7)
H3	0.3534	1.0877	0.3550	0.061*
N4	0.3618 (3)	0.9349 (2)	0.39441 (11)	0.0468 (6)
N5	0.0553 (4)	0.3611 (2)	0.26242 (12)	0.0529 (7)
N6	0.1796 (3)	0.2949 (2)	0.28209 (11)	0.0542 (7)
H6	0.2121	0.2943	0.3217	0.065*
N7	0.1565 (3)	0.2589 (2)	0.18135 (11)	0.0459 (6)
N8	0.1743 (3)	0.2251 (2)	0.11665 (11)	0.0463 (6)
C1	0.0902 (4)	1.0862 (3)	0.70951 (15)	0.0566 (9)
C2	0.1044 (4)	1.1357 (3)	0.65067 (14)	0.0463 (8)
C3	0.0605 (4)	1.2523 (3)	0.64682 (16)	0.0607 (9)
H3A	0.0690	1.2880	0.6081	0.073*
C4	0.0045 (5)	1.3162 (4)	0.6992 (2)	0.0762 (11)
H4	-0.0224	1.3948	0.6960	0.091*
C5	-0.0114 (5)	1.2632 (5)	0.7562 (2)	0.0841 (13)
H5	-0.0512	1.3057	0.7913	0.101*
C6	0.0306 (5)	1.1495 (4)	0.76173 (17)	0.0767 (12)
H6A	0.0192	1.1141	0.8005	0.092*
C7	0.1671 (4)	1.0702 (3)	0.59493 (14)	0.0480 (8)
H7	0.1953	0.9918	0.5973	0.058*
C8	0.2777 (4)	1.1187 (3)	0.43966 (14)	0.0434 (7)

C9	0.4020 (4)	0.8808 (3)	0.34211 (15)	0.0499 (8)
H9	0.4075	0.9231	0.3054	0.060*
C10	0.4389 (4)	0.7564 (3)	0.33816 (14)	0.0455 (8)
C11	0.4785 (4)	0.6972 (3)	0.28059 (15)	0.0515 (8)
C12	0.5155 (5)	0.5792 (3)	0.27706 (19)	0.0674 (10)
H12	0.5434	0.5420	0.2383	0.081*
C13	0.5112 (5)	0.5166 (4)	0.3307 (2)	0.0785 (12)
H13	0.5351	0.4364	0.3283	0.094*
C14	0.4719 (5)	0.5714 (3)	0.3881 (2)	0.0758 (11)
H14	0.4688	0.5284	0.4245	0.091*
C15	0.4369 (4)	0.6898 (3)	0.39182 (16)	0.0596 (9)
H15	0.4114	0.7265	0.4311	0.072*
C16	-0.0109 (4)	0.4711 (3)	0.11266 (15)	0.0498 (8)
C17	-0.0726 (4)	0.3890 (3)	0.15659 (14)	0.0444 (8)
C18	-0.2529 (5)	0.3556 (3)	0.16078 (16)	0.0599 (9)
H18	-0.2967	0.3019	0.1907	0.072*
C19	-0.3681 (5)	0.4010 (4)	0.12104 (19)	0.0719 (11)
H19	-0.4891	0.3765	0.1235	0.086*
C20	-0.3039 (6)	0.4819 (4)	0.07809 (18)	0.0759 (12)
H20	-0.3819	0.5130	0.0516	0.091*
C21	-0.1245 (5)	0.5180 (3)	0.07351 (16)	0.0667 (10)
H21	-0.0814	0.5735	0.0443	0.080*
C22	0.0461 (4)	0.3399 (3)	0.20069 (14)	0.0443 (8)
C23	0.2465 (4)	0.2311 (3)	0.23466 (14)	0.0460 (8)
C24	0.2022 (4)	0.1161 (3)	0.10273 (14)	0.0486 (8)
H24	0.2049	0.0624	0.1346	0.058*
C25	0.2301 (4)	0.0749 (3)	0.03672 (14)	0.0479 (8)
C26	0.2425 (4)	-0.0464 (3)	0.01809 (15)	0.0561 (9)
C27	0.2720 (5)	-0.0836 (4)	-0.0444 (2)	0.0790 (12)
H27	0.2803	-0.1657	-0.0561	0.095*
C28	0.2890 (6)	0.0012 (5)	-0.08861 (19)	0.0903 (14)
H28	0.3108	-0.0233	-0.1306	0.108*
C29	0.2744 (6)	0.1223 (5)	-0.07193 (18)	0.0872 (13)
H29	0.2839	0.1789	-0.1027	0.105*
C30	0.2458 (5)	0.1594 (3)	-0.00994 (15)	0.0651 (10)
H30	0.2366	0.2415	0.0013	0.078*
C31	0.1607 (8)	0.5962 (6)	0.5595 (4)	0.192 (4)
H31A	0.0341	0.5822	0.5574	0.288*
H31B	0.2023	0.5618	0.5966	0.288*
H31C	0.2014	0.5572	0.5217	0.288*
C32	0.2229 (7)	0.7192 (6)	0.5636 (3)	0.137 (2)
H32A	0.3501	0.7290	0.5621	0.164*
H32B	0.1986	0.7524	0.6060	0.164*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0842 (7)	0.0466 (5)	0.0462 (5)	0.0223 (5)	0.0086 (4)	0.0072 (4)

S2	0.0711 (6)	0.0711 (6)	0.0526 (5)	0.0330 (5)	0.0123 (5)	0.0211 (5)
Cl1	0.1144 (9)	0.0774 (7)	0.0609 (6)	0.0123 (6)	0.0128 (6)	0.0226 (5)
Cl2	0.0862 (7)	0.0867 (7)	0.0473 (5)	0.0142 (6)	0.0118 (5)	-0.0027 (5)
Cl3	0.0576 (6)	0.0654 (6)	0.0864 (7)	0.0088 (5)	0.0044 (5)	0.0266 (5)
Cl4	0.1162 (9)	0.0531 (6)	0.0935 (8)	0.0272 (6)	0.0055 (6)	0.0064 (5)
O1	0.086 (2)	0.0733 (18)	0.099 (2)	0.0229 (17)	0.0189 (17)	0.0211 (16)
N1	0.0572 (18)	0.0483 (16)	0.0421 (15)	0.0134 (13)	0.0085 (13)	0.0011 (13)
N2	0.0617 (18)	0.0441 (15)	0.0402 (14)	0.0176 (13)	0.0103 (13)	0.0080 (12)
N3	0.0725 (19)	0.0443 (16)	0.0391 (14)	0.0180 (14)	0.0112 (13)	0.0080 (12)
N4	0.0585 (18)	0.0404 (15)	0.0442 (15)	0.0139 (13)	0.0096 (13)	0.0052 (12)
N5	0.071 (2)	0.0499 (17)	0.0428 (15)	0.0239 (15)	0.0053 (14)	0.0018 (13)
N6	0.073 (2)	0.0553 (17)	0.0381 (15)	0.0198 (15)	0.0002 (14)	0.0066 (13)
N7	0.0568 (17)	0.0475 (16)	0.0378 (14)	0.0186 (13)	0.0076 (13)	0.0071 (12)
N8	0.0552 (17)	0.0499 (17)	0.0374 (14)	0.0180 (13)	0.0080 (12)	0.0049 (12)
C1	0.049 (2)	0.070 (2)	0.0462 (19)	0.0005 (18)	0.0070 (16)	-0.0004 (18)
C2	0.0424 (19)	0.050 (2)	0.0436 (18)	0.0020 (16)	0.0050 (15)	-0.0025 (16)
C3	0.059 (2)	0.066 (2)	0.056 (2)	0.0138 (19)	0.0020 (18)	-0.0050 (19)
C4	0.069 (3)	0.074 (3)	0.084 (3)	0.021 (2)	0.003 (2)	-0.021 (2)
C5	0.072 (3)	0.108 (4)	0.068 (3)	0.016 (3)	0.019 (2)	-0.027 (3)
C6	0.077 (3)	0.101 (3)	0.046 (2)	0.002 (3)	0.016 (2)	-0.005 (2)
C7	0.052 (2)	0.0485 (19)	0.0436 (18)	0.0088 (16)	0.0053 (15)	0.0021 (16)
C8	0.0454 (19)	0.0458 (19)	0.0402 (17)	0.0115 (15)	0.0025 (14)	0.0045 (15)
C9	0.059 (2)	0.047 (2)	0.0444 (18)	0.0098 (17)	0.0050 (16)	0.0072 (16)
C10	0.047 (2)	0.0381 (18)	0.0503 (19)	0.0071 (15)	0.0035 (15)	-0.0002 (15)
C11	0.045 (2)	0.053 (2)	0.055 (2)	0.0086 (16)	-0.0013 (16)	-0.0042 (17)
C12	0.065 (3)	0.058 (2)	0.077 (3)	0.016 (2)	0.001 (2)	-0.017 (2)
C13	0.081 (3)	0.048 (2)	0.107 (3)	0.021 (2)	-0.008 (3)	-0.004 (2)
C14	0.092 (3)	0.054 (2)	0.083 (3)	0.019 (2)	-0.007 (2)	0.013 (2)
C15	0.070 (3)	0.049 (2)	0.061 (2)	0.0120 (18)	0.0052 (19)	0.0037 (18)
C16	0.050 (2)	0.050 (2)	0.0506 (19)	0.0145 (16)	-0.0002 (16)	-0.0009 (16)
C17	0.055 (2)	0.0372 (17)	0.0430 (17)	0.0168 (16)	0.0058 (15)	-0.0046 (14)
C18	0.060 (2)	0.059 (2)	0.063 (2)	0.0171 (19)	0.0163 (19)	-0.0038 (18)
C19	0.052 (2)	0.081 (3)	0.084 (3)	0.023 (2)	0.003 (2)	-0.007 (2)
C20	0.071 (3)	0.096 (3)	0.069 (3)	0.042 (3)	-0.009 (2)	0.003 (2)
C21	0.072 (3)	0.078 (3)	0.058 (2)	0.031 (2)	0.002 (2)	0.018 (2)
C22	0.052 (2)	0.0393 (18)	0.0441 (18)	0.0124 (15)	0.0102 (15)	0.0049 (15)
C23	0.057 (2)	0.0432 (18)	0.0390 (17)	0.0106 (16)	0.0048 (15)	0.0058 (15)
C24	0.060 (2)	0.047 (2)	0.0419 (18)	0.0171 (17)	0.0053 (16)	0.0078 (15)
C25	0.049 (2)	0.055 (2)	0.0431 (18)	0.0173 (16)	0.0067 (15)	0.0044 (16)
C26	0.056 (2)	0.059 (2)	0.054 (2)	0.0198 (18)	-0.0012 (17)	-0.0064 (18)
C27	0.081 (3)	0.083 (3)	0.073 (3)	0.030 (2)	0.003 (2)	-0.025 (2)
C28	0.092 (3)	0.130 (4)	0.048 (2)	0.031 (3)	0.006 (2)	-0.023 (3)
C29	0.106 (4)	0.114 (4)	0.048 (2)	0.030 (3)	0.021 (2)	0.015 (2)
C30	0.078 (3)	0.073 (3)	0.051 (2)	0.027 (2)	0.0156 (19)	0.0117 (19)
C31	0.122 (5)	0.109 (5)	0.359 (11)	0.040 (4)	-0.016 (6)	0.079 (6)
C32	0.084 (4)	0.125 (5)	0.211 (7)	0.027 (4)	-0.004 (4)	0.056 (5)

Geometric parameters (Å, °)

S1—C8	1.682 (3)	C10—C11	1.394 (4)
S2—C23	1.669 (3)	C11—C12	1.373 (4)
C11—C1	1.735 (4)	C12—C13	1.364 (5)
C12—C11	1.739 (3)	C12—H12	0.93
C13—C16	1.730 (3)	C13—C14	1.370 (5)
C14—C26	1.732 (3)	C13—H13	0.93
O1—C32	1.380 (5)	C14—C15	1.371 (4)
O1—H1	0.82	C14—H14	0.93
N1—C7	1.267 (3)	C15—H15	0.93
N1—N2	1.373 (3)	C16—C21	1.371 (4)
N2—C8	1.341 (3)	C16—C17	1.383 (4)
N2—H2	0.86	C17—C18	1.383 (4)
N3—C8	1.338 (3)	C17—C22	1.474 (4)
N3—N4	1.372 (3)	C18—C19	1.378 (5)
N3—H3	0.86	C18—H18	0.93
N4—C9	1.279 (3)	C19—C20	1.365 (5)
N5—C22	1.294 (3)	C19—H19	0.93
N5—N6	1.370 (3)	C20—C21	1.381 (5)
N6—C23	1.340 (3)	C20—H20	0.93
N6—H6	0.86	C21—H21	0.93
N7—C23	1.378 (3)	C24—C25	1.457 (4)
N7—C22	1.382 (3)	C24—H24	0.93
N7—N8	1.396 (3)	C25—C26	1.380 (4)
N8—C24	1.267 (3)	C25—C30	1.398 (4)
C1—C6	1.385 (4)	C26—C27	1.379 (5)
C1—C2	1.389 (4)	C27—C28	1.363 (5)
C2—C3	1.390 (4)	C27—H27	0.93
C2—C7	1.463 (4)	C28—C29	1.374 (6)
C3—C4	1.379 (4)	C28—H28	0.93
C3—H3A	0.93	C29—C30	1.368 (5)
C4—C5	1.372 (5)	C29—H29	0.93
C4—H4	0.93	C30—H30	0.93
C5—C6	1.358 (5)	C31—C32	1.350 (7)
C5—H5	0.93	C31—H31A	0.96
C6—H6A	0.93	C31—H31B	0.96
C7—H7	0.93	C31—H31C	0.96
C9—C10	1.441 (4)	C32—H32A	0.97
C9—H9	0.93	C32—H32B	0.97
C10—C15	1.391 (4)		
C32—O1—H1	109.5	C14—C15—H15	119.3
C7—N1—N2	116.9 (2)	C10—C15—H15	119.3
C8—N2—N1	117.8 (2)	C21—C16—C17	121.3 (3)
C8—N2—H2	121.1	C21—C16—C13	118.6 (3)
N1—N2—H2	121.1	C17—C16—C13	120.0 (2)
C8—N3—N4	121.6 (2)	C16—C17—C18	118.5 (3)

C8—N3—H3	119.2	C16—C17—C22	122.6 (3)
N4—N3—H3	119.2	C18—C17—C22	118.9 (3)
C9—N4—N3	114.8 (2)	C19—C18—C17	120.7 (3)
C22—N5—N6	103.9 (2)	C19—C18—H18	119.7
C23—N6—N5	114.6 (2)	C17—C18—H18	119.7
C23—N6—H6	122.7	C20—C19—C18	119.7 (4)
N5—N6—H6	122.7	C20—C19—H19	120.1
C23—N7—C22	108.6 (2)	C18—C19—H19	120.1
C23—N7—N8	129.7 (2)	C19—C20—C21	120.8 (4)
C22—N7—N8	121.5 (2)	C19—C20—H20	119.6
C24—N8—N7	116.3 (2)	C21—C20—H20	119.6
C6—C1—C2	121.3 (3)	C16—C21—C20	119.1 (3)
C6—C1—C11	118.1 (3)	C16—C21—H21	120.5
C2—C1—C11	120.6 (2)	C20—C21—H21	120.5
C3—C2—C1	117.3 (3)	N5—C22—N7	110.7 (3)
C3—C2—C7	120.6 (3)	N5—C22—C17	125.3 (3)
C1—C2—C7	122.1 (3)	N7—C22—C17	123.9 (3)
C4—C3—C2	121.3 (3)	N6—C23—N7	102.1 (2)
C4—C3—H3A	119.3	N6—C23—S2	127.3 (2)
C2—C3—H3A	119.3	N7—C23—S2	130.5 (2)
C5—C4—C3	119.6 (4)	N8—C24—C25	119.9 (3)
C5—C4—H4	120.2	N8—C24—H24	120.0
C3—C4—H4	120.2	C25—C24—H24	120.0
C6—C5—C4	120.6 (4)	C26—C25—C30	118.0 (3)
C6—C5—H5	119.7	C26—C25—C24	122.0 (3)
C4—C5—H5	119.7	C30—C25—C24	120.0 (3)
C5—C6—C1	119.8 (4)	C27—C26—C25	121.4 (3)
C5—C6—H6A	120.1	C27—C26—C14	118.0 (3)
C1—C6—H6A	120.1	C25—C26—C14	120.6 (2)
N1—C7—C2	118.5 (3)	C28—C27—C26	119.3 (4)
N1—C7—H7	120.8	C28—C27—H27	120.3
C2—C7—H7	120.8	C26—C27—H27	120.3
N3—C8—N2	116.5 (3)	C27—C28—C29	120.8 (4)
N3—C8—S1	119.8 (2)	C27—C28—H28	119.6
N2—C8—S1	123.7 (2)	C29—C28—H28	119.6
N4—C9—C10	122.2 (3)	C30—C29—C28	119.9 (4)
N4—C9—H9	118.9	C30—C29—H29	120.0
C10—C9—H9	118.9	C28—C29—H29	120.0
C15—C10—C11	116.9 (3)	C29—C30—C25	120.6 (4)
C15—C10—C9	121.4 (3)	C29—C30—H30	119.7
C11—C10—C9	121.7 (3)	C25—C30—H30	119.7
C12—C11—C10	121.6 (3)	C32—C31—H31A	109.5
C12—C11—C12	118.3 (3)	C32—C31—H31B	109.5
C10—C11—C12	120.1 (2)	H31A—C31—H31B	109.5
C13—C12—C11	119.8 (3)	C32—C31—H31C	109.5
C13—C12—H12	120.1	H31A—C31—H31C	109.5
C11—C12—H12	120.1	H31B—C31—H31C	109.5
C12—C13—C14	120.3 (4)	C31—C32—O1	122.3 (6)

C12—C13—H13	119.8	C31—C32—H32A	106.8
C14—C13—H13	119.8	O1—C32—H32A	106.8
C13—C14—C15	120.0 (4)	C31—C32—H32B	106.8
C13—C14—H14	120.0	O1—C32—H32B	106.8
C15—C14—H14	120.0	H32A—C32—H32B	106.6
C14—C15—C10	121.4 (3)		
C7—N1—N2—C8	-173.4 (3)	C21—C16—C17—C22	-178.1 (3)
C8—N3—N4—C9	174.7 (3)	C13—C16—C17—C22	-1.6 (4)
C22—N5—N6—C23	-1.1 (4)	C16—C17—C18—C19	1.4 (5)
C23—N7—N8—C24	38.7 (4)	C22—C17—C18—C19	179.0 (3)
C22—N7—N8—C24	-147.1 (3)	C17—C18—C19—C20	-1.4 (5)
C6—C1—C2—C3	1.5 (5)	C18—C19—C20—C21	0.6 (6)
C11—C1—C2—C3	-176.9 (2)	C17—C16—C21—C20	-0.3 (5)
C6—C1—C2—C7	-180.0 (3)	C13—C16—C21—C20	-176.8 (3)
C11—C1—C2—C7	1.6 (4)	C19—C20—C21—C16	0.3 (6)
C1—C2—C3—C4	-0.2 (5)	N6—N5—C22—N7	2.1 (3)
C7—C2—C3—C4	-178.8 (3)	N6—N5—C22—C17	178.9 (3)
C2—C3—C4—C5	-1.1 (6)	C23—N7—C22—N5	-2.5 (4)
C3—C4—C5—C6	1.2 (6)	N8—N7—C22—N5	-177.7 (3)
C4—C5—C6—C1	0.1 (6)	C23—N7—C22—C17	-179.3 (3)
C2—C1—C6—C5	-1.5 (6)	N8—N7—C22—C17	5.5 (4)
C11—C1—C6—C5	176.9 (3)	C16—C17—C22—N5	115.1 (4)
N2—N1—C7—C2	178.9 (3)	C18—C17—C22—N5	-62.5 (4)
C3—C2—C7—N1	0.7 (5)	C16—C17—C22—N7	-68.6 (4)
C1—C2—C7—N1	-177.7 (3)	C18—C17—C22—N7	113.9 (3)
N4—N3—C8—N2	-2.2 (4)	N5—N6—C23—N7	-0.4 (3)
N4—N3—C8—S1	176.7 (2)	N5—N6—C23—S2	176.6 (2)
N1—N2—C8—N3	-178.5 (3)	C22—N7—C23—N6	1.6 (3)
N1—N2—C8—S1	2.6 (4)	N8—N7—C23—N6	176.4 (3)
N3—N4—C9—C10	178.8 (3)	C22—N7—C23—S2	-175.2 (3)
N4—C9—C10—C15	-1.7 (5)	N8—N7—C23—S2	-0.5 (5)
N4—C9—C10—C11	178.5 (3)	N7—N8—C24—C25	-177.3 (3)
C15—C10—C11—C12	-0.6 (5)	N8—C24—C25—C26	-173.1 (3)
C9—C10—C11—C12	179.3 (3)	N8—C24—C25—C30	7.1 (5)
C15—C10—C11—C12	-179.8 (2)	C30—C25—C26—C27	1.0 (5)
C9—C10—C11—C12	0.1 (4)	C24—C25—C26—C27	-178.8 (3)
C10—C11—C12—C13	1.0 (5)	C30—C25—C26—C14	-179.9 (3)
C12—C11—C12—C13	-179.8 (3)	C24—C25—C26—C14	0.3 (4)
C11—C12—C13—C14	-0.6 (6)	C25—C26—C27—C28	-0.2 (6)
C12—C13—C14—C15	-0.1 (6)	C14—C26—C27—C28	-179.3 (3)
C13—C14—C15—C10	0.6 (6)	C26—C27—C28—C29	-1.0 (7)
C11—C10—C15—C14	-0.2 (5)	C27—C28—C29—C30	1.3 (7)
C9—C10—C15—C14	179.9 (3)	C28—C29—C30—C25	-0.4 (6)
C21—C16—C17—C18	-0.5 (5)	C26—C25—C30—C29	-0.7 (5)
C13—C16—C17—C18	175.9 (2)	C24—C25—C30—C29	179.1 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O1	0.86	2.20	2.978 (4)	150
N6—H6 \cdots S1 ⁱ	0.86	2.41	3.261 (3)	170
N3—H3 \cdots S2 ⁱⁱ	0.86	2.56	3.410 (2)	169
O1—H1 \cdots S1 ⁱⁱⁱ	0.82	2.65	3.413 (3)	156

Symmetry codes: (i) $x, y-1, z$; (ii) $x, y+1, z$; (iii) $-x, -y+2, -z+1$.