

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-(3,4-Dihydroxyphenyl)-2-(4-hydroxyphenyl)ethanone

Zhu-Ping Xiao,^{a,b*} Shou-Fu Yi,^b Gao-Yu Ou,^b Fang Zhang^b and Jian Zhu^b

^aKey Laboratory of Plant Resources Conservation and Utilization (Jishou University), College of Hunan Province, Jishou University, Jishou 416000, People's Republic of China, and ^bCollege of Chemistry & Chemical Engineering, Jishou University, Jishou 416000, People's Republic of China

Correspondence e-mail: xiaozhuping2005@163.com


Received 14 November 2009; accepted 18 November 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.055; wR factor = 0.172; data-to-parameter ratio = 13.3.

The title compound, $\text{C}_{14}\text{H}_{12}\text{O}_4$, is a deoxybenzoin derivative in which the dihedral between the carbonyl group and the catechol unit is $5.99(3)^\circ$. The dihedral angle between the two benzene rings is $60.26(13)^\circ$. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds connect molecules, forming a two-dimensional network. In addition, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\pi$ contacts further stabilize the crystal structure.

Related literature

For synthetic applications of deoxybenzoin compounds, see: Xiao *et al.* (2007a, 2008a). For natural occurrences of these compounds, see: Kiuchi *et al.* (1990); Niwa *et al.* (1999); Sanduja *et al.* (1985). For their biological activity, see: Papoutsis *et al.* (2007); Xiao *et al.* (2007b, 2008b); Parmar *et al.* (1996). For a related structure, see: Xiao & Xiao (2008c).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{12}\text{O}_4$
 $M_r = 244.24$
 Triclinic, $P\bar{1}$
 $a = 5.7073(11)$ Å
 $b = 9.3464(19)$ Å
 $c = 11.202(2)$ Å
 $\alpha = 100.112(9)^\circ$
 $\beta = 94.792(9)^\circ$

$\gamma = 100.625(9)^\circ$
 $V = 573.99(19)$ Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.970$, $T_{\max} = 0.980$

3179 measured reflections
 2214 independent reflections
 1756 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.172$
 $S = 1.06$
 2214 reflections

166 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{O4}^{\text{ii}}$	0.82	1.94	2.744 (2)	168
$\text{O2}-\text{H2}\cdots\text{O3}^{\text{ii}}$	0.82	1.91	2.7274 (19)	171
$\text{O4}-\text{H4}\cdots\text{O2}^{\text{iii}}$	0.82	2.00	2.772 (2)	158
$\text{C3}-\text{H3}\cdots\text{O3}^{\text{ii}}$	0.93	2.53	3.191 (2)	129
$\text{C11}-\text{H11}\cdots\text{Cg1}^{\text{iv}}$	0.93	2.85	3.635 (2)	143

Symmetry codes: (i) $x-1, y-1, z-1$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, y+1, z+1$; (iv) $-x+1, -y+2, -z+1$. Cg1 is the centroid of the C1–C6 ring.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

This research was supported financially by the Key Laboratory of Plant Resources Conservation and Utilization (Jishou University), College of Hunan Province (grant No. JSK200904) and the Key Laboratory of Hunan Forest Products and Chemical Industry Engineering of Hunan Province (grant No. JDZ200904).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2957).

References

- Bruker (2007). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Kiuchi, F., Chen, X. & Tsuda, Y. (1990). *Heterocycles*, **31**, 629–36.
- Niwa, K., Hashimoto, M., Morishita, S., Yokoyama, Y., Mori, H. & Tamaya, T. (1999). *J. Cancer Res.* **90**, 726–32.
- Papoutsis, Z., Kassi, E., Fokialakis, N., Mitakou, S., Lambrinidis, G., Mikros, E. & Moutsatsou, P. (2007). *Steroids*, **72**, 693–704.
- Parmar, V. S., Bisht, K. S., Jain, R., Singh, S., Sharma, S. K., Gupta, S., Malhotra, S., Tyagi, O. D., Vardhan, A. & Pati, H. N. (1996). *Indian J. Chem. Sect. B*, **35**, 220–232.
- Sanduja, R., Weinheimer, A. J. & Alam, M. (1985). *J. Chem. Res. Synop.* pp. 56–57.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xiao, Z.-P., Fang, R.-Q., Li, H.-Q., Xue, J.-Y., Zheng, Y. & Zhu, H.-L. (2008b). *Eur. J. Med. Chem.* **43**, 1828–1836.
- Xiao, Z.-P., Fang, R.-Q., Shi, L., Ding, H., Xu, C. & Zhu, H.-L. (2007a). *Can. J. Chem.* **85**, 951–957.
- Xiao, Z.-P., Li, H.-Q., Xue, J.-Y., Shi, L. & Zhu, H.-L. (2008a). *Synth. Commun.* **38**, 525–529.
- Xiao, Z.-P., Shi, D.-H., Li, H.-Q., Zhang, L.-N., Xu, C. & Zhu, H.-L. (2007b). *Bioorg. Med. Chem.* **15**, 3703–3710.
- Xiao, Z.-P. & Xiao, H.-Y. (2008c). *Acta Cryst.* **E64**, o2324.

supporting information

Acta Cryst. (2010). E66, o4 [doi:10.1107/S1600536809049393]

1-(3,4-Dihydroxyphenyl)-2-(4-hydroxyphenyl)ethanone

Zhu-Ping Xiao, Shou-Fu Yi, Gao-Yu Ou, Fang Zhang and Jian Zhu

S1. Comment

Deoxybenzoin compounds are intermediates in the syntheses of isoflavones (Xiao *et al.*, 2008a; Xiao *et al.*, 2007a) and are found in several plants, such as *Glycyrrhiza sp.*, *Trifolium subterraneum* and *Ononis spinosa*, and marine sources (Kiuchi *et al.*, 1990; Niwa *et al.*, 1999; Sanduja *et al.*, 1985). Many deoxybenzoin compounds have shown significant estrogen receptor modulatory, urease inhibitory, antimicrobial and antiviral properties (Papoutsi *et al.*, 2007; Xiao *et al.*, 2007b; Xiao *et al.*, 2008b; Parmar *et al.*, 1996). As part of our work involving the synthesis of a series of deoxybenzoin derivatives for urease inhibitory activity screening we report herein the crystal structure of the title deoxybenzoin derivative (I).


The molecular structure of the title compound is shown in Fig. 1. The two benzene rings form a dihedral angle of 60.26 (13)°. The carbonyl group forms a dihedral angle of 5.99 (3)° with the catechol moiety [O1/O2/C1-C6] while a similar angle is 1.95 (13)° in our previous related crystal structure (Xiao *et al.*, 2008c). In the crystal structure, intermolecular O—H···O connect molecules to form a two-dimensional network. In addition, weak intermolecular C—H···O hydrogen bonds and C—H··· π contacts further stabilize the crystal structure (see Fig. 2).

S2. Experimental


0.55 g (5 mmol) of catechol and 0.76 g (5 mmol) of *p*-hydroxyphenylacetic acid were dissolved into 10 ml of fresh distilled BF₃.Et₂O. The mixture was stirred and heated on an oil bath at 353 K for about 3 h. After cooling, the contents were poured into 150 ml of ice-cold aqueous sodium acetate (w % = 10%) with stirring. Then, the precipitate was filtered and washed three times with water. The resulting solid was crystallized from methanol-water to give colorless blocks of (I) suitable for single-crystal structure determination.

S3. Refinement

All H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H of 0.93 Å for the aromatic atoms, 0.97 Å for the CH₂ groups and 0.82 Å for the OH groups. $U_{iso}(H)$ values were set at 1.2 times $U_{eq}(C)$ for aromatic C and CH₂ groups and 1.5 times $U_{eq}(O)$ for O—H groups.

**Figure 1**

Molecular structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

Part of the crystal structure of (I) with hydrogen bonds indicated by thin dashed lines and C—H... π contacts shown as thick dashed lines.

1-(3,4-Dihydroxyphenyl)-2-(4-hydroxyphenyl)ethanone

Crystal data

$C_{14}H_{12}O_4$
 $M_r = 244.24$
 Triclinic, $P\bar{1}$
 Hall symbol: $-P\ 1$
 $a = 5.7073$ (11) Å
 $b = 9.3464$ (19) Å
 $c = 11.202$ (2) Å
 $\alpha = 100.112$ (9)°
 $\beta = 94.792$ (9)°
 $\gamma = 100.625$ (9)°
 $V = 573.99$ (19) Å³

$Z = 2$
 $F(000) = 256$
 $D_x = 1.413$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 1785 reflections
 $\theta = 2.0$ – 26.0 °
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
 Block, colorless
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD diffractometer	3179 measured reflections
Radiation source: fine-focus sealed tube	2214 independent reflections
Graphite monochromator	1756 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.028$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 1.9^\circ$
$T_{\text{min}} = 0.970$, $T_{\text{max}} = 0.980$	$h = -7 \rightarrow 6$
	$k = -11 \rightarrow 6$
	$l = -13 \rightarrow 13$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.055$	H-atom parameters constrained
$wR(F^2) = 0.172$	$w = 1/[\sigma^2(F_o^2) + (0.1062P)^2 + 0.1004P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2214 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
166 parameters	$\Delta\rho_{\text{max}} = 0.26 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.32 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0231 (3)	0.4370 (2)	0.17089 (17)	0.0357 (4)
C2	0.2221 (3)	0.4285 (2)	0.24884 (17)	0.0342 (4)
C3	0.2784 (3)	0.5201 (2)	0.36129 (17)	0.0357 (5)
H3	0.4123	0.5140	0.4120	0.043*
C4	0.1374 (3)	0.62287 (19)	0.40118 (17)	0.0335 (4)
C5	-0.0606 (3)	0.6309 (2)	0.32327 (18)	0.0364 (5)
H5	-0.1561	0.6986	0.3480	0.044*
C6	-0.1164 (3)	0.5392 (2)	0.20949 (18)	0.0382 (5)
H6	-0.2490	0.5460	0.1582	0.046*
C7	0.2107 (4)	0.7192 (2)	0.52317 (18)	0.0377 (5)
C8	0.0779 (4)	0.8423 (2)	0.56251 (19)	0.0434 (5)
H8A	0.0603	0.8942	0.4956	0.052*
H8B	-0.0821	0.7977	0.5769	0.052*
C9	0.1937 (3)	0.9545 (2)	0.67514 (18)	0.0376 (5)
C10	0.4194 (4)	1.0415 (2)	0.67929 (19)	0.0464 (5)
H10	0.5049	1.0265	0.6126	0.056*

C11	0.5210 (4)	1.1498 (2)	0.77965 (19)	0.0466 (5)
H11	0.6730	1.2070	0.7802	0.056*
C12	0.3970 (3)	1.1730 (2)	0.87899 (17)	0.0375 (5)
C13	0.1733 (4)	1.0874 (2)	0.8778 (2)	0.0497 (6)
H13	0.0887	1.1019	0.9449	0.060*
C14	0.0751 (4)	0.9795 (2)	0.7761 (2)	0.0497 (6)
H14	-0.0763	0.9220	0.7760	0.060*
O1	-0.0188 (3)	0.34464 (17)	0.06112 (13)	0.0505 (4)
H1	-0.1572	0.3388	0.0313	0.076*
O2	0.3534 (2)	0.32578 (15)	0.20545 (12)	0.0420 (4)
H2	0.4435	0.3133	0.2620	0.063*
O3	0.3760 (3)	0.70119 (18)	0.59031 (14)	0.0609 (5)
O4	0.5023 (3)	1.28604 (17)	0.97474 (14)	0.0507 (4)
H4	0.4249	1.2818	1.0329	0.076*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0337 (10)	0.0355 (9)	0.0340 (10)	0.0061 (7)	-0.0017 (7)	0.0002 (8)
C2	0.0314 (10)	0.0336 (9)	0.0374 (10)	0.0096 (7)	0.0032 (7)	0.0034 (8)
C3	0.0340 (10)	0.0344 (9)	0.0373 (10)	0.0113 (8)	-0.0047 (8)	0.0028 (8)
C4	0.0343 (10)	0.0291 (9)	0.0366 (10)	0.0096 (7)	-0.0001 (8)	0.0036 (7)
C5	0.0319 (10)	0.0337 (9)	0.0433 (11)	0.0123 (7)	0.0005 (8)	0.0028 (8)
C6	0.0317 (10)	0.0392 (10)	0.0412 (11)	0.0101 (8)	-0.0056 (8)	0.0027 (8)
C7	0.0404 (11)	0.0330 (9)	0.0402 (11)	0.0141 (8)	-0.0016 (8)	0.0045 (8)
C8	0.0404 (11)	0.0383 (10)	0.0479 (12)	0.0162 (8)	-0.0047 (9)	-0.0052 (9)
C9	0.0377 (11)	0.0328 (9)	0.0413 (11)	0.0149 (8)	-0.0002 (8)	-0.0011 (8)
C10	0.0429 (12)	0.0534 (12)	0.0379 (11)	0.0079 (9)	0.0099 (9)	-0.0047 (9)
C11	0.0358 (11)	0.0521 (12)	0.0443 (12)	0.0021 (9)	0.0057 (9)	-0.0040 (9)
C12	0.0379 (11)	0.0371 (10)	0.0345 (10)	0.0116 (8)	-0.0008 (8)	-0.0030 (8)
C13	0.0486 (13)	0.0489 (12)	0.0470 (13)	0.0061 (10)	0.0174 (10)	-0.0046 (10)
C14	0.0391 (12)	0.0420 (11)	0.0610 (14)	0.0010 (9)	0.0114 (10)	-0.0039 (10)
O1	0.0472 (9)	0.0560 (9)	0.0417 (9)	0.0205 (7)	-0.0100 (7)	-0.0116 (7)
O2	0.0402 (8)	0.0456 (8)	0.0390 (8)	0.0206 (6)	-0.0024 (6)	-0.0043 (6)
O3	0.0755 (12)	0.0577 (10)	0.0467 (9)	0.0401 (8)	-0.0243 (8)	-0.0110 (7)
O4	0.0450 (9)	0.0559 (9)	0.0415 (8)	0.0076 (7)	0.0011 (6)	-0.0119 (7)

Geometric parameters (Å, °)

C1—O1	1.348 (2)	C8—H8B	0.9700
C1—C6	1.388 (3)	C9—C14	1.373 (3)
C1—C2	1.396 (3)	C9—C10	1.383 (3)
C2—C3	1.369 (3)	C10—C11	1.378 (3)
C2—O2	1.372 (2)	C10—H10	0.9300
C3—C4	1.402 (3)	C11—C12	1.375 (3)
C3—H3	0.9300	C11—H11	0.9300
C4—C5	1.390 (2)	C12—C13	1.374 (3)
C4—C7	1.479 (3)	C12—O4	1.374 (2)

C5—C6	1.382 (3)	C13—C14	1.382 (3)
C5—H5	0.9300	C13—H13	0.9300
C6—H6	0.9300	C14—H14	0.9300
C7—O3	1.212 (2)	O1—H1	0.8200
C7—C8	1.514 (3)	O2—H2	0.8200
C8—C9	1.506 (3)	O4—H4	0.8200
C8—H8A	0.9700		
O1—C1—C6	124.22 (17)	C9—C8—H8B	108.4
O1—C1—C2	116.86 (17)	C7—C8—H8B	108.4
C6—C1—C2	118.92 (17)	H8A—C8—H8B	107.4
C3—C2—O2	123.43 (16)	C14—C9—C10	117.12 (19)
C3—C2—C1	120.35 (17)	C14—C9—C8	121.34 (19)
O2—C2—C1	116.22 (16)	C10—C9—C8	121.47 (19)
C2—C3—C4	121.03 (16)	C11—C10—C9	121.7 (2)
C2—C3—H3	119.5	C11—C10—H10	119.1
C4—C3—H3	119.5	C9—C10—H10	119.1
C5—C4—C3	118.43 (17)	C12—C11—C10	119.8 (2)
C5—C4—C7	123.56 (16)	C12—C11—H11	120.1
C3—C4—C7	118.00 (16)	C10—C11—H11	120.1
C6—C5—C4	120.53 (17)	C13—C12—O4	122.75 (18)
C6—C5—H5	119.7	C13—C12—C11	119.75 (19)
C4—C5—H5	119.7	O4—C12—C11	117.47 (18)
C5—C6—C1	120.75 (17)	C12—C13—C14	119.4 (2)
C5—C6—H6	119.6	C12—C13—H13	120.3
C1—C6—H6	119.6	C14—C13—H13	120.3
O3—C7—C4	120.71 (17)	C9—C14—C13	122.2 (2)
O3—C7—C8	120.36 (18)	C9—C14—H14	118.9
C4—C7—C8	118.93 (15)	C13—C14—H14	118.9
C9—C8—C7	115.66 (16)	C1—O1—H1	109.5
C9—C8—H8A	108.4	C2—O2—H2	109.5
C7—C8—H8A	108.4	C12—O4—H4	109.5
O1—C1—C2—C3	179.11 (17)	C3—C4—C7—C8	173.33 (18)
C6—C1—C2—C3	-0.2 (3)	O3—C7—C8—C9	11.5 (3)
O1—C1—C2—O2	-0.5 (3)	C4—C7—C8—C9	-167.73 (17)
C6—C1—C2—O2	-179.76 (17)	C7—C8—C9—C14	-121.6 (2)
O2—C2—C3—C4	-179.83 (17)	C7—C8—C9—C10	61.4 (3)
C1—C2—C3—C4	0.6 (3)	C14—C9—C10—C11	-0.6 (3)
C2—C3—C4—C5	-0.6 (3)	C8—C9—C10—C11	176.50 (18)
C2—C3—C4—C7	-179.35 (16)	C9—C10—C11—C12	0.1 (3)
C3—C4—C5—C6	0.2 (3)	C10—C11—C12—C13	0.4 (3)
C7—C4—C5—C6	178.83 (17)	C10—C11—C12—O4	-177.56 (18)
C4—C5—C6—C1	0.3 (3)	O4—C12—C13—C14	177.42 (18)
O1—C1—C6—C5	-179.49 (18)	C11—C12—C13—C14	-0.5 (3)
C2—C1—C6—C5	-0.3 (3)	C10—C9—C14—C13	0.6 (3)
C5—C4—C7—O3	175.4 (2)	C8—C9—C14—C13	-176.54 (19)
C3—C4—C7—O3	-5.9 (3)	C12—C13—C14—C9	-0.1 (3)

C5—C4—C7—C8 -5.3 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O4 ⁱ	0.82	1.94	2.744 (2)	168
O2—H2 \cdots O3 ⁱⁱ	0.82	1.91	2.7274 (19)	171
O4—H4 \cdots O2 ⁱⁱⁱ	0.82	2.00	2.772 (2)	158
C3—H3 \cdots O3 ⁱⁱ	0.93	2.53	3.191 (2)	129
C11—H11 \cdots Cg1 ^{iv}	0.93	2.85	3.635 (2)	143

Symmetry codes: (i) $x-1, y-1, z-1$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, y+1, z+1$; (iv) $-x+1, -y+2, -z+1$.