

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# N'-(4-Chlorobenzylidene)furan-2-carbohydrazide monohydrate

Jin-He Jiang

 Microscale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China  
 Correspondence e-mail: weifangjih@126.com


Received 21 January 2010; accepted 10 February 2010

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.045;  $wR$  factor = 0.145; data-to-parameter ratio = 16.4.

In the title compound,  $\text{C}_{12}\text{H}_9\text{ClN}_2\text{O}_2 \cdot \text{H}_2\text{O}$ , the dihedral angle between the aromatic rings is  $13.9(2)^\circ$  and an intramolecular  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bond occurs. In the crystal structure, the components are linked by  $\text{N}-\text{H} \cdots \text{O}$ ,  $\text{O}-\text{H} \cdots \text{O}$  and  $\text{C}-\text{H} \cdots \text{O}$  hydrogen bonds.

## Related literature

For background to Schiff bases, see: Cimerman *et al.* (1997).  
 For a related structure, see: Girgis (2006).



## Experimental

## Crystal data

 $\text{C}_{12}\text{H}_9\text{ClN}_2\text{O}_2 \cdot \text{H}_2\text{O}$   
 $M_r = 266.68$   
 Monoclinic,  $P2_1$   
 $a = 4.5480(9)$  Å  
 $b = 12.423(3)$  Å

 $c = 10.971(2)$  Å  
 $\beta = 91.90(3)^\circ$   
 $V = 619.5(2)$  Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation

 $\mu = 0.31$  mm<sup>-1</sup>  
 $T = 293$  K

 $0.25 \times 0.20 \times 0.18$  mm

## Data collection

 Bruker SMART CCD  
 diffractometer  
 5979 measured reflections

 2799 independent reflections  
 1708 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.042$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.145$   
 $S = 1.04$   
 2799 reflections  
 171 parameters  
 1 restraint

 H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.23$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.34$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983),  
 1319 Friedel pairs  
 Flack parameter: 0.01 (11)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O1W}-\text{H1W} \cdots \text{O1}^i$	0.97 (5)	1.90 (5)	2.865 (4)	174 (5)
$\text{O1W}-\text{H2W} \cdots \text{O1}^{ii}$	0.92 (6)	1.97 (6)	2.873 (4)	168 (5)
$\text{N1}-\text{H1A} \cdots \text{O2}$	0.86	2.36	2.715 (4)	106
$\text{N1}-\text{H1A} \cdots \text{O1W}$	0.86	2.05	2.886 (4)	165
$\text{C8}-\text{H8A} \cdots \text{O1W}$	0.93	2.49	3.290 (5)	144

 Symmetry codes: (i)  $-x, y - \frac{1}{2}, -z + 2$ ; (ii)  $-x - 1, y - \frac{1}{2}, -z + 2$ .

Data collection: *SMART* (Bruker 1997); cell refinement: *SAINTE* (Bruker 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5316).

## References

- Bruker (1997). *SMART, SAINTE and SADABS*. Bruker AXS, Inc., Madison, Wisconsin, USA.  
 Cimerman, Z., Galic, N. & Bosner, B. (1997). *Anal. Chim. Acta*, **343**, 145–153.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Girgis, A. S. (2006). *J. Chem. Res.* pp. 81–85.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2010). E66, o627 [doi:10.1107/S1600536810005532]

***N'*-(4-Chlorobenzylidene)furan-2-carbohydrazide monohydrate****Jin-He Jiang****S1. Comment**

Schiff bases have received considerable attention in the literature. They are attractive from several points of view, such as the possibility of analytical application (Cimerman *et al.*, 1997). As part of our search for new schiff base compounds we synthesized the title compound (I), and describe its structure here.


The molecular structure of (I) is shown in Fig. 1. The C7—N1 bond length of 1.342 (4)Å is longer than the C—N double bond [1.281 (2) Å] reported (Girgis, 2006). In the crystal structure, molecules are linked by intermolecular N—H···O hydrogen bonds.

**S2. Experimental**

A mixture of furan-2-carbohydrazide (0.1 mol), and 4-chlorobenzaldehyde (0.1 mol) was stirred in refluxing ethanol (20 mL) for 4 h to afford the title compound (0.087 mol, yield 87%). Colourless blocks of (I) were obtained by recrystallization from ethanol at room temperature.

**S3. Refinement**

H atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances = 0.93-0.97 Å; N—H = 0.86 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C},\text{N})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ .

**Figure 1**

The structure of (I) showing 30% probability displacement ellipsoids.

### *N'*-(4-Chlorobenzylidene)furan-2-carbohydrazide monohydrate

#### *Crystal data*

$C_{12}H_9ClN_2O_2 \cdot H_2O$

$M_r = 266.68$

Monoclinic,  $P2_1$

Hall symbol:  $P\ 2y_b$

$a = 4.5480(9)\ \text{\AA}$

$b = 12.423(3)\ \text{\AA}$

$c = 10.971(2)\ \text{\AA}$

$\beta = 91.90(3)^\circ$

$V = 619.5(2)\ \text{\AA}^3$

$Z = 2$

$F(000) = 276$

$D_x = 1.430\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2022 reflections

$\theta = 3.3\text{--}27.3^\circ$

$\mu = 0.31\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colourless

$0.25 \times 0.20 \times 0.18\ \text{mm}$

*Data collection*

Bruker SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

5979 measured reflections

2799 independent reflections

1708 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\text{max}} = 27.5^\circ$ ,  $\theta_{\text{min}} = 3.3^\circ$

$h = -5 \rightarrow 5$

$k = -16 \rightarrow 16$

$l = -14 \rightarrow 14$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.145$

$S = 1.04$

2799 reflections

171 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H atoms treated by a mixture of independent  
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0702P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.23 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.34 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack (1983), 1319 Friedel  
pairs

Absolute structure parameter: 0.01 (11)

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.8615 (3)	0.97667 (10)	0.39621 (9)	0.0713 (4)
O1	-0.3072 (6)	0.94704 (18)	1.0474 (2)	0.0525 (6)
O2	-0.5944 (6)	0.68010 (17)	1.0623 (2)	0.0457 (6)
N1	-0.1934 (6)	0.7961 (2)	0.9399 (2)	0.0412 (6)
H1A	-0.2136	0.7276	0.9323	0.049*
N2	-0.0118 (6)	0.8522 (2)	0.8633 (2)	0.0428 (7)
C1	0.6164 (11)	0.8151 (4)	0.5209 (4)	0.0677 (12)
H1B	0.7036	0.7692	0.4657	0.081*
C2	0.6554 (8)	0.9240 (3)	0.5124 (3)	0.0527 (10)
C3	0.5303 (9)	0.9909 (3)	0.5956 (3)	0.0568 (9)
H3A	0.5619	1.0647	0.5907	0.068*
C4	0.3591 (9)	0.9509 (3)	0.6861 (3)	0.0544 (10)
H4A	0.2762	0.9975	0.7417	0.065*
C5	0.3098 (8)	0.8405 (3)	0.6946 (3)	0.0466 (8)
C6	0.4448 (10)	0.7733 (3)	0.6129 (4)	0.0616 (11)

H6A	0.4209	0.6992	0.6193	0.074*
C7	-0.3375 (7)	0.8492 (3)	1.0262 (3)	0.0409 (7)
C8	0.1202 (8)	0.7935 (3)	0.7861 (3)	0.0466 (8)
H8A	0.0950	0.7192	0.7881	0.056*
C9	-0.5355 (7)	0.7844 (3)	1.0984 (3)	0.0398 (7)
C10	-0.6880 (9)	0.8080 (3)	1.1980 (3)	0.0483 (9)
H10A	-0.6893	0.8733	1.2393	0.058*
C11	-0.8474 (9)	0.7132 (3)	1.2281 (3)	0.0515 (9)
H11A	-0.9722	0.7044	1.2928	0.062*
C12	-0.7811 (9)	0.6394 (3)	1.1444 (3)	0.0513 (9)
H12A	-0.8536	0.5694	1.1428	0.062*
O1W	-0.1978 (8)	0.5707 (2)	0.8764 (3)	0.0596 (8)
H1W	-0.037 (11)	0.526 (4)	0.906 (4)	0.084 (15)*
H2W	-0.372 (13)	0.537 (4)	0.893 (4)	0.093 (18)*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C11	0.0710 (7)	0.0849 (8)	0.0595 (5)	-0.0026 (6)	0.0271 (5)	0.0050 (5)
O1	0.0543 (16)	0.0340 (13)	0.0700 (15)	-0.0032 (11)	0.0163 (12)	-0.0029 (12)
O2	0.0529 (15)	0.0354 (12)	0.0497 (13)	-0.0044 (11)	0.0163 (11)	-0.0023 (10)
N1	0.0388 (16)	0.0333 (14)	0.0522 (15)	-0.0034 (12)	0.0113 (12)	0.0006 (13)
N2	0.0353 (15)	0.0413 (15)	0.0521 (16)	-0.0009 (13)	0.0071 (13)	0.0022 (13)
C1	0.081 (3)	0.064 (3)	0.060 (2)	0.003 (2)	0.034 (2)	-0.012 (2)
C2	0.044 (2)	0.064 (3)	0.051 (2)	-0.0025 (18)	0.0087 (16)	-0.0037 (17)
C3	0.065 (2)	0.050 (2)	0.056 (2)	-0.007 (2)	0.0161 (18)	-0.0020 (19)
C4	0.060 (2)	0.048 (2)	0.058 (2)	-0.0010 (18)	0.0225 (18)	-0.0051 (18)
C5	0.042 (2)	0.051 (2)	0.0475 (19)	-0.0007 (17)	0.0098 (15)	-0.0014 (16)
C6	0.072 (3)	0.051 (2)	0.063 (2)	-0.008 (2)	0.025 (2)	-0.0125 (19)
C7	0.0359 (18)	0.0370 (18)	0.0503 (18)	0.0050 (15)	0.0092 (14)	0.0040 (15)
C8	0.044 (2)	0.0413 (18)	0.0550 (19)	-0.0018 (16)	0.0117 (16)	-0.0042 (16)
C9	0.0392 (19)	0.0361 (17)	0.0442 (16)	0.0035 (15)	0.0045 (14)	0.0028 (14)
C10	0.052 (2)	0.045 (2)	0.0493 (18)	0.0023 (16)	0.0125 (16)	-0.0033 (16)
C11	0.054 (2)	0.048 (2)	0.053 (2)	0.0092 (17)	0.0221 (17)	0.0093 (17)
C12	0.059 (2)	0.042 (2)	0.055 (2)	-0.0053 (17)	0.0195 (18)	0.0072 (16)
O1W	0.053 (2)	0.0392 (14)	0.087 (2)	-0.0005 (14)	0.0169 (16)	-0.0001 (14)

*Geometric parameters (Å, °)*

C11—C2	1.735 (4)	C4—H4A	0.9300
O1—C7	1.244 (4)	C5—C6	1.384 (5)
O2—C12	1.356 (4)	C5—C8	1.466 (5)
O2—C9	1.379 (4)	C6—H6A	0.9300
N1—C7	1.342 (4)	C7—C9	1.461 (5)
N1—N2	1.386 (4)	C8—H8A	0.9300
N1—H1A	0.8600	C9—C10	1.346 (5)
N2—C8	1.282 (4)	C10—C11	1.427 (6)
C1—C2	1.368 (6)	C10—H10A	0.9300

C1—C6	1.396 (6)	C11—C12	1.339 (5)
C1—H1B	0.9300	C11—H11A	0.9300
C2—C3	1.372 (5)	C12—H12A	0.9300
C3—C4	1.375 (5)	O1W—H1W	0.97 (5)
C3—H3A	0.9300	O1W—H2W	0.92 (6)
C4—C5	1.393 (5)		
C12—O2—C9	106.2 (3)	C5—C6—H6A	119.5
C7—N1—N2	119.7 (3)	C1—C6—H6A	119.5
C7—N1—H1A	120.1	O1—C7—N1	123.9 (3)
N2—N1—H1A	120.1	O1—C7—C9	120.2 (3)
C8—N2—N1	114.6 (3)	N1—C7—C9	115.8 (3)
C2—C1—C6	119.6 (4)	N2—C8—C5	121.7 (3)
C2—C1—H1B	120.2	N2—C8—H8A	119.2
C6—C1—H1B	120.2	C5—C8—H8A	119.2
C1—C2—C3	119.7 (4)	C10—C9—O2	109.7 (3)
C1—C2—C11	119.8 (3)	C10—C9—C7	131.6 (3)
C3—C2—C11	120.5 (3)	O2—C9—C7	118.6 (3)
C2—C3—C4	121.3 (4)	C9—C10—C11	106.7 (3)
C2—C3—H3A	119.4	C9—C10—H10A	126.6
C4—C3—H3A	119.4	C11—C10—H10A	126.6
C3—C4—C5	120.0 (4)	C12—C11—C10	106.2 (3)
C3—C4—H4A	120.0	C12—C11—H11A	126.9
C5—C4—H4A	120.0	C10—C11—H11A	126.9
C6—C5—C4	118.4 (4)	C11—C12—O2	111.1 (3)
C6—C5—C8	119.1 (3)	C11—C12—H12A	124.4
C4—C5—C8	122.5 (3)	O2—C12—H12A	124.4
C5—C6—C1	121.0 (4)	H1W—O1W—H2W	109 (5)
C7—N1—N2—C8	178.5 (3)	C6—C5—C8—N2	-179.5 (3)
C6—C1—C2—C3	1.2 (7)	C4—C5—C8—N2	0.2 (6)
C6—C1—C2—C11	-178.5 (3)	C12—O2—C9—C10	-1.6 (4)
C1—C2—C3—C4	-1.7 (6)	C12—O2—C9—C7	180.0 (3)
C11—C2—C3—C4	178.0 (3)	O1—C7—C9—C10	-6.7 (6)
C2—C3—C4—C5	0.0 (6)	N1—C7—C9—C10	172.3 (4)
C3—C4—C5—C6	2.1 (6)	O1—C7—C9—O2	171.4 (3)
C3—C4—C5—C8	-177.6 (3)	N1—C7—C9—O2	-9.6 (4)
C4—C5—C6—C1	-2.6 (6)	O2—C9—C10—C11	1.1 (4)
C8—C5—C6—C1	177.1 (4)	C7—C9—C10—C11	179.3 (3)
C2—C1—C6—C5	1.0 (7)	C9—C10—C11—C12	-0.3 (4)
N2—N1—C7—O1	-4.2 (5)	C10—C11—C12—O2	-0.7 (4)
N2—N1—C7—C9	176.9 (3)	C9—O2—C12—C11	1.4 (4)
N1—N2—C8—C5	177.2 (3)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1W $\cdots$ O1 <sup>i</sup>	0.97 (5)	1.90 (5)	2.865 (4)	174 (5)

---

O1 <i>W</i> —H2 <i>W</i> ···O1 <sup>ii</sup>	0.92 (6)	1.97 (6)	2.873 (4)	168 (5)
N1—H1 <i>A</i> ···O2	0.86	2.36	2.715 (4)	106
N1—H1 <i>A</i> ···O1 <i>W</i>	0.86	2.05	2.886 (4)	165
C8—H8 <i>A</i> ···O1 <i>W</i>	0.93	2.49	3.290 (5)	144

---

Symmetry codes: (i)  $-x, y-1/2, -z+2$ ; (ii)  $-x-1, y-1/2, -z+2$ .