

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(2-Chlorophenyl)succinimide

 B. S. Saraswathi,^a B. Thimme Gowda,^{a*} Sabine Foro^b and Hartmut Fuess^b
^aDepartment of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 11 March 2010; accepted 20 March 2010

 Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.025; wR factor = 0.072; data-to-parameter ratio = 12.5.

In the title compound, $\text{C}_{10}\text{H}_8\text{ClNO}_2$, the dihedral angle between the aromatic benzene ring and the imide segment is 69.5 (1)°. In the crystal structure, molecules are linked by very weak $\text{C}-\text{H}\cdots\pi$ interactions along the [001] direction.

Related literature

For our study of the effect of ring and side-chain substitutions on the structures of this class of compounds, see: Gowda *et al.* (2007); Saraswathi *et al.* (2010a,b).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_8\text{ClNO}_2$
 $M_r = 209.62$

 Orthorhombic, $Pca2_1$
 $a = 10.616$ (1) Å

 $b = 11.191$ (2) Å

 $c = 8.220$ (1) Å

 $V = 976.6$ (2) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.36$ mm⁻¹
 $T = 299$ K

 $0.50 \times 0.48 \times 0.40$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector

 Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)

 $T_{\min} = 0.840$, $T_{\max} = 0.869$

2417 measured reflections

1600 independent reflections

 1500 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.072$
 $S = 1.04$

1600 reflections

128 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Absolute structure: Flack (1983),

525 Friedel pairs

Flack parameter: 0.01 (7)

Table 1

 $\text{C}-\text{H}\cdots\pi$ interaction geometry (Å, °).

 Cg1 is the centroid of the C1–C6 ring.

D—H \cdots A	D—H	H \cdots A	D \cdots A	D—H \cdots A
C3—H3 \cdots Cg1 ⁱ	0.93	2.93	3.76 (2)	149

 Symmetry code: (i) $-x + 1, -y, z + \frac{1}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

BSS thanks the University Grants Commission, Government of India, New Delhi, for the award of a research fellowship under its faculty improvement program.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2270).

References

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Gowda, B. T., Kozisek, J., Svoboda, I. & Fuess, H. (2007). *Z. Naturforsch. Teil A*, **62**, 91–100.
 Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
 Saraswathi, B. S., Gowda, B. T., Foro, S. & Fuess, H. (2010a). *Acta Cryst.* **E66**, o325.
 Saraswathi, B. S., Gowda, B. T., Foro, S. & Fuess, H. (2010b). *Acta Cryst.* **E66**, o390.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2010). E66, o921 [doi:10.1107/S1600536810010457]

***N*-(2-Chlorophenyl)succinimide**

B. S. Saraswathi, B. Thimme Gowda, Sabine Foro and Hartmut Fuess

S1. Comment

The amide moiety is an important constituent of many biologically significant compounds. As a part of studying the effect of ring and side chain substitutions on the structures of this class of compounds (Gowda *et al.*, 2007; Saraswathi *et al.*, 2010*a,b*), the crystal structure of *N,N*-(2-chlorophenyl)succinimide has been determined (Fig. 1). In the structure of the title compound, C₁₀H₈ClNO₂, the molecule is non-planar with the benzene and pyrrolidine rings tilted by 69.5 (1)° with respect to one another. In the crystal structure, the molecules are linked by weak C—H···π interactions.


S2. Experimental

The solution of succinic anhydride (2.5 g) in toluene (25 ml) was treated dropwise with the solution of 2-chloroaniline (2.5 g) also in toluene (20 ml) with constant stirring. The resulting mixture was stirred for about one hour and set aside for an additional hour at room temperature for completion of the reaction. The mixture was then treated with dilute hydrochloric acid to remove the unreacted 2-chloroaniline. The resultant solid *N*-(2-chlorophenyl)succinamic acid was filtered under suction and washed thoroughly with water to remove the unreacted succinic anhydride and succinic acid. It was recrystallized to constant melting point from ethanol. *N*-(2-chlorophenyl)succinamic acid was then heated for 2 hours and then allowed to cool slowly to room temperature to get crystals of *N*-(2-chlorophenyl)succinimide. The purity of the compound was checked and characterized by its infrared spectra.


The prism like colourless single crystals of the compound used in X-ray diffraction studies were grown in ethanolic solution by a slow evaporation at room temperature.

S3. Refinement

The H atoms were positioned with idealized geometry using a riding model with C—H in the range 0.93–0.97 Å. U_{iso}(H) values were set equal to 1.2U_{eq}(parent atom).

**Figure 1**

Molecular structure of the title compound, showing the atom labelling scheme. The displacement ellipsoids are drawn at the 50% probability level. The H atoms are represented as small spheres of arbitrary radii.

**Figure 2**

Molecular packing of the title compound.

***N*-(2-Chlorophenyl)succinimide**

Crystal data

$C_{10}H_8ClNO_2$

$M_r = 209.62$

Orthorhombic, $Pca2_1$

Hall symbol: $P\ 2c\ -2ac$

$a = 10.616\ (1)\ \text{\AA}$

$b = 11.191\ (2)\ \text{\AA}$

$c = 8.220 (1) \text{ \AA}$
 $V = 976.6 (2) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 432$
 $D_x = 1.426 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1700 reflections
 $\theta = 2.6\text{--}27.7^\circ$
 $\mu = 0.36 \text{ mm}^{-1}$
 $T = 299 \text{ K}$
 Prism, colourless
 $0.50 \times 0.48 \times 0.40 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur
 diffractometer with a Sapphire CCD detector
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Rotation method data acquisition using ω and ϕ
 scans
 Absorption correction: multi-scan
 (CrysAlis RED; Oxford Diffraction, 2009)
 $T_{\min} = 0.840$, $T_{\max} = 0.869$

2417 measured reflections
 1600 independent reflections
 1500 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$
 $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.6^\circ$
 $h = -6 \rightarrow 13$
 $k = -13 \rightarrow 8$
 $l = -7 \rightarrow 10$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.072$
 $S = 1.04$
 1600 reflections
 128 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0446P)^2 + 0.1451P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.036 (3)
 Absolute structure: Flack (1983), 525 Friedel
 pairs
 Absolute structure parameter: 0.01 (7)

Special details

Experimental. (CrysAlis RED; Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.40098 (5)	0.22415 (4)	0.92831 (9)	0.0664 (2)
O1	0.77333 (15)	0.38146 (14)	0.8513 (2)	0.0691 (5)
O2	0.42907 (15)	0.30911 (14)	0.5374 (2)	0.0625 (4)
N1	0.60155 (12)	0.31896 (14)	0.70608 (18)	0.0386 (4)

C1	0.61026 (16)	0.19473 (17)	0.7438 (2)	0.0388 (4)
C2	0.52257 (16)	0.14047 (17)	0.8445 (3)	0.0438 (4)
C3	0.5306 (2)	0.02002 (18)	0.8789 (3)	0.0507 (5)
H3	0.4708	-0.0160	0.9455	0.061*
C4	0.6275 (2)	-0.04682 (19)	0.8141 (3)	0.0529 (5)
H4	0.6333	-0.1279	0.8375	0.063*
C5	0.71622 (19)	0.00642 (19)	0.7144 (3)	0.0540 (5)
H5	0.7817	-0.0388	0.6712	0.065*
C6	0.70742 (17)	0.12731 (18)	0.6789 (3)	0.0477 (5)
H6	0.7668	0.1631	0.6116	0.057*
C7	0.68670 (17)	0.40365 (17)	0.7621 (2)	0.0437 (4)
C8	0.65005 (18)	0.52249 (17)	0.6916 (3)	0.0481 (5)
H8A	0.7161	0.5527	0.6211	0.058*
H8B	0.6348	0.5802	0.7773	0.058*
C9	0.53041 (17)	0.49950 (18)	0.5953 (3)	0.0495 (5)
H9A	0.4599	0.5421	0.6428	0.059*
H9B	0.5405	0.5251	0.4833	0.059*
C10	0.50941 (17)	0.36713 (18)	0.6038 (3)	0.0429 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0671 (4)	0.0458 (3)	0.0864 (4)	-0.0093 (2)	0.0334 (3)	-0.0081 (3)
O1	0.0631 (9)	0.0671 (10)	0.0771 (11)	-0.0155 (8)	-0.0345 (9)	0.0026 (8)
O2	0.0574 (8)	0.0553 (9)	0.0749 (11)	-0.0085 (7)	-0.0267 (8)	0.0040 (8)
N1	0.0364 (7)	0.0401 (8)	0.0394 (9)	-0.0075 (6)	-0.0032 (7)	0.0028 (7)
C1	0.0378 (8)	0.0416 (9)	0.0370 (10)	-0.0080 (7)	-0.0070 (7)	0.0018 (8)
C2	0.0422 (9)	0.0431 (10)	0.0461 (11)	-0.0087 (8)	0.0032 (8)	-0.0038 (8)
C3	0.0550 (11)	0.0439 (10)	0.0531 (13)	-0.0117 (9)	0.0033 (10)	0.0039 (9)
C4	0.0633 (12)	0.0419 (10)	0.0535 (12)	-0.0019 (9)	-0.0108 (10)	0.0058 (10)
C5	0.0504 (11)	0.0549 (11)	0.0565 (13)	0.0114 (9)	-0.0025 (10)	0.0015 (10)
C6	0.0408 (9)	0.0567 (11)	0.0457 (11)	-0.0003 (8)	-0.0001 (9)	0.0052 (10)
C7	0.0433 (9)	0.0465 (10)	0.0414 (10)	-0.0120 (8)	0.0020 (8)	-0.0039 (8)
C8	0.0518 (10)	0.0430 (10)	0.0495 (11)	-0.0090 (9)	0.0068 (9)	-0.0039 (9)
C9	0.0443 (11)	0.0441 (10)	0.0601 (12)	0.0007 (8)	0.0014 (9)	0.0021 (10)
C10	0.0401 (9)	0.0447 (10)	0.0439 (10)	-0.0024 (8)	0.0001 (9)	0.0005 (8)

Geometric parameters (\AA , $^\circ$)

Cl1—C2	1.737 (2)	C4—H4	0.9300
O1—C7	1.202 (2)	C5—C6	1.387 (3)
O2—C10	1.203 (2)	C5—H5	0.9300
N1—C7	1.388 (2)	C6—H6	0.9300
N1—C10	1.398 (2)	C7—C8	1.502 (3)
N1—C1	1.427 (2)	C8—C9	1.519 (3)
C1—C6	1.385 (3)	C8—H8A	0.9700
C1—C2	1.386 (3)	C8—H8B	0.9700
C2—C3	1.380 (3)	C9—C10	1.500 (3)

C3—C4	1.379 (3)	C9—H9A	0.9700
C3—H3	0.9300	C9—H9B	0.9700
C4—C5	1.383 (3)		
C7—N1—C10	113.08 (16)	C5—C6—H6	120.0
C7—N1—C1	123.41 (15)	O1—C7—N1	124.02 (19)
C10—N1—C1	123.47 (14)	O1—C7—C8	128.07 (18)
C6—C1—C2	119.44 (18)	N1—C7—C8	107.91 (16)
C6—C1—N1	119.70 (16)	C7—C8—C9	105.54 (15)
C2—C1—N1	120.86 (17)	C7—C8—H8A	110.6
C3—C2—C1	120.59 (18)	C9—C8—H8A	110.6
C3—C2—C11	119.42 (15)	C7—C8—H8B	110.6
C1—C2—C11	119.99 (15)	C9—C8—H8B	110.6
C4—C3—C2	119.79 (19)	H8A—C8—H8B	108.8
C4—C3—H3	120.1	C10—C9—C8	105.50 (16)
C2—C3—H3	120.1	C10—C9—H9A	110.6
C3—C4—C5	120.20 (19)	C8—C9—H9A	110.6
C3—C4—H4	119.9	C10—C9—H9B	110.6
C5—C4—H4	119.9	C8—C9—H9B	110.6
C4—C5—C6	119.92 (19)	H9A—C9—H9B	108.8
C4—C5—H5	120.0	O2—C10—N1	124.13 (18)
C6—C5—H5	120.0	O2—C10—C9	128.10 (19)
C1—C6—C5	120.06 (18)	N1—C10—C9	107.77 (16)
C1—C6—H6	120.0		
C7—N1—C1—C6	-69.2 (2)	C4—C5—C6—C1	0.2 (3)
C10—N1—C1—C6	108.2 (2)	C10—N1—C7—O1	-179.95 (19)
C7—N1—C1—C2	110.9 (2)	C1—N1—C7—O1	-2.3 (3)
C10—N1—C1—C2	-71.7 (2)	C10—N1—C7—C8	-0.1 (2)
C6—C1—C2—C3	-0.8 (3)	C1—N1—C7—C8	177.56 (16)
N1—C1—C2—C3	179.19 (17)	O1—C7—C8—C9	-177.3 (2)
C6—C1—C2—C11	179.40 (15)	N1—C7—C8—C9	2.9 (2)
N1—C1—C2—C11	-0.7 (2)	C7—C8—C9—C10	-4.4 (2)
C1—C2—C3—C4	0.8 (3)	C7—N1—C10—O2	177.3 (2)
C11—C2—C3—C4	-179.36 (17)	C1—N1—C10—O2	-0.3 (3)
C2—C3—C4—C5	-0.3 (3)	C7—N1—C10—C9	-2.8 (2)
C3—C4—C5—C6	-0.2 (3)	C1—N1—C10—C9	179.55 (16)
C2—C1—C6—C5	0.2 (3)	C8—C9—C10—O2	-175.7 (2)
N1—C1—C6—C5	-179.71 (18)	C8—C9—C10—N1	4.4 (2)
