

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Bis(1,3-dibenzylimidazolium) $\mu$ -oxido-bis[trichloridoferrate(III)]

E. M. Mutambi,\* C. J. Adams and A. G. Orpen

University of Bristol, School of Chemistry, Cantock's Close, Bristol BS8 1TS, England  
Correspondence e-mail: chemm@bristol.ac.uk


Received 7 June 2010; accepted 21 June 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  
R factor = 0.033;  $wR$  factor = 0.081; data-to-parameter ratio = 25.2.

In the title compound  $(\text{C}_{17}\text{H}_{17}\text{N}_2)_2[\text{Fe}_2\text{Cl}_6\text{O}]$ , obtained from the solid-state reaction of  $\text{FeCl}_2$  and  $N,N'$ -dibenzylimidazolium chloride, the complex anion has approximate  $D_{3d}$  symmetry with crystallographically imposed inversion symmetry coincident with the bridging  $\mu$ -O atom. The stereochemistry about each  $\text{FeCl}_3\text{O}$  centre is distorted tetrahedral [ $\text{Fe}-\text{Cl} = 2.2176(5)$ – $2.2427(5)$  Å and  $\text{Fe}-\text{O} = 1.7545(2)$  Å]. The Cl atoms are involved in weak anion-cation  $\text{C}-\text{H}\cdots\text{Cl}$  interactions, giving a network structure.

## Related literature

For literature relating to the intended product, see: Yoshida *et al.* (2005); Zhong *et al.* (2007). For literature relating to anions see: Molins *et al.* (1998); Kohn *et al.* (1996); Vasilevsky *et al.* (1988).



## Experimental

### Crystal data

 $(\text{C}_{17}\text{H}_{17}\text{N}_2)_2[\text{Fe}_2\text{Cl}_6\text{O}]$  $M_r = 839.05$ Orthorhombic,  $Pbca$  $a = 16.2468(5)$  Å $b = 12.8841(4)$  Å $c = 17.6041(5)$  Å $V = 3684.99(19)$  Å<sup>3</sup> $Z = 4$ Mo  $K\alpha$  radiation $\mu = 1.26$  mm<sup>-1</sup> $T = 100$  K $0.31 \times 0.21 \times 0.16$  mm

### Data collection

Bruker SMART CCD area-detector  
diffractometerAbsorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996) $T_{\min} = 0.681$ ,  $T_{\max} = 0.815$ 

42672 measured reflections

5399 independent reflections

4111 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.052$ 

### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.033$  $wR(F^2) = 0.081$  $S = 1.04$ 

5399 reflections

214 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.77$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.44$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}-\text{H1}\cdots\text{Cl1}^{\text{i}}$	0.95	2.82	3.6579 (16)	147
$\text{C3}-\text{H3}\cdots\text{Cl1}$	0.95	2.81	3.4853 (17)	129
$\text{C11}-\text{H11A}\cdots\text{Cl1}^{\text{i}}$	0.99	2.85	3.7440 (18)	151
$\text{C13}-\text{H13}\cdots\text{Cl2}^{\text{ii}}$	0.95	2.86	3.5619 (18)	131
$\text{C15}-\text{H15}\cdots\text{Cl2}^{\text{iii}}$	0.95	2.91	3.8403 (18)	167
$\text{C2}-\text{H2}\cdots\text{Cl3}$	0.95	2.91	3.8223 (17)	162

Symmetry codes: (i)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $-x + \frac{3}{2}, -y, z - \frac{1}{2}$ ; (iii)  $x, -y - \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXL97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

EMM thanks the University of Bristol and Overseas Research Scholarship Awards Scheme (ORSAS) for funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2046).

## References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Kohn, R. D., Seifert, G. & Kociok-Kohn, G. (1996). *Angew. Chem. Int. Ed.* **35**, 2879–2881.
- Molins, E., Roig, A., Miravittles, C., Moreno-Manas, M., Vallribera, A., Galvez, N. & Serra, N. (1998). *Struct. Chem.* **9**, 203–208.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Vasilevsky, I., Stenkamp, R. E., Lingafelter, E. C. & Rose, N. J. (1988). *J. Coord. Chem.* **19**, 171–187.
- Yoshida, Y., Otsuka, A., Saito, G., Natsume, S., Nishibori, E., Takata, M., Sakata, M., Takahashi, M. & Yoko, T. (2005). *Bull. Chem. Soc. Jpn.* **78**, 1921–1928.
- Zhong, C., Sasaki, T., Jimbo-Kobayashi, A., Fujiwara, E., Kobayashi, A., Tada, M. & Iwasawa, Y. (2007). *Bull. Chem. Soc. Jpn.* **80**, 2365–2374.

## supporting information

*Acta Cryst.* (2010). E66, m846 [doi:10.1107/S1600536810024098]

**Bis(1,3-dibenzylimidazolium)  $\mu$ -oxido-bis[trichloridoferrate(III)]**

**E. M. Mutambi, C. J. Adams and A. G. Orpen**

**S1. Comment**

We sought to widen the synthetic routes for the synthesis of  $[\text{FeCl}_4]^-$ - $[\text{HIBz}]^+$  compounds (IBz = *N,N'*-dibenzylimidazole) by using solid state reactions without resorting to solution methods. The aim was to form the salts by grinding together the reactants and this was done with  $\text{FeCl}_2$  and  $\text{IBz}\cdot\text{HCl}$  resulting in formation of the title compound ( $\text{C}_{17}\text{H}_{17}\text{N}_2$ )<sub>2</sub>  $[\text{Cl}_6\text{Fe}_2\text{O}]$  (I), which was obtained during the recrystallization process after extracting with dichloromethane and the structure is reported here. The asymmetric unit of (I) contains one  $\text{HIBz}^+$  cation and half of  $[\text{Fe}_2\text{OCl}_6]^{2-}$  dianion has  $D_{3d}$  symmetry with crystallographically imposed inversion symmetry coincident with the bridging O atom (Fig. 1). The stereochemistry about each  $\text{FeCl}_3\text{O}$  centre is tetrahedral [ $\text{Fe}-\text{Cl}$ , 2.2176 (5)–2.2427 (5) Å;  $\text{Fe}-\text{O}$ , 1.7545 (2) Å]. The  $\text{Fe}-\text{O}$  bond distance compares very well with the average bond length of 1.758 Å for similar anions reported in the CSD. However, the  $\text{Fe}-\text{Cl}$  bond distances are slightly longer compared to the mean  $\text{Fe}-\text{Cl}$  distance of 2.181 Å in the tetrahedral  $\text{FeCl}_4$ , but agree well with the average distance of 2.214 Å reported in the CSD for anions similar to (I). In the cation the phenyl rings lie approximately perpendicular to the plane of the HIBz ring [torsion angles  $\text{C}1-\text{N}1-\text{C}4-\text{C}5$ , 86.90 (19)°;  $\text{C}1-\text{N}2-\text{C}11-\text{C}12$ , 109.17 (18)°].


The crystal structure shows weak cation–anion  $\text{C}-\text{H}\cdots\text{Cl}$  hydrogen-bonding interactions (Table 1, Fig. 2), giving a network structure (Figure 3).

**S2. Experimental**

$\text{FeCl}_2$  (0.0299 g; 0.1 mmol) and *N,N'*-dibenzylimidazolium chloride (0.0858 g; 0.2 mmol) were ground together for *ca.* 10 minutes using a mortar and pestle. A small amount of the crystalline product was dissolved in dichloromethane and allowed to slowly evaporate at room temperature, giving after 3 days, yellow plate-like crystals of (I) suitable for X-ray diffraction.


**S3. Refinement**

H atoms were positioned geometrically and refined using a riding model with  $\text{C}-\text{H} = 0.93$  Å (aromatic) or 0.99 Å (aliphatic) and  $U_{\text{iso}}(\text{H}) = 1.2$  times  $U_{\text{eq}}(\text{C}, \text{N})$ .




**Figure 1**

Molecular structure and atom numbering scheme for the IBz cation and the complex dianion in (I). The dianion lies across an inversion centre: for primed atoms, the symmetry code is  $-x + 1, -y, -z + 2$ . Non-H atoms are drawn at the 50% probability level.



**Figure 2**

Cation–anion C—H···Cl interactions in (I).

**Figure 3**

Crystal packing showing the three-dimensional networks formed in (I).

### Bis(1,3-dibenzylimidazolium) $\mu$ -oxido-bis[trichloridoferrate(III)]

#### Crystal data

$(C_{17}H_{17}N_2)_2[Fe_2Cl_6O]$

$M_r = 839.05$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 16.2468$  (5) Å

$b = 12.8841$  (4) Å

$c = 17.6041$  (5) Å

$V = 3684.99$  (19) Å<sup>3</sup>

$Z = 4$

$F(000) = 1712$

$D_x = 1.512$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 8129 reflections

$\theta = 2.3$ – $29.0^\circ$

$\mu = 1.26$  mm<sup>-1</sup>

$T = 100$  K

Plate, yellow

$0.31 \times 0.21 \times 0.16$  mm

#### Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: sealed tube

Graphite monochromator

Detector resolution: 9.091 pixels mm<sup>-1</sup>

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.681$ ,  $T_{\max} = 0.815$

42672 measured reflections

5399 independent reflections

4111 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 30.1^\circ$ ,  $\theta_{\min} = 3.4^\circ$

$h = -22 \rightarrow 22$

$k = -10 \rightarrow 18$

$l = -24 \rightarrow 24$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.081$   
 $S = 1.04$   
 5399 reflections  
 214 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0331P)^2 + 1.977P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.77 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.572901 (17)	0.002174 (18)	0.926492 (14)	0.01778 (7)
N1	0.69781 (9)	0.34450 (10)	0.70045 (8)	0.0158 (3)
N2	0.60555 (9)	0.24363 (10)	0.65187 (8)	0.0162 (3)
C1	0.65919 (11)	0.31864 (12)	0.63681 (9)	0.0168 (3)
H1	0.6684	0.3489	0.5883	0.020*
C2	0.60972 (11)	0.22098 (13)	0.72855 (9)	0.0191 (3)
H2	0.5782	0.1704	0.7550	0.023*
C3	0.66719 (11)	0.28445 (13)	0.75867 (9)	0.0191 (3)
H3	0.6835	0.2872	0.8105	0.023*
C4	0.75812 (11)	0.42987 (12)	0.70917 (9)	0.0180 (3)
H4A	0.7904	0.4374	0.6618	0.022*
H4B	0.7967	0.4132	0.7509	0.022*
C5	0.71438 (11)	0.53029 (12)	0.72638 (9)	0.0158 (3)
C6	0.70433 (11)	0.56215 (13)	0.80122 (10)	0.0192 (3)
H6	0.7271	0.5220	0.8413	0.023*
C7	0.66112 (12)	0.65228 (14)	0.81769 (11)	0.0255 (4)
H7	0.6553	0.6744	0.8689	0.031*
C8	0.62652 (12)	0.70995 (14)	0.75939 (12)	0.0283 (4)
H8	0.5958	0.7708	0.7706	0.034*
C9	0.63691 (13)	0.67870 (14)	0.68495 (12)	0.0289 (4)
H9	0.6136	0.7186	0.6450	0.035*
C10	0.68116 (12)	0.58936 (14)	0.66809 (10)	0.0226 (4)
H10	0.6887	0.5688	0.6167	0.027*
C11	0.54927 (11)	0.19507 (13)	0.59652 (10)	0.0182 (3)
H11A	0.5494	0.2364	0.5491	0.022*

H11B	0.4926	0.1961	0.6173	0.022*
C12	0.57281 (11)	0.08439 (12)	0.57817 (9)	0.0159 (3)
C13	0.64112 (11)	0.06417 (13)	0.53308 (9)	0.0185 (3)
H13	0.6745	0.1199	0.5158	0.022*
C14	0.66090 (12)	-0.03689 (14)	0.51314 (9)	0.0209 (4)
H14	0.7082	-0.0503	0.4829	0.025*
C15	0.61153 (12)	-0.11859 (13)	0.53742 (10)	0.0219 (4)
H15	0.6244	-0.1877	0.5228	0.026*
C16	0.54355 (12)	-0.09921 (14)	0.58294 (10)	0.0213 (4)
H16	0.5102	-0.1551	0.5999	0.026*
C17	0.52421 (12)	0.00244 (13)	0.60382 (10)	0.0184 (3)
H17	0.4780	0.0157	0.6354	0.022*
Cl1	0.63474 (3)	0.15767 (3)	0.93098 (2)	0.02198 (10)
Cl2	0.66715 (3)	-0.12117 (3)	0.94427 (3)	0.02821 (11)
Cl3	0.51889 (3)	-0.02728 (3)	0.81283 (3)	0.02493 (10)
O4	0.5000	0.0000	1.0000	0.0421 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Fe1	0.02114 (14)	0.01791 (12)	0.01430 (12)	-0.00244 (9)	0.00495 (10)	0.00096 (8)
N1	0.0180 (8)	0.0140 (6)	0.0153 (6)	0.0019 (5)	0.0011 (6)	-0.0013 (5)
N2	0.0190 (8)	0.0135 (6)	0.0160 (6)	0.0021 (5)	0.0017 (6)	-0.0007 (5)
C1	0.0204 (9)	0.0156 (7)	0.0144 (7)	0.0025 (6)	0.0018 (7)	0.0000 (6)
C2	0.0234 (10)	0.0164 (8)	0.0174 (8)	0.0015 (7)	0.0040 (7)	0.0035 (6)
C3	0.0252 (10)	0.0170 (8)	0.0151 (7)	0.0038 (7)	0.0017 (7)	0.0026 (6)
C4	0.0169 (9)	0.0170 (8)	0.0202 (8)	-0.0002 (6)	0.0008 (7)	-0.0017 (6)
C5	0.0147 (8)	0.0145 (7)	0.0182 (8)	-0.0022 (6)	0.0011 (6)	0.0006 (6)
C6	0.0189 (9)	0.0187 (8)	0.0200 (8)	-0.0023 (6)	0.0013 (7)	0.0001 (6)
C7	0.0246 (10)	0.0211 (9)	0.0308 (10)	-0.0048 (7)	0.0083 (8)	-0.0076 (7)
C8	0.0206 (10)	0.0137 (8)	0.0506 (12)	-0.0014 (7)	0.0110 (9)	-0.0004 (8)
C9	0.0252 (11)	0.0218 (9)	0.0397 (11)	0.0013 (7)	0.0011 (9)	0.0168 (8)
C10	0.0251 (10)	0.0222 (9)	0.0204 (8)	-0.0006 (7)	0.0022 (7)	0.0058 (6)
C11	0.0173 (9)	0.0170 (8)	0.0202 (8)	0.0001 (6)	-0.0016 (7)	-0.0014 (6)
C12	0.0179 (9)	0.0155 (7)	0.0144 (7)	0.0020 (6)	-0.0029 (7)	-0.0002 (5)
C13	0.0191 (9)	0.0206 (8)	0.0159 (7)	-0.0007 (7)	-0.0012 (7)	0.0012 (6)
C14	0.0224 (10)	0.0255 (9)	0.0148 (8)	0.0065 (7)	-0.0015 (7)	-0.0010 (6)
C15	0.0297 (11)	0.0196 (8)	0.0163 (8)	0.0058 (7)	-0.0071 (7)	-0.0024 (6)
C16	0.0250 (10)	0.0182 (8)	0.0208 (8)	-0.0020 (7)	-0.0053 (7)	0.0027 (6)
C17	0.0189 (9)	0.0203 (8)	0.0160 (7)	0.0015 (7)	-0.0016 (7)	0.0006 (6)
Cl1	0.0318 (3)	0.01609 (19)	0.01806 (19)	-0.00222 (16)	0.00283 (17)	-0.00090 (14)
Cl2	0.0343 (3)	0.0168 (2)	0.0335 (2)	-0.00020 (18)	-0.0127 (2)	0.00265 (16)
Cl3	0.0263 (2)	0.0239 (2)	0.0246 (2)	0.00011 (17)	-0.00685 (19)	-0.00070 (16)
O4	0.0488 (15)	0.0435 (13)	0.0342 (12)	-0.0089 (10)	0.0261 (11)	-0.0015 (9)

*Geometric parameters (Å, °)*

Fe1—O4	1.7545 (2)	C7—H7	0.9500
Fe1—Cl3	2.2176 (5)	C8—C9	1.381 (3)
Fe1—Cl2	2.2290 (5)	C8—H8	0.9500
Fe1—Cl1	2.2427 (5)	C9—C10	1.389 (3)
N1—C1	1.327 (2)	C9—H9	0.9500
N1—C3	1.377 (2)	C10—H10	0.9500
N1—C4	1.481 (2)	C11—C12	1.511 (2)
N2—C1	1.328 (2)	C11—H11A	0.9900
N2—C2	1.383 (2)	C11—H11B	0.9900
N2—C11	1.476 (2)	C12—C13	1.389 (2)
C1—H1	0.9500	C12—C17	1.394 (2)
C2—C3	1.350 (2)	C13—C14	1.386 (2)
C2—H2	0.9500	C13—H13	0.9500
C3—H3	0.9500	C14—C15	1.391 (3)
C4—C5	1.507 (2)	C14—H14	0.9500
C4—H4A	0.9900	C15—C16	1.387 (3)
C4—H4B	0.9900	C15—H15	0.9500
C5—C10	1.387 (2)	C16—C17	1.396 (2)
C5—C6	1.390 (2)	C16—H16	0.9500
C6—C7	1.387 (2)	C17—H17	0.9500
C6—H6	0.9500	O4—Fe1 <sup>i</sup>	1.7545 (2)
C7—C8	1.386 (3)		
O4—Fe1—Cl3	113.304 (18)	C9—C8—C7	119.78 (17)
O4—Fe1—Cl2	110.417 (18)	C9—C8—H8	120.1
O4—Fe1—Cl1	106.901 (16)	C7—C8—H8	120.1
Cl3—Fe1—Cl1	111.221 (19)	C8—C9—C10	120.49 (17)
Cl2—Fe1—Cl1	108.92 (2)	C8—C9—H9	119.8
C1—N1—C3	108.45 (14)	C10—C9—H9	119.8
C1—N1—C4	125.94 (14)	C5—C10—C9	119.86 (17)
C3—N1—C4	125.39 (14)	C5—C10—H10	120.1
C1—N2—C2	108.44 (14)	C9—C10—H10	120.1
C1—N2—C11	125.69 (14)	N2—C11—C12	112.61 (14)
C2—N2—C11	125.84 (15)	N2—C11—H11A	109.1
N1—C1—N2	108.93 (14)	C12—C11—H11A	109.1
N1—C1—H1	125.5	N2—C11—H11B	109.1
N2—C1—H1	125.5	C12—C11—H11B	109.1
C3—C2—N2	106.84 (15)	H11A—C11—H11B	107.8
C3—C2—H2	126.6	C13—C12—C17	119.72 (15)
N2—C2—H2	126.6	C13—C12—C11	120.09 (15)
C2—C3—N1	107.33 (15)	C17—C12—C11	120.14 (16)
C2—C3—H3	126.3	C14—C13—C12	120.40 (16)
N1—C3—H3	126.3	C14—C13—H13	119.8
N1—C4—C5	110.28 (14)	C12—C13—H13	119.8
N1—C4—H4A	109.6	C13—C14—C15	119.96 (17)
C5—C4—H4A	109.6	C13—C14—H14	120.0



N1—C4—H4B	109.6	C15—C14—H14	120.0
C5—C4—H4B	109.6	C16—C15—C14	120.04 (16)
H4A—C4—H4B	108.1	C16—C15—H15	120.0
C10—C5—C6	119.58 (16)	C14—C15—H15	120.0
C10—C5—C4	120.39 (15)	C15—C16—C17	120.01 (17)
C6—C5—C4	119.98 (15)	C15—C16—H16	120.0
C7—C6—C5	120.31 (17)	C17—C16—H16	120.0
C7—C6—H6	119.8	C12—C17—C16	119.86 (17)
C5—C6—H6	119.8	C12—C17—H17	120.1
C8—C7—C6	119.95 (17)	C16—C17—H17	120.1
C8—C7—H7	120.0	Fe1—O4—Fe1 <sup>i</sup>	180.0
C6—C7—H7	120.0		
C3—N1—C1—N2	-0.60 (19)	C7—C8—C9—C10	0.6 (3)
C4—N1—C1—N2	-175.46 (14)	C6—C5—C10—C9	-1.2 (3)
C2—N2—C1—N1	0.37 (19)	C4—C5—C10—C9	176.32 (17)
C11—N2—C1—N1	178.63 (14)	C8—C9—C10—C5	0.8 (3)
C1—N2—C2—C3	0.01 (19)	C1—N2—C11—C12	109.77 (18)
C11—N2—C2—C3	-178.25 (15)	C2—N2—C11—C12	-72.3 (2)
N2—C2—C3—N1	-0.37 (19)	N2—C11—C12—C13	-73.19 (19)
C1—N1—C3—C2	0.60 (19)	N2—C11—C12—C17	109.41 (18)
C4—N1—C3—C2	175.50 (15)	C17—C12—C13—C14	0.4 (2)
C1—N1—C4—C5	86.90 (19)	C11—C12—C13—C14	-177.04 (15)
C3—N1—C4—C5	-87.12 (19)	C12—C13—C14—C15	0.9 (3)
N1—C4—C5—C10	-82.77 (19)	C13—C14—C15—C16	-1.4 (3)
N1—C4—C5—C6	94.71 (18)	C14—C15—C16—C17	0.6 (3)
C10—C5—C6—C7	0.2 (3)	C13—C12—C17—C16	-1.2 (3)
C4—C5—C6—C7	-177.29 (16)	C11—C12—C17—C16	176.24 (15)
C5—C6—C7—C8	1.1 (3)	C15—C16—C17—C12	0.7 (3)
C6—C7—C8—C9	-1.5 (3)		

Symmetry code: (i)  $-x+1, -y, -z+2$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 $\cdots$ C11 <sup>ii</sup>	0.95	2.82	3.6579 (16)	147
C3—H3 $\cdots$ C11	0.95	2.81	3.4853 (17)	129
C11—H11A $\cdots$ C11 <sup>ii</sup>	0.99	2.85	3.7440 (18)	151
C13—H13 $\cdots$ C12 <sup>iii</sup>	0.95	2.86	3.5619 (18)	131
C15—H15 $\cdots$ C12 <sup>iv</sup>	0.95	2.91	3.8403 (18)	167
C2—H2 $\cdots$ C13	0.95	2.91	3.8223 (17)	162

Symmetry codes: (ii)  $x, -y+1/2, z-1/2$ ; (iii)  $-x+3/2, -y, z-1/2$ ; (iv)  $x, -y-1/2, z-1/2$ .