

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

9,9-Bis[4-(prop-2-ynoxy)phenyl]-9H-fluorene

 Kiramat Shah,^a Muhammad Raza Shah^a and Seik Weng Ng^{b*}
^aH. E. J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and

^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 30 June 2010; accepted 1 July 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.032; wR factor = 0.082; data-to-parameter ratio = 9.3.

In the title compound, $\text{C}_{31}\text{H}_{22}\text{O}_2$, the bond angle at the C atom belonging to the five-membered ring of the fluorene system is opened to 112.64 (12)°. The two benzene rings are twisted with respect to the fluorene ring system at dihedral angles of 72.81 (6) and 81.83 (6)°. One $\text{C}_{\text{aryl}}-\text{O}-\text{C}\equiv$ fragment is extended, with a $\text{C}-\text{O}-\text{C}-\text{C}$ torsion angle of -178.77 (13)°, but the other $\text{C}_{\text{aryl}}-\text{O}-\text{C}-\text{C}\equiv$ fragment is bent, with a $\text{C}-\text{O}-\text{C}-\text{C}$ torsion angle of 64.78 (19)°. Intermolecular weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonding is present in the crystal structure.

Related literature

For the synthesis of copolyethers having 1,3,4-oxadiazole rings and fluorene groups from the polymerization of 9,9-bis(4-propargyloxyphenyl)fluorene, see: Hamciuc *et al.* (2009). For a related structure, see: Shah *et al.* (2010).



Experimental

Crystal data

$\text{C}_{31}\text{H}_{22}\text{O}_2$	$V = 2203.1$ (3) Å ³
$M_r = 426.49$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 11.0047$ (8) Å	$\mu = 0.08$ mm ⁻¹
$b = 12.9686$ (9) Å	$T = 100$ K
$c = 15.4371$ (11) Å	$0.35 \times 0.25 \times 0.15$ mm

Data collection

Bruker SMART APEX diffractometer	2861 independent reflections
21245 measured reflections	2658 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$	$\Delta\rho_{\text{max}} = 0.18$ e Å ⁻³
$wR(F^2) = 0.082$	$\Delta\rho_{\text{min}} = -0.20$ e Å ⁻³
$S = 1.04$	Absolute structure: 2204 Friedel pairs were merged
2861 reflections	Flack parameter: ?
306 parameters	Rogers parameter: ?
2 restraints	
H atoms treated by a mixture of independent and constrained refinement	

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}29-\text{H}29\text{A}\cdots\text{O}1^i$	0.99	2.46	3.358 (2)	151

 Symmetry code: (i) $-x + 2, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors thank the Higher Education Commission of Pakistan and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2792).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2009). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hamciuc, C., Hamciuc, E., Ipate, A. M., Cristea, M. & Okrasa, L. (2009). *J. Appl. Polym. Sci.* **113**, 383–391.
 Shah, K., Yousuf, S., Raza Shah, M. & Ng, S. W. (2010). *Acta Cryst.* **E66**, o1705.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2010). E66, o1939 [https://doi.org/10.1107/S1600536810025833]

9,9-Bis[4-(prop-2-ynyloxy)phenyl]-9H-fluorene**Kiramat Shah, Muhammad Raza Shah and Seik Weng Ng****S1. Comment**

9,9-Bis(4-propargyloxyphenyl)fluorene is the monomer required for the synthesis of other copolyethers (Hamciuc *et al.*, 2009); these polymers typically possess good solubility in inorganic solvents, high thermal stability and high glass transition temperatures. The compound is readily synthesized from commercially available 9,9-bis(4-hydroxyphenyl)fluorene and propargyl bromide. A previous report detailed the structure of the product of its reaction with *t*-butyl bromoacetate (Shah *et al.*, 2010). In the C₃₁H₂₂O₂ molecule (Scheme I, Fig. 1), the angle at the carbon atom belonging to the five-membered fluorenyl ring that is connected to two *p*-phenylene rings is open to 116.1 (1) ° by the rings. One four-atom C_{aryl}-O-C-C≡ fragment is coplanar with the ring [C—O—C—C torsion angle 1.2 (1) °] but the other is bent [torsion angle 64.8 (2) °].

S2. Experimental

9,9-Bis(4-hydroxyphenyl)fluorene (0.5 g, 1.4 mmol) was dissolved in acetone (25 ml) to give a clear solution. Potassium carbonate (0.7 g, 5 mmol) was added and the mixture stirred for an hour. Propargyl bromide (0.7 ml, 5 mmol) was added and stirring continued overnight. The mixture was filtered; prismatic crystals separated from the solution in 80% yield.

S3. Refinement

The acetylenic H-atoms were located in a difference Fourier map, and were refined with a distance restraint of C—H 0.95±0.01 Å; their temperature factors were refined. Other H atoms were placed in calculated positions [C—H 0.95–0.99 Å, $U(H) = 1.2U(C)$] and were included in the refinement in the riding model approximation. Friedel pairs were merged as no significant anomalous scattering.




Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $C_{31}H_{22}O_2$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

9,9-Bis[4-(prop-2-ynyloxy)phenyl]-9H-fluorene

Crystal data

$C_{31}H_{22}O_2$
 $M_r = 426.49$
 Orthorhombic, $P2_12_12_1$
 Hall symbol: P 2ac 2ab
 $a = 11.0047$ (8) Å
 $b = 12.9686$ (9) Å
 $c = 15.4371$ (11) Å
 $V = 2203.1$ (3) Å³
 $Z = 4$

$F(000) = 896$
 $D_x = 1.286$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 7024 reflections
 $\theta = 2.3$ – 27.7°
 $\mu = 0.08$ mm⁻¹
 $T = 100$ K
 Block, colorless
 $0.35 \times 0.25 \times 0.15$ mm

Data collection

Bruker SMART APEX
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 ω scans
 21245 measured reflections
 2861 independent reflections

2658 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.042$
 $\theta_{max} = 27.5^\circ$, $\theta_{min} = 2.1^\circ$
 $h = -14 \rightarrow 14$
 $k = -16 \rightarrow 16$
 $l = -19 \rightarrow 20$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.082$
 $S = 1.04$
 2861 reflections
 306 parameters
 2 restraints

Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0504P)^2 + 0.3105P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$$

Absolute structure: 2204 Friedel pairs were merged

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.24715 (9)	0.37529 (9)	1.06380 (7)	0.0184 (3)
O2	0.72277 (11)	0.74489 (10)	0.66169 (7)	0.0228 (3)
C1	0.84334 (14)	0.71897 (12)	1.07308 (10)	0.0146 (3)
C2	0.93879 (15)	0.78830 (12)	1.08005 (11)	0.0174 (3)
H2	1.0072	0.7832	1.0428	0.021*
C3	0.93255 (15)	0.86561 (13)	1.14272 (11)	0.0200 (3)
H3	0.9979	0.9130	1.1488	0.024*
C4	0.83115 (16)	0.87398 (13)	1.19668 (11)	0.0205 (3)
H4	0.8278	0.9276	1.2385	0.025*
C5	0.73513 (16)	0.80494 (13)	1.18992 (11)	0.0188 (3)
H5	0.6660	0.8111	1.2264	0.023*
C6	0.74238 (14)	0.72624 (13)	1.12841 (10)	0.0156 (3)
C7	0.65875 (14)	0.64169 (13)	1.10835 (10)	0.0169 (3)
C8	0.54672 (15)	0.61615 (14)	1.14465 (11)	0.0206 (3)
H8	0.5128	0.6565	1.1900	0.025*
C9	0.48567 (15)	0.52999 (15)	1.11285 (12)	0.0226 (4)
H9	0.4097	0.5109	1.1372	0.027*
C10	0.53482 (15)	0.47162 (14)	1.04579 (12)	0.0211 (4)
H10	0.4922	0.4129	1.0251	0.025*
C11	0.64577 (15)	0.49837 (13)	1.00872 (11)	0.0189 (3)
H11	0.6782	0.4593	0.9621	0.023*
C12	0.70800 (14)	0.58268 (13)	1.04078 (10)	0.0157 (3)
C13	0.82912 (13)	0.62794 (12)	1.01042 (10)	0.0143 (3)
C14	0.93668 (14)	0.55407 (12)	1.02154 (10)	0.0145 (3)
C15	0.93030 (15)	0.46837 (13)	1.07548 (11)	0.0167 (3)
H15	0.8553	0.4519	1.1027	0.020*
C16	1.03105 (14)	0.40609 (13)	1.09059 (10)	0.0170 (3)
H16	1.0248	0.3477	1.1275	0.020*
C17	1.14088 (14)	0.43008 (13)	1.05113 (11)	0.0151 (3)
C18	1.14950 (14)	0.51457 (12)	0.99593 (11)	0.0160 (3)
H18	1.2243	0.5303	0.9682	0.019*
C19	1.04803 (14)	0.57562 (12)	0.98172 (10)	0.0160 (3)
H19	1.0542	0.6334	0.9441	0.019*
C20	1.23372 (16)	0.26883 (13)	1.08654 (12)	0.0210 (4)
H20A	1.1907	0.2313	1.0399	0.025*
H20B	1.1859	0.2623	1.1406	0.025*
C21	1.35523 (17)	0.22575 (14)	1.09882 (11)	0.0222 (4)
C22	1.4531 (2)	0.19125 (17)	1.11001 (14)	0.0342 (5)
C23	0.81257 (14)	0.66357 (13)	0.91600 (10)	0.0153 (3)
C24	0.76842 (14)	0.76168 (13)	0.89833 (10)	0.0163 (3)

H24	0.7588	0.8092	0.9447	0.020*
C25	0.73786 (15)	0.79255 (13)	0.81480 (11)	0.0186 (3)
H25	0.7075	0.8600	0.8043	0.022*
C26	0.75237 (15)	0.72339 (14)	0.74710 (10)	0.0182 (3)
C27	0.79946 (15)	0.62612 (15)	0.76239 (11)	0.0210 (4)
H27	0.8119	0.5799	0.7155	0.025*
C28	0.82856 (15)	0.59607 (13)	0.84623 (11)	0.0192 (3)
H28	0.8597	0.5288	0.8563	0.023*
C29	0.67080 (16)	0.84363 (14)	0.64399 (11)	0.0221 (4)
H29A	0.6628	0.8518	0.5805	0.027*
H29B	0.7272	0.8977	0.6649	0.027*
C30	0.55120 (16)	0.86010 (14)	0.68398 (11)	0.0203 (3)
C31	0.45729 (17)	0.87683 (15)	0.71807 (12)	0.0254 (4)
H22	1.5337 (13)	0.167 (2)	1.1183 (19)	0.073 (10)*
H31	0.3818 (13)	0.8907 (17)	0.7453 (13)	0.036 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0141 (5)	0.0153 (6)	0.0257 (6)	0.0011 (5)	0.0006 (5)	0.0027 (5)
O2	0.0289 (6)	0.0249 (6)	0.0146 (5)	0.0056 (5)	-0.0034 (5)	0.0000 (5)
C1	0.0168 (7)	0.0140 (7)	0.0130 (7)	0.0024 (6)	-0.0032 (6)	0.0020 (6)
C2	0.0182 (7)	0.0176 (8)	0.0164 (8)	-0.0001 (6)	-0.0016 (6)	0.0003 (6)
C3	0.0233 (8)	0.0157 (8)	0.0209 (8)	-0.0031 (7)	-0.0052 (7)	0.0008 (7)
C4	0.0292 (8)	0.0168 (8)	0.0154 (7)	0.0034 (7)	-0.0040 (7)	-0.0016 (6)
C5	0.0212 (8)	0.0205 (8)	0.0147 (7)	0.0045 (7)	-0.0007 (6)	0.0001 (7)
C6	0.0161 (7)	0.0167 (8)	0.0140 (7)	0.0031 (6)	-0.0021 (6)	0.0020 (6)
C7	0.0165 (7)	0.0184 (8)	0.0159 (7)	0.0021 (6)	-0.0024 (6)	0.0027 (6)
C8	0.0160 (7)	0.0267 (9)	0.0192 (8)	0.0023 (7)	-0.0003 (6)	0.0019 (7)
C9	0.0140 (7)	0.0283 (9)	0.0256 (9)	-0.0019 (7)	0.0001 (7)	0.0063 (8)
C10	0.0167 (7)	0.0196 (8)	0.0269 (9)	-0.0035 (6)	-0.0053 (7)	0.0031 (7)
C11	0.0171 (7)	0.0181 (8)	0.0215 (8)	0.0007 (6)	-0.0025 (7)	0.0006 (7)
C12	0.0145 (7)	0.0161 (8)	0.0164 (7)	0.0017 (6)	-0.0026 (6)	0.0036 (6)
C13	0.0144 (7)	0.0131 (7)	0.0153 (7)	0.0007 (6)	-0.0004 (6)	0.0003 (6)
C14	0.0146 (7)	0.0151 (8)	0.0138 (7)	0.0007 (6)	-0.0012 (6)	-0.0017 (6)
C15	0.0154 (7)	0.0188 (8)	0.0158 (8)	-0.0021 (6)	0.0008 (6)	0.0007 (6)
C16	0.0186 (7)	0.0163 (8)	0.0159 (8)	-0.0001 (6)	-0.0007 (6)	0.0031 (6)
C17	0.0149 (7)	0.0142 (8)	0.0162 (7)	0.0000 (6)	-0.0018 (6)	-0.0032 (6)
C18	0.0154 (7)	0.0160 (8)	0.0166 (8)	-0.0025 (6)	0.0014 (6)	-0.0009 (6)
C19	0.0192 (7)	0.0126 (7)	0.0160 (8)	-0.0018 (6)	0.0005 (6)	0.0012 (6)
C20	0.0219 (8)	0.0149 (8)	0.0263 (9)	-0.0003 (7)	-0.0036 (7)	0.0033 (7)
C21	0.0292 (9)	0.0181 (8)	0.0195 (8)	0.0046 (7)	0.0028 (7)	0.0010 (7)
C22	0.0337 (10)	0.0411 (12)	0.0279 (10)	0.0200 (10)	0.0063 (9)	0.0081 (9)
C23	0.0137 (7)	0.0172 (8)	0.0151 (7)	-0.0003 (6)	-0.0010 (6)	-0.0002 (6)
C24	0.0173 (7)	0.0151 (8)	0.0165 (7)	0.0005 (6)	-0.0011 (6)	-0.0031 (6)
C25	0.0203 (8)	0.0151 (8)	0.0205 (8)	0.0011 (7)	-0.0041 (7)	0.0005 (7)
C26	0.0172 (7)	0.0227 (8)	0.0149 (7)	-0.0005 (6)	-0.0009 (6)	-0.0002 (7)
C27	0.0224 (8)	0.0235 (9)	0.0169 (8)	0.0036 (7)	-0.0017 (6)	-0.0040 (7)

C28	0.0208 (8)	0.0161 (8)	0.0206 (8)	0.0037 (6)	-0.0023 (7)	-0.0021 (7)
C29	0.0247 (8)	0.0233 (9)	0.0184 (8)	0.0012 (7)	-0.0012 (7)	0.0044 (7)
C30	0.0265 (8)	0.0175 (8)	0.0170 (8)	-0.0020 (7)	-0.0052 (7)	0.0025 (7)
C31	0.0265 (9)	0.0253 (10)	0.0243 (9)	-0.0038 (8)	0.0013 (7)	-0.0011 (8)

Geometric parameters (Å, °)

O1—C17	1.3824 (18)	C14—C19	1.399 (2)
O1—C20	1.432 (2)	C15—C16	1.391 (2)
O2—C26	1.387 (2)	C15—H15	0.9500
O2—C29	1.429 (2)	C16—C17	1.389 (2)
C1—C2	1.387 (2)	C16—H16	0.9500
C1—C6	1.405 (2)	C17—C18	1.391 (2)
C1—C13	1.534 (2)	C18—C19	1.386 (2)
C2—C3	1.395 (2)	C18—H18	0.9500
C2—H2	0.9500	C19—H19	0.9500
C3—C4	1.397 (2)	C20—C21	1.462 (2)
C3—H3	0.9500	C20—H20A	0.9900
C4—C5	1.389 (2)	C20—H20B	0.9900
C4—H4	0.9500	C21—C22	1.179 (3)
C5—C6	1.396 (2)	C22—H22	0.949 (10)
C5—H5	0.9500	C23—C24	1.389 (2)
C6—C7	1.465 (2)	C23—C28	1.399 (2)
C7—C8	1.394 (2)	C24—C25	1.391 (2)
C7—C12	1.403 (2)	C24—H24	0.9500
C8—C9	1.393 (3)	C25—C26	1.386 (2)
C8—H8	0.9500	C25—H25	0.9500
C9—C10	1.392 (3)	C26—C27	1.384 (2)
C9—H9	0.9500	C27—C28	1.389 (2)
C10—C11	1.392 (2)	C27—H27	0.9500
C10—H10	0.9500	C28—H28	0.9500
C11—C12	1.382 (2)	C29—C30	1.469 (2)
C11—H11	0.9500	C29—H29A	0.9900
C12—C13	1.530 (2)	C29—H29B	0.9900
C13—C14	1.532 (2)	C30—C31	1.180 (3)
C13—C23	1.540 (2)	C31—H31	0.949 (10)
C14—C15	1.390 (2)		
C17—O1—C20	116.28 (12)	C14—C15—H15	119.2
C26—O2—C29	117.13 (13)	C16—C15—H15	119.2
C2—C1—C6	120.56 (15)	C17—C16—C15	119.34 (15)
C2—C1—C13	128.68 (14)	C17—C16—H16	120.3
C6—C1—C13	110.75 (13)	C15—C16—H16	120.3
C1—C2—C3	118.85 (15)	O1—C17—C16	124.00 (14)
C1—C2—H2	120.6	O1—C17—C18	115.70 (14)
C3—C2—H2	120.6	C16—C17—C18	120.29 (15)
C2—C3—C4	120.58 (16)	C19—C18—C17	119.46 (15)
C2—C3—H3	119.7	C19—C18—H18	120.3

C4—C3—H3	119.7	C17—C18—H18	120.3
C5—C4—C3	120.86 (15)	C18—C19—C14	121.46 (15)
C5—C4—H4	119.6	C18—C19—H19	119.3
C3—C4—H4	119.6	C14—C19—H19	119.3
C4—C5—C6	118.61 (15)	O1—C20—C21	107.81 (14)
C4—C5—H5	120.7	O1—C20—H20A	110.1
C6—C5—H5	120.7	C21—C20—H20A	110.1
C5—C6—C1	120.52 (15)	O1—C20—H20B	110.1
C5—C6—C7	130.92 (15)	C21—C20—H20B	110.1
C1—C6—C7	108.56 (14)	H20A—C20—H20B	108.5
C8—C7—C12	120.73 (16)	C22—C21—C20	179.0 (2)
C8—C7—C6	130.43 (16)	C21—C22—H22	177 (2)
C12—C7—C6	108.84 (14)	C24—C23—C28	117.77 (15)
C9—C8—C7	118.38 (17)	C24—C23—C13	120.14 (14)
C9—C8—H8	120.8	C28—C23—C13	121.74 (14)
C7—C8—H8	120.8	C23—C24—C25	122.01 (15)
C8—C9—C10	120.73 (16)	C23—C24—H24	119.0
C8—C9—H9	119.6	C25—C24—H24	119.0
C10—C9—H9	119.6	C26—C25—C24	118.99 (15)
C11—C10—C9	120.73 (17)	C26—C25—H25	120.5
C11—C10—H10	119.6	C24—C25—H25	120.5
C9—C10—H10	119.6	C27—C26—O2	115.68 (15)
C12—C11—C10	118.97 (16)	C27—C26—C25	120.28 (15)
C12—C11—H11	120.5	O2—C26—C25	124.04 (15)
C10—C11—H11	120.5	C26—C27—C28	120.06 (16)
C11—C12—C7	120.44 (15)	C26—C27—H27	120.0
C11—C12—C13	128.72 (15)	C28—C27—H27	120.0
C7—C12—C13	110.80 (14)	C27—C28—C23	120.85 (16)
C12—C13—C14	113.50 (13)	C27—C28—H28	119.6
C12—C13—C1	101.01 (12)	C23—C28—H28	119.6
C14—C13—C1	109.36 (12)	O2—C29—C30	114.11 (14)
C12—C13—C23	107.59 (12)	O2—C29—H29A	108.7
C14—C13—C23	112.64 (12)	C30—C29—H29A	108.7
C1—C13—C23	112.21 (13)	O2—C29—H29B	108.7
C15—C14—C19	117.84 (14)	C30—C29—H29B	108.7
C15—C14—C13	121.86 (14)	H29A—C29—H29B	107.6
C19—C14—C13	120.16 (14)	C31—C30—C29	177.1 (2)
C14—C15—C16	121.61 (15)	C30—C31—H31	179.6 (16)
C6—C1—C2—C3	-0.3 (2)	C12—C13—C14—C15	-17.1 (2)
C13—C1—C2—C3	-179.70 (15)	C1—C13—C14—C15	94.84 (17)
C1—C2—C3—C4	-0.9 (2)	C23—C13—C14—C15	-139.66 (15)
C2—C3—C4—C5	0.8 (3)	C12—C13—C14—C19	167.42 (14)
C3—C4—C5—C6	0.5 (2)	C1—C13—C14—C19	-80.65 (17)
C4—C5—C6—C1	-1.7 (2)	C23—C13—C14—C19	44.9 (2)
C4—C5—C6—C7	177.91 (16)	C19—C14—C15—C16	0.7 (2)
C2—C1—C6—C5	1.6 (2)	C13—C14—C15—C16	-174.94 (15)
C13—C1—C6—C5	-178.88 (14)	C14—C15—C16—C17	0.2 (3)

C2—C1—C6—C7	-178.09 (14)	C20—O1—C17—C16	27.2 (2)
C13—C1—C6—C7	1.44 (17)	C20—O1—C17—C18	-153.71 (14)
C5—C6—C7—C8	0.7 (3)	C15—C16—C17—O1	177.96 (15)
C1—C6—C7—C8	-179.64 (16)	C15—C16—C17—C18	-1.1 (2)
C5—C6—C7—C12	-179.73 (16)	O1—C17—C18—C19	-178.06 (14)
C1—C6—C7—C12	-0.10 (18)	C16—C17—C18—C19	1.0 (2)
C12—C7—C8—C9	0.7 (2)	C17—C18—C19—C14	-0.2 (2)
C6—C7—C8—C9	-179.79 (16)	C15—C14—C19—C18	-0.7 (2)
C7—C8—C9—C10	-0.7 (3)	C13—C14—C19—C18	174.99 (14)
C8—C9—C10—C11	-0.3 (3)	C17—O1—C20—C21	-178.77 (13)
C9—C10—C11—C12	1.4 (3)	C12—C13—C23—C24	86.88 (17)
C10—C11—C12—C7	-1.4 (2)	C14—C13—C23—C24	-147.28 (15)
C10—C11—C12—C13	-178.96 (15)	C1—C13—C23—C24	-23.34 (19)
C8—C7—C12—C11	0.3 (2)	C12—C13—C23—C28	-86.21 (17)
C6—C7—C12—C11	-179.24 (14)	C14—C13—C23—C28	39.6 (2)
C8—C7—C12—C13	178.31 (14)	C1—C13—C23—C28	163.57 (15)
C6—C7—C12—C13	-1.28 (18)	C28—C23—C24—C25	1.5 (2)
C11—C12—C13—C14	-63.3 (2)	C13—C23—C24—C25	-171.88 (15)
C7—C12—C13—C14	118.92 (15)	C23—C24—C25—C26	-0.3 (2)
C11—C12—C13—C1	179.74 (16)	C29—O2—C26—C27	178.32 (14)
C7—C12—C13—C1	1.99 (16)	C29—O2—C26—C25	-2.0 (2)
C11—C12—C13—C23	62.0 (2)	C24—C25—C26—C27	-1.6 (2)
C7—C12—C13—C23	-115.75 (15)	C24—C25—C26—O2	178.69 (15)
C2—C1—C13—C12	177.43 (15)	O2—C26—C27—C28	-178.08 (15)
C6—C1—C13—C12	-2.06 (16)	C25—C26—C27—C28	2.2 (3)
C2—C1—C13—C14	57.5 (2)	C26—C27—C28—C23	-0.9 (3)
C6—C1—C13—C14	-121.99 (14)	C24—C23—C28—C27	-0.9 (2)
C2—C1—C13—C23	-68.3 (2)	C13—C23—C28—C27	172.36 (15)
C6—C1—C13—C23	112.26 (14)	C26—O2—C29—C30	-64.78 (19)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C29—H29 <i>A</i> \cdots O1 ⁱ	0.99	2.46	3.358 (2)	151

Symmetry code: (i) $-x+2, y+1/2, -z+3/2$.