

catena-Poly[[[diaquacadmium(II)]-bis[μ -3,5-bis(isonicotinamido)benzoato]] tetrahydrate]

Man-Sheng Chen, Yi-Fang Deng, Chun-Hua Zhang and Dai-Zhi Kuang*

Key Laboratory of Functional Organometallic Materials, Department of Chemistry and Materials Science, Hengyang Normal University, Hengyang, Hunan 421008, People's Republic of China

Correspondence e-mail: cmsniu@163.com


Received 21 July 2010; accepted 8 September 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.047; wR factor = 0.110; data-to-parameter ratio = 12.9.

The title compound, $\{[\text{Cd}(\text{C}_{19}\text{H}_{13}\text{N}_4\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 4\text{H}_2\text{O}\}_n$ or $\{[\text{Cd}(\text{BBA})_2(\text{H}_2\text{O})_2] \cdot 4\text{H}_2\text{O}\}_n$, where BBA is 3,5-bis(isonicotinamido)benzoate, is isotypic with its Mn isologue [Chen *et al.* (2009). *J. Coord. Chem.* **62**, 2421–2428]. The cation sits on a twofold axis and is six-coordinated in a slightly distorted octahedral geometry; the polyhedra are linked into zigzag chains, which are further connected by $\text{N}-\text{H} \cdots \text{O}$, $\text{O}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds as well as $\pi-\pi$ interactions [centroid-centroid distance of 3.639 (2) Å], giving a three-dimensional supramolecular framework.

Related literature

For the isotypic Mn structure, see: Chen *et al.* (2009). For the properties of coordination polymers, see: Evans & Lin (2002); Yaghi *et al.* (2003); Kitagawa *et al.* (2004); Biradha *et al.* (2006); Wu *et al.* (2009). For the rational design and synthesis of new supramolecular frameworks by covalent and weak intra/intermolecular interactions, see: Eddaoudi *et al.* (2001); Moulton & Zaworotko (2001); Cheng *et al.* (2002); Zhang *et al.* (2003); Go *et al.* (2004). For the coordination capacities of carboxylate, pyridine and amide groups, see: Bent (1968); Huyskens (1977); Lee & Kumler (1962); Wang *et al.* (2007).



Experimental

Crystal data

$[\text{Cd}(\text{C}_{19}\text{H}_{13}\text{N}_4\text{O}_4)_2(\text{H}_2\text{O})_2] \cdot 4\text{H}_2\text{O}$
 $M_r = 943.16$
 Monoclinic, $C2/c$
 $a = 17.584$ (3) Å
 $b = 10.8568$ (19) Å
 $c = 21.891$ (4) Å
 $\beta = 103.801$ (2)°

$V = 4058.5$ (12) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.62$ mm⁻¹
 $T = 293$ K
 $0.20 \times 0.16 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.887$, $T_{\max} = 0.941$

10338 measured reflections
 3867 independent reflections
 3391 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.110$
 $S = 1.08$
 3867 reflections
 300 parameters
 13 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.80$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.48$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N}3-\text{H}3 \cdots \text{O}6$	0.86	2.10	2.885 (5)	151
$\text{N}4-\text{H}4 \cdots \text{O}3^i$	0.86	2.26	3.096 (4)	164
$\text{O}5-\text{H}5B \cdots \text{O}2$	0.83 (3)	1.98 (3)	2.710 (4)	145 (4)
$\text{O}5-\text{H}5A \cdots \text{N}1^{ii}$	0.83 (2)	2.00 (2)	2.825 (4)	172 (4)
$\text{O}6-\text{H}6B \cdots \text{O}5^{iii}$	0.84 (2)	2.40 (4)	3.163 (6)	151 (6)
$\text{O}6-\text{H}6A \cdots \text{O}7$	0.82 (3)	1.92 (4)	2.681 (6)	148 (5)
$\text{O}7-\text{H}7A \cdots \text{O}2^{iv}$	0.85 (2)	1.96 (3)	2.767 (4)	162 (5)
$\text{O}7-\text{H}7B \cdots \text{O}4^v$	0.84 (2)	1.99 (3)	2.791 (4)	159 (6)

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $x, -y + 1, z + \frac{1}{2}$; (iv) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z$; (v) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINTE* (Bruker, 2000); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Construct Program of Key Disciplines in Hunan Province, Distinguished Young Cadremans of Hunan Province (2008) and Distinguished Young Cadremans of Hengyang Normal University (2007). We also acknowledge support from the Foundation of Education Department of Hunan Province (10C0473)

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2362).

References

- Bent, H. A. (1968). *Chem. Rev.* **68**, 587–648.
- Biradha, K., Sarkar, M. & Rajput, L. (2006). *Chem. Commun.* pp. 4169–4179.
- Bruker (2000). *SMART*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, M. S., Chen, S. S., Okamura, T.-A., Su, Z., Sun, W. Y. & Ueyama, N. (2009). *J. Coord. Chem.* **62**, 2421–2428.
- Cheng, D. P., Khan, M. A. & Houser, R. P. (2002). *J. Chem. Soc. Dalton Trans.* pp. 4555–4560.
- Eddaoudi, M., Moler, D. B., Li, H. L., Chen, B. L., Reineke, T. M., O’Keeffe, M. & Yaghi, O. M. (2001). *Acc. Chem. Res.* **34**, 319–330.
- Evans, O. R. & Lin, W. (2002). *Acc. Chem. Res.* **35**, 511–522.
- Go, Y. B., Wang, X. Q., Anokhina, E. V. & Jacobson, A. J. (2004). *Inorg. Chem.* **43**, 5360–5367.
- Huyskens, P. L. (1977). *J. Am. Chem. Soc.* **99**, 2578–2582.
- Kitagawa, S., Kitaura, R. & Noro, S. (2004). *Angew. Chem. Int. Ed.* **43**, 2334–2375.
- Lee, C. M. & Kumler, W. D. (1962). *J. Am. Chem. Soc.* **84**, 571–578.
- Moulton, B. & Zaworotko, M. J. (2001). *Chem. Rev.* **101**, 1629–1658.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, Y., Huang, Y. Q., Liu, G. X., Okamura, T.-A., Doi, M., Sheng, Y. W., Sun, W. Y. & Ueyama, N. (2007). *Chem. Eur. J.* **13**, 7523–7531.
- Wu, S. T., Ma, L. Q., Long, L. S., Zheng, L. S. & Lin, W. B. (2009). *Inorg. Chem.* **48**, 2436–2442.
- Yaghi, O. M., O’Keeffe, M., Ockwig, N. W., Chae, H. K., Eddaoudi, M. & Kim, J. (2003). *Nature (London)*, **423**, 705–714.
- Zhang, L. Y., Liu, G. F., Zheng, S. L., Ye, B. H. & Chen, X. M. (2003). *Eur. J. Inorg. Chem.* pp. 2965–2971.

supporting information

Acta Cryst. (2010). E66, m1253–m1254 [doi:10.1107/S1600536810036147]

catena-Poly[[[diaquacadmium(II)]-bis[μ -3,5-bis(isonicotinamido)benzoato]] tetrahydrate]

Man-Sheng Chen, Yi-Fang Deng, Chun-Hua Zhang and Dai-Zhi Kuang

S1. Comment

Coordination polymeric structures are one of the most attractive areas of materials research due to their intriguing structural topologies and functional properties such as molecular adsorption, magnetism and luminescence (Evans & Lin, 2002; Yaghi *et al.*, 2003; Kitagawa *et al.*, 2004; Biradha *et al.*, 2006; Wu *et al.*, 2009) but in spite of this interest the rational design and synthesis of new supramolecular frameworks by covalent and weak intra/intermolecular interactions is still a challenge (Eddaoudi, *et al.* 2001; Moulton & Zaworotko, 2001; Cheng, *et al.* 2002; Zhang, *et al.* 2003; Go, *et al.* 2004). On the other hand, it is well known that carboxylate and pyridine groups have good coordination capacities as well as the amide group, a fascinating functional group with two different types of hydrogen bonding sites: the –NH moiety which acts as an electron acceptor while the –C=O group acts as an electron donor (Lee, *et al.*, 1962; Bent, 1968; Huyskens, 1977; Wang, *et al.*, 2007). we have recently pursued systematic investigations into the assembly of polymers through ligands containing both carboxylate and amido-pyridine groups, in order to study the influence of different metal ions. In the present work we report the structure of a new cadmium coordination polymer of the bridging ligand 3,5-bis(isonicotinamido)benzoate (BBA⁻), namely {[Cd(BBA)₂(H₂O)₂].4(H₂O)}_n, (I), which is isomorphous to its Mn isologue [Mn(BBA)₂(H₂O)₂](Chen, *et al.* 2009).

As shown in Fig. 1, the asymmetric unit of (I) is composed of one-half of a Cd^{II} cation, which sits on a crystallographic twofold axis, one BBA⁻ ligand, one coordinated water molecule and two solvent water molecules. The central Cd^{II} atom has the [CdN₂O₄] octahedral coordination geometry with four coordination sites in one plane occupied by two *cis*-positioned N atoms from two BBA⁻ ligands and two water molecules, and the other two coordination sites taken up by two *trans*-positioned carboxylate O atoms from two further BBA⁻ ligands. The ligand (3,5-bis(isonicotinamido)-benzoic acid) is fully deprotonated (BBA⁻) and the carboxylate group adopts a monodentate mode, coordinating only through one O atom to one Cd^{II} centre. On the other hand, the dihedral angles between the central benzene and terminal pyridine ring are 15.66 (10) ° and 19.72 (11)°, respectively.

It is noteworthy that each BBA⁻ ligand in turn uses its carboxylate group and one of the two pyridinyl groups to connect two metal centers, while the other pyridinyl group does not coordinate. Then, two Cd(II) and two BBA⁻ ligands form a Cd₂(BBA)₂ macrocyclic ring with Cd...Cd distance of 12.313 (18) Å. Such M₂(BBA)₂ macrocyclic rings are further connected by Cd—N and Cd—O coordination bonds to give an infinite one-dimensional zigzag chain structure (Fig. 2).

In addition, there are non-bonding interactions which consolidate the framework structure, in particular some O—H...O, N—H...O and O—H...N hydrogen bonds (Table 1) as well as π - π interactions between the central benzene ring of BBA⁻ anions (C2-->C7) and its symmetry related counterpart, symmetry code: 1/2 - x, 3/2 - y, -z) with a centroid-centroid distance of 3.639 (2) Å, and slippage and interplanar distances of 1.514 and 3.308 Å, respectively.

S2. Experimental

All reagents and solvents were used as obtained commercially without further purification. A mixture containing $\text{Cd}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (31.1 mg, 0.1 mmol), HL (36.5 mg, 0.12 mmol), $\text{N}(\text{CH}_2\text{CH}_3)_3$ (0.5 mL, 0.1 mmol), 10 ml H_2O was sealed in a 16 ml Teflon-lined stainless steel container and heated at 393 K for 3 days. After cooling to room temperature within 12 h, block colorless crystals of (I) suitable for X-ray diffraction analysis were obtained in 39% Yield. Anal. Calcd for $\text{C}_{38}\text{H}_{38}\text{CdN}_8\text{O}_{14}$: C, 48.39; H, 4.06; N, 11.88%; Found: C, 48.43; H, 4.11; N, 11.76%.

S3. Refinement

H atoms bonded to C atoms were placed geometrically and treated as riding, with C—H distances 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The H atoms of water molecules were determined from a difference Fourier synthesis and refined with restrained O—H distances 0.85 (2) Å. The amide H atoms were located from difference maps and refined with the N—H distances restrained to 0.8600 Å.





Figure 1

The ORTEP drawing of the title compound (I). Displacement ellipsoids are drawn at 30% probability level [symmetry code: (i) $-x, y, -z - 1/2$ (ii) $x, 2 - y, -1/2 + z$ (iii) $-x, 2 - y, -z$].

**Figure 2**

Projection showing the one-dimensional structure of the compound (I).

catena-Poly[[[diaquacadmium(II)]-bis[μ -3,5-bis(isonicotinamido)benzoato]] tetrahydrate]

Crystal data

[Cd(C₁₉H₁₃N₄O₄)₂(H₂O)₂] \cdot 4H₂O

$M_r = 943.16$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 17.584$ (3) Å

$b = 10.8568$ (19) Å

$c = 21.891$ (4) Å

$\beta = 103.801$ (2)°

$V = 4058.5$ (12) Å³

$Z = 4$

$F(000) = 1928$

$D_x = 1.544$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3272 reflections

$\theta = 2.3$ – 25.3 °

$\mu = 0.62$ mm⁻¹

$T = 293$ K

Block, colorless

$0.20 \times 0.16 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ϕ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2000)

$T_{\min} = 0.887$, $T_{\max} = 0.941$

10338 measured reflections

3867 independent reflections

3391 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.055$

$\theta_{\max} = 25.8$ °, $\theta_{\min} = 1.9$ °

$h = -21 \rightarrow 21$

$k = -13 \rightarrow 6$

$l = -26 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.110$

$S = 1.08$

3867 reflections

300 parameters

13 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0527P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.022$

$\Delta\rho_{\max} = 0.80$ e Å⁻³

$\Delta\rho_{\min} = -0.48$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.0000	0.74023 (3)	-0.2500	0.03203 (14)
C1	0.1391 (2)	0.6471 (3)	-0.14433 (15)	0.0341 (8)
C2	0.16577 (18)	0.6286 (3)	-0.07425 (14)	0.0304 (7)
C3	0.21138 (19)	0.5274 (3)	-0.04962 (15)	0.0340 (8)
H3A	0.2266	0.4705	-0.0762	0.041*
C4	0.23363 (19)	0.5125 (3)	0.01481 (16)	0.0334 (8)
C5	0.2124 (2)	0.5954 (3)	0.05570 (15)	0.0360 (8)
H5	0.2276	0.5832	0.0990	0.043*
C6	0.16780 (19)	0.6977 (3)	0.03069 (15)	0.0312 (7)
C7	0.14483 (19)	0.7127 (3)	-0.03395 (15)	0.0316 (7)
H7	0.1147	0.7807	-0.0505	0.038*
C8	0.3530 (2)	0.3931 (3)	0.03346 (16)	0.0390 (8)
C9	0.3967 (2)	0.2847 (3)	0.06802 (18)	0.0406 (9)
C10	0.4431 (2)	0.2140 (4)	0.0395 (2)	0.0550 (11)
H10	0.4453	0.2292	-0.0018	0.066*
C11	0.4868 (3)	0.1191 (4)	0.0739 (2)	0.0638 (13)
H11	0.5172	0.0701	0.0541	0.077*
C12	0.4417 (3)	0.1640 (4)	0.1594 (2)	0.0609 (12)
H12	0.4406	0.1472	0.2008	0.073*
C13	0.3957 (3)	0.2595 (3)	0.1292 (2)	0.0532 (11)
H13	0.3648	0.3054	0.1499	0.064*
C14	0.1521 (2)	0.7883 (3)	0.13085 (16)	0.0372 (8)
C15	0.1209 (2)	0.9006 (3)	0.15632 (15)	0.0352 (8)
C16	0.1481 (3)	0.9300 (4)	0.21868 (17)	0.0524 (11)
H16	0.1864	0.8819	0.2446	0.063*
C17	0.1180 (3)	1.0315 (4)	0.24232 (18)	0.0534 (11)
H17	0.1380	1.0517	0.2844	0.064*
C18	0.0367 (2)	1.0727 (4)	0.14846 (17)	0.0487 (10)
H18	-0.0018	1.1220	0.1236	0.058*
C19	0.0636 (2)	0.9741 (3)	0.12058 (17)	0.0460 (9)
H19	0.0433	0.9572	0.0782	0.055*
N1	0.4871 (2)	0.0953 (3)	0.13331 (19)	0.0625 (10)
N2	0.06210 (18)	1.1019 (3)	0.20837 (13)	0.0398 (7)
N3	0.28104 (17)	0.4100 (3)	0.04134 (13)	0.0402 (7)
H3	0.2624	0.3573	0.0632	0.048*

N4	0.14702 (16)	0.7905 (3)	0.06830 (12)	0.0353 (7)
H4	0.1288	0.8572	0.0491	0.042*
O1	0.09323 (16)	0.7351 (2)	-0.16214 (11)	0.0429 (6)
O2	0.16316 (16)	0.5753 (2)	-0.17948 (11)	0.0505 (7)
O3	0.38414 (17)	0.4597 (2)	0.00154 (13)	0.0585 (8)
O4	0.17829 (19)	0.7016 (2)	0.16454 (12)	0.0571 (8)
O5	0.07154 (19)	0.6011 (3)	-0.29733 (14)	0.0501 (7)
O6	0.1941 (3)	0.3124 (5)	0.1269 (2)	0.1053 (14)
O7	0.2760 (3)	0.1472 (3)	0.20727 (17)	0.0782 (11)
H5A	0.050 (2)	0.539 (3)	-0.3152 (17)	0.055 (13)*
H7B	0.299 (3)	0.154 (6)	0.2454 (12)	0.12 (2)*
H7A	0.283 (3)	0.073 (2)	0.198 (2)	0.11 (2)*
H5B	0.109 (2)	0.575 (4)	-0.2698 (18)	0.088 (19)*
H6B	0.150 (2)	0.325 (7)	0.135 (3)	0.16 (3)*
H6A	0.207 (4)	0.243 (3)	0.143 (3)	0.112 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.0418 (2)	0.0269 (2)	0.0257 (2)	0.000	0.00478 (15)	0.000
C1	0.0364 (19)	0.0347 (19)	0.0303 (18)	-0.0015 (16)	0.0064 (15)	-0.0028 (15)
C2	0.0296 (17)	0.0320 (17)	0.0274 (17)	0.0024 (14)	0.0027 (14)	0.0007 (13)
C3	0.038 (2)	0.0279 (18)	0.0353 (19)	0.0029 (15)	0.0069 (16)	-0.0035 (14)
C4	0.0333 (19)	0.0292 (18)	0.0377 (19)	0.0076 (14)	0.0082 (15)	0.0049 (14)
C5	0.043 (2)	0.0360 (19)	0.0276 (18)	0.0080 (16)	0.0061 (15)	0.0047 (14)
C6	0.0327 (18)	0.0293 (17)	0.0324 (18)	0.0043 (14)	0.0095 (15)	0.0024 (14)
C7	0.0304 (18)	0.0292 (17)	0.0348 (18)	0.0057 (14)	0.0069 (15)	0.0022 (14)
C8	0.045 (2)	0.0354 (19)	0.035 (2)	0.0108 (17)	0.0064 (17)	0.0046 (15)
C9	0.040 (2)	0.0330 (19)	0.047 (2)	0.0070 (16)	0.0064 (17)	0.0066 (16)
C10	0.055 (3)	0.049 (2)	0.062 (3)	0.016 (2)	0.015 (2)	0.010 (2)
C11	0.053 (3)	0.045 (2)	0.095 (4)	0.016 (2)	0.020 (3)	0.013 (2)
C12	0.054 (3)	0.063 (3)	0.060 (3)	0.001 (2)	0.003 (2)	0.024 (2)
C13	0.053 (2)	0.052 (3)	0.053 (3)	0.013 (2)	0.010 (2)	0.0152 (19)
C14	0.041 (2)	0.0374 (19)	0.0326 (19)	0.0035 (16)	0.0077 (16)	-0.0008 (15)
C15	0.043 (2)	0.0338 (18)	0.0314 (18)	0.0054 (16)	0.0139 (16)	0.0018 (14)
C16	0.066 (3)	0.049 (2)	0.038 (2)	0.023 (2)	0.0029 (19)	-0.0039 (18)
C17	0.071 (3)	0.054 (3)	0.030 (2)	0.014 (2)	0.0031 (19)	-0.0071 (18)
C18	0.056 (2)	0.053 (2)	0.037 (2)	0.023 (2)	0.0090 (18)	0.0032 (18)
C19	0.054 (2)	0.053 (2)	0.0300 (19)	0.014 (2)	0.0076 (17)	-0.0028 (17)
N1	0.044 (2)	0.050 (2)	0.090 (3)	0.0065 (17)	0.006 (2)	0.027 (2)
N2	0.0495 (19)	0.0383 (17)	0.0319 (16)	0.0079 (14)	0.0104 (14)	0.0013 (13)
N3	0.0494 (19)	0.0319 (16)	0.0403 (18)	0.0120 (14)	0.0127 (15)	0.0111 (12)
N4	0.0453 (18)	0.0305 (15)	0.0307 (16)	0.0105 (13)	0.0104 (14)	0.0038 (12)
O1	0.0536 (16)	0.0396 (14)	0.0300 (13)	0.0159 (12)	-0.0009 (11)	-0.0014 (10)
O2	0.0682 (19)	0.0500 (16)	0.0320 (14)	0.0208 (14)	0.0094 (13)	-0.0039 (12)
O3	0.0618 (18)	0.0565 (18)	0.0640 (18)	0.0204 (15)	0.0286 (15)	0.0286 (15)
O4	0.094 (2)	0.0429 (15)	0.0350 (15)	0.0257 (16)	0.0179 (15)	0.0076 (12)
O5	0.0571 (19)	0.0382 (16)	0.0509 (18)	0.0037 (13)	0.0048 (14)	-0.0158 (13)

O6	0.105 (3)	0.114 (3)	0.112 (3)	0.032 (3)	0.057 (3)	0.061 (3)
O7	0.124 (3)	0.058 (2)	0.053 (2)	0.027 (2)	0.021 (2)	-0.0028 (17)

Geometric parameters (Å, °)

Cd1—O1	2.210 (2)	C11—H11	0.9300
Cd1—O1 ⁱ	2.210 (2)	C12—N1	1.318 (5)
Cd1—N2 ⁱⁱ	2.331 (3)	C12—C13	1.382 (5)
Cd1—N2 ⁱⁱⁱ	2.331 (3)	C12—H12	0.9300
Cd1—O5 ⁱ	2.358 (3)	C13—H13	0.9300
Cd1—O5	2.358 (3)	C14—O4	1.217 (4)
C1—O2	1.239 (4)	C14—N4	1.351 (4)
C1—O1	1.252 (4)	C14—C15	1.498 (5)
C1—C2	1.507 (4)	C15—C16	1.372 (5)
C2—C7	1.379 (4)	C15—C19	1.374 (5)
C2—C3	1.391 (4)	C16—C17	1.376 (5)
C3—C4	1.380 (4)	C16—H16	0.9300
C3—H3A	0.9300	C17—N2	1.325 (5)
C4—C5	1.382 (4)	C17—H17	0.9300
C4—N3	1.428 (4)	C18—N2	1.319 (4)
C5—C6	1.394 (4)	C18—C19	1.371 (5)
C5—H5	0.9300	C18—H18	0.9300
C6—C7	1.386 (4)	C19—H19	0.9300
C6—N4	1.404 (4)	N2—Cd1 ⁱⁱ	2.331 (3)
C7—H7	0.9300	N3—H3	0.8600
C8—O3	1.222 (4)	N4—H4	0.8600
C8—N3	1.330 (4)	O5—H5A	0.829 (19)
C8—C9	1.507 (5)	O5—H5B	0.83 (3)
C9—C13	1.372 (6)	O6—H6B	0.84 (2)
C9—C10	1.373 (6)	O6—H6A	0.83 (2)
C10—C11	1.394 (5)	O7—H7B	0.84 (2)
C10—H10	0.9300	O7—H7A	0.85 (2)
C11—N1	1.325 (5)		
O1—Cd1—O1 ⁱ	177.13 (12)	C11—C10—H10	120.7
O1—Cd1—N2 ⁱⁱ	89.86 (9)	N1—C11—C10	123.1 (4)
O1 ⁱ —Cd1—N2 ⁱⁱ	92.25 (10)	N1—C11—H11	118.5
O1—Cd1—N2 ⁱⁱⁱ	92.25 (10)	C10—C11—H11	118.5
O1 ⁱ —Cd1—N2 ⁱⁱⁱ	89.86 (9)	N1—C12—C13	124.4 (4)
N2 ⁱⁱ —Cd1—N2 ⁱⁱⁱ	85.36 (15)	N1—C12—H12	117.8
O1—Cd1—O5 ⁱ	87.95 (10)	C13—C12—H12	117.8
O1 ⁱ —Cd1—O5 ⁱ	90.21 (10)	C9—C13—C12	118.2 (4)
N2 ⁱⁱ —Cd1—O5 ⁱ	87.17 (11)	C9—C13—H13	120.9
N2 ⁱⁱⁱ —Cd1—O5 ⁱ	172.53 (10)	C12—C13—H13	120.9
O1—Cd1—O5	90.21 (10)	O4—C14—N4	123.5 (3)
O1 ⁱ —Cd1—O5	87.95 (10)	O4—C14—C15	121.6 (3)
N2 ⁱⁱ —Cd1—O5	172.53 (10)	N4—C14—C15	114.9 (3)
N2 ⁱⁱⁱ —Cd1—O5	87.17 (11)	C16—C15—C19	117.8 (3)

O5 ⁱ —Cd1—O5	100.30 (15)	C16—C15—C14	119.2 (3)
O2—C1—O1	125.3 (3)	C19—C15—C14	123.0 (3)
O2—C1—C2	118.7 (3)	C15—C16—C17	119.1 (4)
O1—C1—C2	116.1 (3)	C15—C16—H16	120.5
C7—C2—C3	119.5 (3)	C17—C16—H16	120.5
C7—C2—C1	119.8 (3)	N2—C17—C16	123.5 (4)
C3—C2—C1	120.7 (3)	N2—C17—H17	118.2
C4—C3—C2	119.0 (3)	C16—C17—H17	118.2
C4—C3—H3A	120.5	N2—C18—C19	124.0 (3)
C2—C3—H3A	120.5	N2—C18—H18	118.0
C3—C4—C5	122.1 (3)	C19—C18—H18	118.0
C3—C4—N3	120.2 (3)	C15—C19—C18	118.9 (3)
C5—C4—N3	117.7 (3)	C15—C19—H19	120.5
C4—C5—C6	118.5 (3)	C18—C19—H19	120.5
C4—C5—H5	120.7	C12—N1—C11	117.0 (4)
C6—C5—H5	120.7	C18—N2—C17	116.7 (3)
C7—C6—C5	119.6 (3)	C18—N2—Cd1 ⁱⁱ	119.1 (2)
C7—C6—N4	117.5 (3)	C17—N2—Cd1 ⁱⁱ	124.0 (2)
C5—C6—N4	122.9 (3)	C8—N3—C4	122.4 (3)
C2—C7—C6	121.2 (3)	C8—N3—H3	118.8
C2—C7—H7	119.4	C4—N3—H3	118.8
C6—C7—H7	119.4	C14—N4—C6	128.2 (3)
O3—C8—N3	124.2 (3)	C14—N4—H4	115.9
O3—C8—C9	120.3 (3)	C6—N4—H4	115.9
N3—C8—C9	115.5 (3)	C1—O1—Cd1	125.4 (2)
C13—C9—C10	118.7 (3)	Cd1—O5—H5A	120 (3)
C13—C9—C8	121.5 (3)	Cd1—O5—H5B	109 (3)
C10—C9—C8	119.7 (4)	H5A—O5—H5B	105 (4)
C9—C10—C11	118.6 (4)	H6B—O6—H6A	104 (6)
C9—C10—H10	120.7	H7B—O7—H7A	104 (5)
O2—C1—C2—C7	175.0 (3)	O4—C14—C15—C19	-151.8 (4)
O1—C1—C2—C7	-5.5 (5)	N4—C14—C15—C19	26.8 (5)
O2—C1—C2—C3	-5.1 (5)	C19—C15—C16—C17	-0.9 (6)
O1—C1—C2—C3	174.5 (3)	C14—C15—C16—C17	-178.5 (4)
C7—C2—C3—C4	1.0 (5)	C15—C16—C17—N2	1.8 (7)
C1—C2—C3—C4	-179.0 (3)	C16—C15—C19—C18	0.3 (6)
C2—C3—C4—C5	-0.4 (5)	C14—C15—C19—C18	177.8 (4)
C2—C3—C4—N3	-179.4 (3)	N2—C18—C19—C15	-0.5 (6)
C3—C4—C5—C6	-0.7 (5)	C13—C12—N1—C11	-1.2 (7)
N3—C4—C5—C6	178.4 (3)	C10—C11—N1—C12	1.9 (7)
C4—C5—C6—C7	1.1 (5)	C19—C18—N2—C17	1.3 (6)
C4—C5—C6—N4	-176.4 (3)	C19—C18—N2—Cd1 ⁱⁱ	-173.3 (3)
C3—C2—C7—C6	-0.6 (5)	C16—C17—N2—C18	-1.9 (6)
C1—C2—C7—C6	179.4 (3)	C16—C17—N2—Cd1 ⁱⁱ	172.4 (3)
C5—C6—C7—C2	-0.5 (5)	O3—C8—N3—C4	-2.7 (6)
N4—C6—C7—C2	177.1 (3)	C9—C8—N3—C4	176.3 (3)
O3—C8—C9—C13	135.5 (4)	C3—C4—N3—C8	62.3 (5)

N3—C8—C9—C13	-43.5 (5)	C5—C4—N3—C8	-116.8 (4)
O3—C8—C9—C10	-40.0 (5)	O4—C14—N4—C6	1.7 (6)
N3—C8—C9—C10	141.0 (4)	C15—C14—N4—C6	-176.9 (3)
C13—C9—C10—C11	0.4 (6)	C7—C6—N4—C14	169.1 (3)
C8—C9—C10—C11	176.0 (4)	C5—C6—N4—C14	-13.4 (6)
C9—C10—C11—N1	-1.5 (7)	O2—C1—O1—Cd1	31.5 (5)
C10—C9—C13—C12	0.2 (6)	C2—C1—O1—Cd1	-148.0 (2)
C8—C9—C13—C12	-175.3 (4)	N2 ⁱⁱ —Cd1—O1—C1	160.5 (3)
N1—C12—C13—C9	0.2 (7)	N2 ⁱⁱⁱ —Cd1—O1—C1	-114.2 (3)
O4—C14—C15—C16	25.7 (6)	O5 ⁱ —Cd1—O1—C1	73.3 (3)
N4—C14—C15—C16	-155.7 (3)	O5—Cd1—O1—C1	-27.0 (3)

Symmetry codes: (i) $-x, y, -z-1/2$; (ii) $-x, -y+2, -z$; (iii) $x, -y+2, z-1/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N3—H3...O6	0.86	2.10	2.885 (5)	151
N4—H4...O3 ^{iv}	0.86	2.26	3.096 (4)	164
O5—H5B...O2	0.83 (3)	1.98 (3)	2.710 (4)	145 (4)
O5—H5A...N1 ^v	0.83 (2)	2.00 (2)	2.825 (4)	172 (4)
O6—H6B...O5 ^{vi}	0.84 (2)	2.40 (4)	3.163 (6)	151 (6)
O6—H6A...O7	0.82 (3)	1.92 (4)	2.681 (6)	148 (5)
O7—H7A...O2 ^{vii}	0.85 (2)	1.96 (3)	2.767 (4)	162 (5)
O7—H7B...O4 ^{viii}	0.84 (2)	1.99 (3)	2.791 (4)	159 (6)

Symmetry codes: (iv) $-x+1/2, -y+3/2, -z$; (v) $x-1/2, -y+1/2, z-1/2$; (vi) $x, -y+1, z+1/2$; (vii) $-x+1/2, -y+1/2, -z$; (viii) $-x+1/2, y-1/2, -z+1/2$.