

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(1*E*,2*E*)-1,2-Bis(2,3,4-trimethoxybenzylidene)hydrazine

 Malai Haniti S. A. Hamid,^a Mohammad Akbar Ali,^a
 Aminul Huq Mirza,^a Gan Ai Len^a and Ray J. Butcher^{b*}
^aFaculty of Science, Universiti Brunei Darussalam, Jln Tungku Link, BE 1410, Negara Brunei Darussalam, and ^bDepartment of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA

Correspondence e-mail: rbutcher99@yahoo.com


Received 25 August 2010; accepted 31 August 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.048; wR factor = 0.123; data-to-parameter ratio = 16.9.

The title compound, $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$, was obtained as an unexpected product by the reaction of hydrazinium dithiocarbazate with 2,3,4-trimethoxybenzaldehyde in refluxing ethanol. The molecule lies on a center of inversion. The crystal packing is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions.

Related literature

The surprising formation of the title hydrazone was probably due to the decomposition of hydrazinium dithiocarbazate in solution resulting in the formation of hydrazine, which then reacted with 2,3,4-trimethoxybenzaldehyde. Hydrazinium dithiocarbazates are known to decompose on heating (Rudorf, 2007). For the biological activity of Schiff bases, see: Akbar Ali *et al.* (2008); Chan *et al.* (2008). For a previous report of the title compound (the X-ray structure was not provided), see: Praefcke *et al.* (1991). For comparison bond lengths in an aroyl hydrazone, see: Ji *et al.* (2010).



Experimental

Crystal data

 $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$
 $M_r = 388.41$

 Monoclinic, $P2_1/n$
 $a = 10.0380$ (9) Å
 $b = 7.0713$ (7) Å
 $c = 13.9586$ (14) Å
 $\beta = 102.800$ (2)°
 $V = 966.18$ (16) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.60 \times 0.36 \times 0.04$ mm

Data collection

 Bruker SMART CCD area-detector
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.943$, $T_{\max} = 0.996$

 6576 measured reflections
 2203 independent reflections
 1846 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.123$
 $S = 1.05$
 2196 reflections

 130 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.20$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C7}-\text{H7B}\cdots\text{O3}^{\text{ii}}$	0.98	2.54	3.3620 (18)	142
$\text{C8}-\text{H8B}\cdots\text{O1}^{\text{iii}}$	0.98	2.62	3.3735 (19)	134
$\text{C8}-\text{H8B}\cdots\text{O2}^{\text{iii}}$	0.98	2.62	3.5711 (18)	164

 Symmetry codes: (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT-Plus (Bruker, 1998); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

MHSAH, MAA, AHM and GAL thank Universiti Brunei Darussalam for support. The X-ray Diffraction Laboratory, Department of Chemistry, National University of Singapore, is acknowledged for the collection of the X-ray diffraction data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5339).

References

- Akbar Ali, M., Abu Bakar, J. H., Mirza, A. H., Smith, S. J., Gahan, L. R. & Bernhardt, P. V. (2008). *Polyhedron*, **27**, 71–79.
 Bruker (1998). *SAINTE-Plus* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chan, M.-H. E., Crouse, K. A., Tahir, M. I. M., Rosli, R., Umar-Tsafe, N. & Cowley, A. R. (2008). *Polyhedron*, **27**, 1141–1149.
 Ji, N.-N., Shi, A.-Q., Zhao, R.-G., Zheng, Z.-B. & Li, Z.-F. (2010). *Bull. Korean Chem. Soc.* **31**, 881–886.
 Praefcke, K., Kohne, B., Guendogan, B., Singer, D. & Demus, D. (1991). *Mol. Cryst. Liq. Cryst.* **198**, 393–405.
 Rudorf, W.-D. (2007). *J. Sulfur Chem.* **28**, 295–339.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2010). E66, o2557 [doi:10.1107/S160053681003518X]

(1*E*,2*E*)-1,2-Bis(2,3,4-trimethoxybenzylidene)hydrazine

Malai Haniti S. A. Hamid, Mohammad Akbar Ali, Aminul Huq Mirza, Gan Ai Len and Ray J. Butcher

S1. Comment

The compound, C₂₀H₂₄N₂O₆ (I) was obtained by the reaction of hydrazinium dithiocarbazate and 2,3,4-trimethoxybenzaldehyde in boiling ethanol. The surprising formation of the hydrazone was probably due to the decomposition of hydrazinium dithiocarbazate in solution resulting in the formation of hydrazine, which then reacted with 2,3,4-trimethoxybenzaldehyde to form the corresponding hydrazone (I). Hydrazinium dithiocarbazates are known to decompose on heating (Rudorf, 2007).

Schiff bases have attracted considerable attention because they can act as chelating agents for metal ions and many of them also exhibit useful biological activities (Akbar Ali *et al.*, 2008; Chan *et al.*, 2008). Although the compound has previously been reported its X-ray structure has not been provided (Praefcke *et al.*, 1991). Hydrazones derived from the reactions of hydrazines with aldehydes or ketones are common but bis-hydrazones are not.

The molecular structure of (I) is shown in Figure 1 and its selected bond lengths and angles are given in Table 1. Like most thiosemicarbazones and Schiff bases, the imine moiety in [I] shows an E configuration about the C10—N1 [1.283 (2) Å] and N1A—C10A bonds. The C10—N1 and N1A—C10A bond distances also compare well with C=N double bonds in other related compounds. A comparison of the N(1)—N(1 A) distance [1.413 (3) Å] with that in an aroyl hydrazone [1.377 (3) Å] (Ji *et al.* 2010) shows that the bond is shorter than a single N—N bond (1.44 Å) indicating that a significant π -charge delocalization occurs along the C—N—N—C moiety. As the bond angles C6—C10—N1 (121.68°) and C6A—C10A—N1A (121.68°) are close to that of a sp^2 -hybridized carbon atom (*ca* 120°), the molecule does not have a distorted geometry. Due the fact that the molecule lies on a center of inversion the dihedral angle between the two phenyl rings is 0.0°.

Figure 2 shows the packing of (I) in the unit cell. The packing diagram shows that there are intermolecular hydrogen bonds between one of the CH₃ hydrogen atoms of one molecule with an ether oxygen of another molecule.

S2. Experimental

2,3,4-trimethoxybenzaldehyde (0.24 g, 1.24 mmol) dissolved in absolute ethanol (5 ml) was mixed with a solution of hydrazinium dithiocarbazate (0.93 g, 0.66 mmol) in the same solvent (45 ml). After refluxing for two hours, the resulting clear yellow solution was left to stand at room temperature for five days to afford crystalline yellow plates. The crystals were filtered, washed with cold absolute ethanol and dried *in vacuo*. Yield: 0.152 g (63%); m.p. 192–194 °C; IR (KBr, cm⁻¹): 2968, 2937, 2832, 1614, 1590, 1494, 1457, 1431, 1410, 1286, 1229, 1199, 1166, 1090, 1023, 1008, 943, 898, 809, 699, 667, 594, 540, 433; ¹H NMR (400 MHz, CDCl₃, 30 °C): δ 8.92 (2*H*, s, CH=N), 7.84 (2*H*, d, ArH), 6.76 (2*H*, d, ArH), 3.96 (6*H*, s, OCH₃), 3.92 (6*H*, s, OCH₃), 3.90 (6*H*, s, OCH₃); Anal. Calcd. for C₂₀H₂₄N₂O₆ (388.42): C 61.85, H 6.23, N 7.21. Found: C 62.17, H 6.17, N 7.47%.

IR spectrum was recorded as a KBr disc with 13 mm KBr discs SPECAC accessory on a Perkin-Elmer 1600 FT IR spectrometer. ^1H NMR spectrum was run in CDCl_3 on a Varian 400-NMR spectrometer at Universiti Brunei Darussalam. Elemental analysis for C, H and N was done by the Elemental Analysis Laboratory, Department of Chemistry, National University of Singapore. The X-ray data were collected at the X-ray Diffraction Laboratory, Department of Chemistry, National University of Singapore using a Bruker-AXS Smart Apex CCD single-crystal diffractometer.

S3. Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with a C—H distances of 0.95 Å and 0.98 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

The title compound, $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$ with atom labeling. Displacement ellipsoids are at the 50 % probability level.




Figure 2

The molecular packing for $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_6$ viewed down the a axis showing the intermolecular C—H...O interactions.

(1E,2E)-1,2-Bis(2,3,4-trimethoxybenzylidene)hydrazine*Crystal data*C₂₀H₂₄N₂O₆ $M_r = 388.41$ Monoclinic, $P2_1/n$ $a = 10.0380$ (9) Å $b = 7.0713$ (7) Å $c = 13.9586$ (14) Å $\beta = 102.800$ (2)° $V = 966.18$ (16) Å³ $Z = 2$ $F(000) = 412$ $D_x = 1.335$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1770 reflections

 $\theta = 2.8$ – 27.1 ° $\mu = 0.10$ mm⁻¹ $T = 100$ K

Plate, yellow

 $0.60 \times 0.36 \times 0.04$ mm*Data collection*Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scansAbsorption correction: multi-scan
(*SADABS*; Sheldrick, 1996) $T_{\min} = 0.943$, $T_{\max} = 0.996$

6576 measured reflections

2203 independent reflections

1846 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.033$ $\theta_{\max} = 27.5$ °, $\theta_{\min} = 2.3$ ° $h = -13 \rightarrow 12$ $k = 0 \rightarrow 9$ $l = 0 \rightarrow 18$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.048$ $wR(F^2) = 0.123$ $S = 1.05$

2196 reflections

130 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0608P)^2 + 0.353P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.35$ e Å⁻³ $\Delta\rho_{\min} = -0.20$ e Å⁻³*Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.62846 (10)	0.45681 (14)	0.85047 (7)	0.0180 (2)
O2	0.61624 (9)	0.20990 (14)	0.70188 (7)	0.0165 (2)
O3	0.38623 (10)	0.18335 (14)	0.55439 (7)	0.0175 (2)
C10	0.15962 (14)	0.4221 (2)	0.54359 (10)	0.0154 (3)
H10	0.1582	0.3285	0.4945	0.018*

C1	0.29226 (14)	0.5791 (2)	0.69604 (11)	0.0172 (3)
H1	0.2190	0.6649	0.6944	0.021*
C2	0.40646 (15)	0.5899 (2)	0.77231 (10)	0.0180 (3)
H2	0.4113	0.6834	0.8218	0.022*
C3	0.51481 (14)	0.4635 (2)	0.77677 (10)	0.0153 (3)
C4	0.50841 (13)	0.33003 (19)	0.70148 (10)	0.0146 (3)
C5	0.39310 (14)	0.32122 (19)	0.62483 (10)	0.0147 (3)
C6	0.28211 (14)	0.44466 (19)	0.62113 (10)	0.0149 (3)
C7	0.64403 (15)	0.6012 (2)	0.92428 (11)	0.0202 (3)
H7A	0.6488	0.7252	0.8938	0.030*
H7B	0.7282	0.5789	0.9740	0.030*
H7C	0.5657	0.5983	0.9555	0.030*
C8	0.60816 (16)	0.0407 (2)	0.75753 (11)	0.0218 (3)
H8A	0.6081	0.0745	0.8256	0.033*
H8B	0.6871	-0.0401	0.7563	0.033*
H8C	0.5238	-0.0274	0.7286	0.033*
C9	0.44808 (17)	0.2382 (2)	0.47521 (11)	0.0250 (4)
H9A	0.3970	0.3440	0.4393	0.038*
H9B	0.4466	0.1309	0.4305	0.038*
H9C	0.5428	0.2769	0.5017	0.038*
N1	0.05345 (12)	0.52567 (17)	0.53984 (8)	0.0165 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0149 (5)	0.0161 (5)	0.0206 (5)	0.0020 (4)	-0.0011 (4)	-0.0043 (4)
O2	0.0135 (5)	0.0140 (5)	0.0223 (5)	0.0028 (4)	0.0046 (4)	0.0010 (4)
O3	0.0191 (5)	0.0148 (5)	0.0188 (5)	-0.0001 (4)	0.0043 (4)	-0.0027 (4)
C10	0.0161 (6)	0.0143 (7)	0.0160 (6)	0.0000 (5)	0.0043 (5)	0.0016 (5)
C1	0.0152 (7)	0.0160 (7)	0.0206 (7)	0.0040 (5)	0.0045 (5)	0.0010 (5)
C2	0.0199 (7)	0.0159 (7)	0.0183 (7)	0.0008 (6)	0.0043 (6)	-0.0041 (5)
C3	0.0119 (6)	0.0155 (7)	0.0175 (7)	-0.0021 (5)	0.0014 (5)	0.0012 (5)
C4	0.0130 (6)	0.0130 (7)	0.0189 (7)	0.0015 (5)	0.0055 (5)	0.0020 (5)
C5	0.0167 (6)	0.0128 (7)	0.0155 (6)	-0.0011 (5)	0.0055 (5)	0.0002 (5)
C6	0.0136 (6)	0.0146 (7)	0.0164 (6)	-0.0006 (5)	0.0028 (5)	0.0025 (5)
C7	0.0203 (7)	0.0190 (7)	0.0196 (7)	-0.0001 (6)	0.0009 (6)	-0.0035 (6)
C8	0.0277 (8)	0.0176 (8)	0.0212 (7)	0.0069 (6)	0.0082 (6)	0.0040 (6)
C9	0.0320 (8)	0.0244 (8)	0.0206 (7)	0.0016 (7)	0.0098 (6)	-0.0024 (6)
N1	0.0146 (6)	0.0185 (6)	0.0154 (6)	-0.0004 (5)	0.0012 (5)	0.0020 (5)

Geometric parameters (\AA , $^\circ$)

O1—C3	1.3572 (16)	C3—C4	1.4036 (19)
O1—C7	1.4344 (17)	C4—C5	1.3926 (19)
O2—C4	1.3750 (16)	C5—C6	1.4072 (19)
O2—C8	1.4383 (17)	C7—H7A	0.9800
O3—C5	1.3755 (17)	C7—H7B	0.9800
O3—C9	1.4357 (18)	C7—H7C	0.9800

C10—N1	1.2846 (19)	C8—H8A	0.9800
C10—C6	1.4556 (19)	C8—H8B	0.9800
C10—H10	0.9500	C8—H8C	0.9800
C1—C2	1.383 (2)	C9—H9A	0.9800
C1—C6	1.400 (2)	C9—H9B	0.9800
C1—H1	0.9500	C9—H9C	0.9800
C2—C3	1.398 (2)	N1—N1 ⁱ	1.411 (2)
C2—H2	0.9500		
C3—O1—C7	117.32 (11)	C1—C6—C10	122.64 (13)
C4—O2—C8	112.18 (11)	C5—C6—C10	119.40 (13)
C5—O3—C9	113.41 (11)	O1—C7—H7A	109.5
N1—C10—C6	121.62 (13)	O1—C7—H7B	109.5
N1—C10—H10	119.2	H7A—C7—H7B	109.5
C6—C10—H10	119.2	O1—C7—H7C	109.5
C2—C1—C6	121.54 (13)	H7A—C7—H7C	109.5
C2—C1—H1	119.2	H7B—C7—H7C	109.5
C6—C1—H1	119.2	O2—C8—H8A	109.5
C1—C2—C3	120.20 (13)	O2—C8—H8B	109.5
C1—C2—H2	119.9	H8A—C8—H8B	109.5
C3—C2—H2	119.9	O2—C8—H8C	109.5
O1—C3—C2	124.83 (13)	H8A—C8—H8C	109.5
O1—C3—C4	115.81 (12)	H8B—C8—H8C	109.5
C2—C3—C4	119.36 (12)	O3—C9—H9A	109.5
O2—C4—C5	119.67 (12)	O3—C9—H9B	109.5
O2—C4—C3	120.47 (12)	H9A—C9—H9B	109.5
C5—C4—C3	119.85 (12)	O3—C9—H9C	109.5
O3—C5—C4	118.84 (12)	H9A—C9—H9C	109.5
O3—C5—C6	119.97 (12)	H9B—C9—H9C	109.5
C4—C5—C6	121.13 (13)	C10—N1—N1 ⁱ	111.38 (15)
C1—C6—C5	117.88 (13)		
C6—C1—C2—C3	-0.8 (2)	O2—C4—C5—O3	4.25 (19)
C7—O1—C3—C2	-6.1 (2)	C3—C4—C5—O3	-176.89 (12)
C7—O1—C3—C4	174.78 (12)	O2—C4—C5—C6	-178.71 (12)
C1—C2—C3—O1	-177.00 (13)	C3—C4—C5—C6	0.2 (2)
C1—C2—C3—C4	2.1 (2)	C2—C1—C6—C5	-0.9 (2)
C8—O2—C4—C5	-93.85 (15)	C2—C1—C6—C10	175.88 (13)
C8—O2—C4—C3	87.29 (15)	O3—C5—C6—C1	178.16 (12)
O1—C3—C4—O2	-3.77 (19)	C4—C5—C6—C1	1.1 (2)
C2—C3—C4—O2	177.08 (13)	O3—C5—C6—C10	1.3 (2)
O1—C3—C4—C5	177.38 (12)	C4—C5—C6—C10	-175.69 (12)
C2—C3—C4—C5	-1.8 (2)	N1—C10—C6—C1	0.1 (2)
C9—O3—C5—C4	-87.17 (15)	N1—C10—C6—C5	176.74 (13)
C9—O3—C5—C6	95.76 (15)	C6—C10—N1—N1 ⁱ	-178.28 (13)

Symmetry code: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C7—H7B \cdots O3 ⁱⁱ	0.98	2.54	3.3620 (18)	142
C8—H8B \cdots O1 ⁱⁱⁱ	0.98	2.62	3.3735 (19)	134
C8—H8B \cdots O2 ⁱⁱⁱ	0.98	2.62	3.5711 (18)	164

Symmetry codes: (ii) $x+1/2, -y+1/2, z+1/2$; (iii) $-x+3/2, y-1/2, -z+3/2$.