

catena-Poly[(μ_2 -3-carboxy-5-nitrobenzoato)(μ_3 -5-nitrobenzene-1,3-dicarboxylato)(1,10-phenanthroline)-gadolinium(III)]

Jun Wang^{a*} and Zhi-Li Fang^b

^aZhongshan Polytechnic, Zhongshan, Guangdong 528404, People's Republic of China, and ^bSchool of Basic Science, East China Jiaotong University, Nanchang 330013, People's Republic of China

Correspondence e-mail: wangjun7203@126.com


Received 19 August 2010; accepted 2 September 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.012$ Å; R factor = 0.045; wR factor = 0.121; data-to-parameter ratio = 11.9.

The crystal structure of the title complex, $[\text{Gd}(\text{C}_8\text{H}_3\text{NO}_6)(\text{C}_8\text{H}_4\text{NO}_6)(\text{C}_{12}\text{H}_8\text{N}_2)]_n$, contains polymeric chains made up of Gd^{III} atoms, 1,10-phenanthroline and fully or half-deprotonated 5-nitrobenzene-1,3-dicarboxylic acid (H_2L) ligands. The Gd^{III} atom is coordinated in a distorted bicapped trigonal-prismatic fashion by six O atoms from two HL^- and three L^{2-} ligands, and by two N atoms from the 1,10-phenanthroline ligand. The L^{2-} ligands bridge the Gd-phenanthroline units, forming chains running parallel to $[100]$. $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonding as well as $\pi-\pi$ stacking interactions with an interplanar distance of 3.599 (2) Å assemble neighboring polymeric chains.

Related literature

For background to $\pi-\pi$ stacking in biological systems, see: Deisenhofer & Michel (1989). For some crystal structures of metal complexes exhibiting $\pi-\pi$ stacking, see: Li *et al.* (2005); Pan & Xu (2004); Wu *et al.* (2003); Qiu *et al.* (2009).



Experimental

Crystal data

$[\text{Gd}(\text{C}_8\text{H}_3\text{NO}_6)(\text{C}_8\text{H}_4\text{NO}_6)(\text{C}_{12}\text{H}_8\text{N}_2)]_n$
 $M_r = 756.69$
 Triclinic, $P\bar{1}$
 $a = 10.300$ (3) Å
 $b = 12.030$ (3) Å
 $c = 12.150$ (3) Å
 $\alpha = 70.581$ (4)°

$\beta = 85.925$ (4)°
 $\gamma = 76.512$ (2)°
 $V = 1380.6$ (7) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 2.48$ mm⁻¹
 $T = 298$ K
 $0.28 \times 0.26 \times 0.22$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.544$, $T_{\text{max}} = 0.612$

6876 measured reflections
 4837 independent reflections
 4259 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.121$
 $S = 1.04$
 4837 reflections

407 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 2.51$ e Å⁻³
 $\Delta\rho_{\text{min}} = -2.59$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O5}-\text{H5A}\cdots\text{O7}^i$	0.82	2.03	2.737 (7)	145

Symmetry code: (i) $x, y, z + 1$.

Data collection: SMART (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported financially by Zhongshan Polytechnic.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2394).

References

- Bruker (2004). *SMART* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Deisenhofer, J. & Michel, H. (1989). *EMBO J.* **8**, 2149–2170.
- Li, H., Yin, K.-L. & Xu, D.-J. (2005). *Acta Cryst.* **C61**, m19–m21.
- Pan, T.-T. & Xu, D.-J. (2004). *Acta Cryst.* **E60**, m56–m58.
- Qiu, Y.-C., Deng, H., Yang, S.-H., Mou, J.-X., Daiguebonne, C., Kerbellec, N., Guillou, O. & Batten, S. R. (2009). *Inorg. Chem.* **48**, 3976–3981.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, Z.-Y., Xue, Y.-H. & Xu, D.-J. (2003). *Acta Cryst.* **E59**, m809–m811.

supporting information

Acta Cryst. (2010). E66, m1230–m1231 [doi:10.1107/S1600536810035452]

catena-Poly[(μ_2 -3-carboxy-5-nitrobenzoato)(μ_3 -5-nitrobenzene-1,3-dicarboxylato)(1,10-phenanthroline)gadolinium(III)]**Jun Wang and Zhi-Li Fang****S1. Comment**

Recently, we became interested in the nature of π - π stacking as it plays an important role in some biological processes (Deisenhofer & Michel, 1989). A series of metal complexes incorporating different aromatic ligands has been prepared and their crystal structures provide useful information about π - π stacking (Wu *et al.*, 2003; Pan & Xu, 2004; Li *et al.*, 2005; Qiu *et al.*, 2009). As part of our ongoing investigations, the title complex, (I), incorporating 1,10-phenanthroline, has been prepared.


As depicted in Fig. 1, the Gd^{III} atom has a distorted bicapped trigonal-prismatic coordination, defined by six O atoms from two 1-carboxy-5-nitro-3-benzoate (HL^- , where L is 5-nitro-1,3-benzenedicarboxylic acid) and three 5-nitro-1,3-benzenedicarboxylate ligands (L^{2-}), and two N atoms from the 1,10-phenanthroline ligand. The L^{2-} ligands link Gd-phenanthroline moieties, forming an infinite polymeric chain running along [100]. The HL^- and L^{2-} ligands in this complex have two different coordination modes: the first ligand uses one of its carboxylate groups to link one Gd^{III} ions in a chelating coordination mode, the other carboxylate group links two Gd^{III} ions in a bis-monodentate coordination mode, whereas the other ligand uses its carboxylate groups to link two Gd^{III} ions in a bridging coordination mode. In the chain, the closest Gd \cdots Gd separation is 4.333 (3) Å. These chains interact with each other by π - π stacking of adjacent phenanthroline groups (interplanar distance of 3.599 (2) Å). O—H \cdots O hydrogen bonding interactions between carboxyl and carboxylate groups of neighbouring ligands help to consolidate the structure (Table 1; Fig. 2).

S2. Experimental


A sample of Gd₂O₃ (0.0732 g, 0.20 mmol), 5-nitro-1,3-benzenedicarboxylic acid (0.1015 g, 0.50 mmol), 1,10-phenanthroline (0.0991 g, 0.50 mmol) and distilled water (8 ml) were mixed in a Teflon-lined stainless steel vessel with 15 ml capacity. The mixture was heated under autogenous pressure at 393 K for 48 h and cooled slowly to room temperature.

S3. Refinement

The C— and O—bound H atoms were included in the riding-model approximation, with C—H = 0.97 Å and O—H = 0.82 Å, and $U_{iso}(H) = 1.2U_{eq}(C)$ and $1.5U_{eq}(O)$. The highest and the deepest hole the final difference Fourier map are located 1.03 and 1.05 Å, respectively, from the Gd1 atom.

**Figure 1**

Part of the structure of (I), showing the coordination of the Gd atom with the atomic numbering scheme. Non-H atoms are shown with 30% probability displacement ellipsoids, H atoms as spheres of arbitrary radius. [Symmetry codes: (i) $2 - x, -y, 1 - z$; (ii) $1 - x, y, 1 - z$; (iii) $1 + x, y, z$.]

**Figure 2**

View of π - π interactions consolidating the three-dimensional network of (I).

catena-Poly[(μ_2 -3-carboxy-5-nitrobenzoato)(μ_3 -5-nitrobenzene-1,3-dicarboxylato)(1,10-phenanthroline)gadolinium(III)]

Crystal data

[Gd(C₈H₃NO₆)(C₈H₄NO₆)(C₁₂H₈N₂)]

$M_r = 756.69$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.300$ (3) Å

$b = 12.030$ (3) Å

$c = 12.150$ (3) Å

$\alpha = 70.581$ (4)°

$\beta = 85.925$ (4)°

$\gamma = 76.512$ (2)°

$V = 1380.6$ (7) Å³

$Z = 2$

$F(000) = 742$

$D_x = 1.820$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5111 reflections

$\theta = 2.6$ – 27.7 °

$\mu = 2.48$ mm⁻¹

$T = 298$ K

Prism, colorless

$0.28 \times 0.26 \times 0.22$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scan

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.544$, $T_{\max} = 0.612$

6896 measured reflections

4860 independent reflections

4259 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 25.3$ °, $\theta_{\min} = 1.8$ °

$h = -12 \rightarrow 12$

$k = -13 \rightarrow 14$

$l = -14 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.121$

$S = 1.04$

4837 reflections

407 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0811P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 2.51$ e Å⁻³

$\Delta\rho_{\min} = -2.59$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0366 (7)	0.0929 (6)	0.6625 (5)	0.0330 (14)
C2	1.0511 (6)	0.1117 (6)	0.7766 (5)	0.0308 (13)

C3	1.1673 (7)	0.0513 (6)	0.8416 (6)	0.0407 (16)
H3	1.2371	0.0040	0.8132	0.049*
C4	1.1762 (8)	0.0635 (7)	0.9504 (6)	0.0465 (18)
C5	1.0753 (7)	0.1320 (6)	0.9960 (6)	0.0423 (16)
H5	1.0841	0.1378	1.0694	0.051*
C6	0.9615 (7)	0.1914 (7)	0.9309 (6)	0.0419 (16)
C7	0.9488 (7)	0.1835 (6)	0.8200 (6)	0.0374 (15)
H7	0.8723	0.2261	0.7755	0.045*
C8	0.8451 (9)	0.2616 (8)	0.9788 (7)	0.057 (2)
C9	0.5970 (6)	0.1875 (6)	0.3097 (5)	0.0292 (13)
C10	0.4627 (6)	0.2185 (5)	0.2498 (5)	0.0270 (12)
C11	0.4353 (6)	0.3072 (6)	0.1421 (5)	0.0321 (13)
H11	0.5013	0.3440	0.0995	0.039*
C12	0.3063 (6)	0.3388 (6)	0.1007 (5)	0.0328 (14)
C13	0.2038 (6)	0.2870 (6)	0.1622 (5)	0.0314 (13)
H13	0.1169	0.3128	0.1331	0.038*
C14	0.2355 (6)	0.1959 (5)	0.2679 (5)	0.0256 (12)
C15	0.3643 (6)	0.1606 (5)	0.3105 (5)	0.0279 (12)
H15	0.3858	0.0976	0.3804	0.033*
C17	0.5715 (9)	0.1824 (8)	0.6347 (8)	0.061 (2)
H17	0.5663	0.1088	0.6280	0.073*
C18	0.4954 (10)	0.2208 (11)	0.7209 (9)	0.085 (4)
H18	0.4424	0.1727	0.7704	0.103*
C19	0.4993 (10)	0.3254 (11)	0.7313 (9)	0.081 (3)
H19	0.4484	0.3519	0.7878	0.097*
C20	0.5823 (9)	0.3984 (9)	0.6552 (8)	0.064 (3)
C21	0.5923 (12)	0.5114 (12)	0.6605 (11)	0.091 (4)
H21	0.5419	0.5420	0.7148	0.109*
C22	0.6719 (14)	0.5746 (11)	0.5897 (12)	0.096 (4)
H22	0.6781	0.6476	0.5973	0.115*
C23	0.7507 (11)	0.5337 (9)	0.4998 (10)	0.074 (3)
C24	0.8317 (13)	0.5984 (10)	0.4198 (12)	0.095 (4)
H24	0.8409	0.6722	0.4234	0.114*
C25	0.8966 (12)	0.5558 (9)	0.3378 (11)	0.088 (3)
H25	0.9508	0.5991	0.2843	0.106*
C26	0.8809 (9)	0.4432 (7)	0.3339 (8)	0.057 (2)
H26	0.9240	0.4151	0.2753	0.068*
C27	0.7400 (8)	0.4227 (7)	0.4915 (7)	0.0451 (18)
C28	0.6557 (7)	0.3526 (7)	0.5716 (6)	0.0447 (18)
Gd1	0.82752 (3)	0.14792 (3)	0.44144 (2)	0.02604 (13)
N1	1.2997 (8)	-0.0016 (8)	1.0209 (7)	0.072 (2)
N2	0.2724 (6)	0.4320 (6)	-0.0147 (5)	0.0500 (16)
N3	0.6506 (6)	0.2452 (5)	0.5622 (5)	0.0448 (15)
N4	0.8089 (6)	0.3768 (5)	0.4088 (5)	0.0414 (14)
O1	0.9505 (5)	0.1690 (4)	0.5913 (4)	0.0412 (11)
O2	1.1124 (5)	0.0033 (5)	0.6459 (4)	0.0466 (12)
O3	1.3864 (8)	-0.0642 (9)	0.9807 (8)	0.118 (4)
O4	1.3055 (9)	0.0089 (8)	1.1154 (7)	0.113 (3)

O5	0.8641 (6)	0.2486 (6)	1.0881 (5)	0.0647 (16)
H5A	0.7931	0.2735	1.1164	0.097*
O6	0.7481 (8)	0.3226 (11)	0.9241 (7)	0.140 (5)
O7	0.6825 (4)	0.2498 (4)	0.2637 (4)	0.0335 (10)
O8	0.6158 (4)	0.1067 (4)	0.4063 (4)	0.0413 (11)
O9	0.3605 (6)	0.4789 (5)	-0.0694 (5)	0.0593 (15)
O10	0.1591 (7)	0.4607 (8)	-0.0498 (6)	0.103 (3)
C16	0.1297 (6)	0.1363 (6)	0.3374 (5)	0.0300 (13)
O11	0.0114 (4)	0.1968 (4)	0.3252 (4)	0.0421 (11)
O12	0.1665 (5)	0.0297 (4)	0.4038 (4)	0.0448 (12)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.038 (4)	0.044 (4)	0.022 (3)	-0.013 (3)	-0.002 (3)	-0.014 (3)
C2	0.039 (4)	0.037 (3)	0.020 (3)	-0.010 (3)	-0.004 (3)	-0.012 (3)
C3	0.039 (4)	0.048 (4)	0.032 (4)	-0.001 (3)	-0.007 (3)	-0.013 (3)
C4	0.047 (4)	0.059 (5)	0.030 (4)	-0.007 (3)	-0.014 (3)	-0.010 (3)
C5	0.055 (4)	0.052 (4)	0.023 (3)	-0.011 (3)	-0.004 (3)	-0.016 (3)
C6	0.049 (4)	0.053 (4)	0.025 (4)	-0.010 (3)	0.001 (3)	-0.016 (3)
C7	0.043 (4)	0.052 (4)	0.020 (3)	-0.010 (3)	-0.002 (3)	-0.014 (3)
C8	0.055 (5)	0.080 (6)	0.041 (5)	0.001 (4)	0.000 (4)	-0.038 (4)
C9	0.019 (3)	0.044 (4)	0.024 (3)	-0.006 (2)	-0.001 (2)	-0.012 (3)
C10	0.020 (3)	0.039 (3)	0.021 (3)	-0.005 (2)	0.000 (2)	-0.009 (3)
C11	0.025 (3)	0.045 (4)	0.024 (3)	-0.011 (3)	0.004 (2)	-0.006 (3)
C12	0.038 (4)	0.041 (3)	0.017 (3)	-0.015 (3)	0.000 (3)	-0.002 (3)
C13	0.024 (3)	0.046 (4)	0.024 (3)	-0.007 (3)	-0.001 (2)	-0.011 (3)
C14	0.023 (3)	0.035 (3)	0.019 (3)	-0.008 (2)	0.001 (2)	-0.008 (2)
C15	0.027 (3)	0.036 (3)	0.019 (3)	-0.006 (2)	-0.002 (2)	-0.006 (3)
C17	0.054 (5)	0.060 (5)	0.057 (6)	-0.004 (4)	0.025 (4)	-0.014 (4)
C18	0.073 (7)	0.093 (8)	0.062 (7)	0.011 (6)	0.037 (5)	-0.014 (6)
C19	0.070 (7)	0.101 (8)	0.052 (6)	0.010 (6)	0.029 (5)	-0.024 (6)
C20	0.065 (6)	0.072 (6)	0.053 (5)	0.019 (4)	-0.007 (4)	-0.039 (5)
C21	0.084 (8)	0.117 (10)	0.088 (9)	-0.001 (7)	0.008 (7)	-0.071 (8)
C22	0.113 (10)	0.080 (7)	0.116 (10)	0.008 (7)	-0.021 (8)	-0.075 (8)
C23	0.089 (7)	0.057 (5)	0.079 (7)	-0.002 (5)	-0.022 (6)	-0.033 (5)
C24	0.114 (10)	0.056 (6)	0.128 (11)	-0.026 (6)	-0.008 (8)	-0.039 (7)
C25	0.106 (9)	0.049 (5)	0.099 (9)	-0.029 (5)	0.006 (7)	-0.004 (6)
C26	0.072 (6)	0.041 (4)	0.050 (5)	-0.014 (4)	0.002 (4)	-0.006 (4)
C27	0.047 (4)	0.041 (4)	0.047 (5)	0.002 (3)	-0.015 (4)	-0.019 (3)
C28	0.038 (4)	0.054 (5)	0.036 (4)	0.004 (3)	-0.005 (3)	-0.017 (4)
Gd1	0.02026 (18)	0.0381 (2)	0.01841 (18)	-0.00571 (12)	0.00062 (11)	-0.00808 (13)
N1	0.068 (5)	0.094 (6)	0.053 (5)	0.011 (4)	-0.035 (4)	-0.034 (4)
N2	0.049 (4)	0.065 (4)	0.023 (3)	-0.016 (3)	-0.006 (3)	0.006 (3)
N3	0.041 (3)	0.051 (4)	0.036 (3)	0.003 (3)	0.011 (3)	-0.017 (3)
N4	0.046 (3)	0.042 (3)	0.030 (3)	-0.004 (3)	-0.002 (3)	-0.007 (3)
O1	0.052 (3)	0.048 (3)	0.026 (2)	-0.009 (2)	-0.013 (2)	-0.014 (2)
O2	0.043 (3)	0.059 (3)	0.046 (3)	-0.004 (2)	-0.002 (2)	-0.031 (3)

O3	0.082 (5)	0.163 (8)	0.098 (6)	0.055 (6)	-0.056 (5)	-0.072 (6)
O4	0.113 (6)	0.153 (8)	0.066 (5)	0.039 (6)	-0.060 (5)	-0.058 (5)
O5	0.060 (4)	0.104 (5)	0.033 (3)	-0.009 (3)	0.009 (3)	-0.035 (3)
O6	0.092 (6)	0.236 (11)	0.086 (6)	0.077 (7)	-0.041 (5)	-0.112 (7)
O7	0.027 (2)	0.044 (2)	0.024 (2)	-0.0110 (18)	-0.0016 (18)	-0.0002 (19)
O8	0.026 (2)	0.056 (3)	0.030 (3)	-0.014 (2)	-0.0062 (19)	0.006 (2)
O9	0.060 (4)	0.069 (4)	0.031 (3)	-0.023 (3)	0.008 (3)	0.012 (3)
O10	0.065 (5)	0.137 (7)	0.063 (5)	-0.040 (4)	-0.032 (4)	0.044 (4)
C16	0.032 (3)	0.045 (4)	0.014 (3)	-0.013 (3)	0.001 (2)	-0.009 (3)
O11	0.026 (2)	0.056 (3)	0.040 (3)	-0.012 (2)	0.008 (2)	-0.010 (2)
O12	0.040 (3)	0.054 (3)	0.032 (3)	-0.021 (2)	-0.005 (2)	0.004 (2)

Geometric parameters (Å, °)

C1—O2	1.242 (8)	C19—H19	0.9300
C1—O1	1.256 (8)	C20—C28	1.405 (10)
C1—C2	1.500 (8)	C20—C21	1.411 (16)
C2—C3	1.390 (9)	C21—C22	1.322 (18)
C2—C7	1.391 (9)	C21—H21	0.9300
C3—C4	1.390 (10)	C22—C23	1.463 (16)
C3—H3	0.9300	C22—H22	0.9300
C4—C5	1.374 (10)	C23—C24	1.391 (16)
C4—N1	1.487 (10)	C23—C27	1.402 (12)
C5—C6	1.370 (10)	C24—C25	1.338 (16)
C5—H5	0.9300	C24—H24	0.9300
C6—C7	1.399 (9)	C25—C26	1.417 (13)
C6—C8	1.498 (10)	C25—H25	0.9300
C7—H7	0.9300	C26—N4	1.309 (10)
C8—O6	1.187 (10)	C26—H26	0.9300
C8—O5	1.308 (9)	C27—N4	1.380 (9)
C9—O8	1.243 (8)	C27—C28	1.445 (11)
C9—O7	1.268 (7)	C28—N3	1.348 (9)
C9—C10	1.519 (8)	Gd1—O12 ⁱ	2.322 (5)
C10—C11	1.383 (9)	Gd1—O2 ⁱⁱ	2.342 (4)
C10—C15	1.392 (8)	Gd1—O11 ⁱⁱⁱ	2.349 (4)
C11—C12	1.376 (9)	Gd1—O1	2.403 (4)
C11—H11	0.9300	Gd1—O8	2.442 (4)
C12—C13	1.396 (9)	Gd1—O7	2.499 (4)
C12—N2	1.479 (8)	Gd1—N3	2.571 (5)
C13—C14	1.383 (9)	Gd1—N4	2.611 (6)
C13—H13	0.9300	N1—O4	1.204 (9)
C14—C15	1.378 (8)	N1—O3	1.214 (10)
C14—C16	1.498 (8)	N2—O10	1.204 (9)
C15—H15	0.9300	N2—O9	1.220 (8)
C17—N3	1.324 (11)	O2—Gd1 ⁱⁱ	2.342 (4)
C17—C18	1.401 (12)	O5—H5A	0.8200
C17—H17	0.9300	C16—O12	1.252 (8)
C18—C19	1.316 (16)	C16—O11	1.257 (7)

C18—H18	0.9300	O11—Gd1 ^{iv}	2.349 (4)
C19—C20	1.434 (15)	O12—Gd1 ⁱ	2.322 (5)
O2—C1—O1	125.4 (6)	C25—C24—C23	120.8 (10)
O2—C1—C2	116.8 (6)	C25—C24—H24	119.6
O1—C1—C2	117.8 (6)	C23—C24—H24	119.6
C3—C2—C7	120.1 (6)	C24—C25—C26	118.6 (10)
C3—C2—C1	118.6 (6)	C24—C25—H25	120.7
C7—C2—C1	121.3 (6)	C26—C25—H25	120.7
C4—C3—C2	118.1 (6)	N4—C26—C25	123.3 (9)
C4—C3—H3	121.0	N4—C26—H26	118.3
C2—C3—H3	121.0	C25—C26—H26	118.3
C5—C4—C3	122.9 (7)	N4—C27—C23	121.9 (9)
C5—C4—N1	118.7 (6)	N4—C27—C28	118.2 (6)
C3—C4—N1	118.4 (7)	C23—C27—C28	119.9 (8)
C6—C5—C4	118.5 (6)	N3—C28—C20	122.7 (8)
C6—C5—H5	120.8	N3—C28—C27	118.0 (6)
C4—C5—H5	120.8	C20—C28—C27	119.4 (8)
C5—C6—C7	120.7 (7)	O12 ⁱ —Gd1—O2 ⁱⁱ	76.27 (19)
C5—C6—C8	120.9 (6)	O12 ⁱ —Gd1—O11 ⁱⁱⁱ	124.82 (17)
C7—C6—C8	118.3 (7)	O2 ⁱⁱ —Gd1—O11 ⁱⁱⁱ	76.07 (18)
C2—C7—C6	119.7 (6)	O12 ⁱ —Gd1—O1	75.73 (17)
C2—C7—H7	120.1	O2 ⁱⁱ —Gd1—O1	126.15 (17)
C6—C7—H7	120.1	O11 ⁱⁱⁱ —Gd1—O1	83.73 (17)
O6—C8—O5	124.0 (7)	O12 ⁱ —Gd1—O8	80.85 (15)
O6—C8—C6	124.2 (7)	O2 ⁱⁱ —Gd1—O8	75.42 (17)
O5—C8—C6	111.8 (7)	O11 ⁱⁱⁱ —Gd1—O8	134.60 (16)
O8—C9—O7	122.3 (5)	O1—Gd1—O8	141.63 (17)
O8—C9—C10	118.4 (5)	O12 ⁱ —Gd1—O7	132.49 (15)
O7—C9—C10	119.1 (5)	O2 ⁱⁱ —Gd1—O7	81.73 (16)
C11—C10—C15	120.8 (5)	O11 ⁱⁱⁱ —Gd1—O7	88.64 (15)
C11—C10—C9	121.4 (5)	O1—Gd1—O7	147.63 (16)
C15—C10—C9	117.6 (5)	O8—Gd1—O7	52.84 (14)
C12—C11—C10	117.3 (6)	O12 ⁱ —Gd1—N3	84.99 (19)
C12—C11—H11	121.4	O2 ⁱⁱ —Gd1—N3	145.6 (2)
C10—C11—H11	121.4	O11 ⁱⁱⁱ —Gd1—N3	137.47 (19)
C11—C12—C13	123.4 (6)	O1—Gd1—N3	74.72 (18)
C11—C12—N2	119.3 (6)	O8—Gd1—N3	73.32 (19)
C13—C12—N2	117.3 (6)	O7—Gd1—N3	90.53 (17)
C14—C13—C12	117.9 (6)	O12 ⁱ —Gd1—N4	138.41 (18)
C14—C13—H13	121.1	O2 ⁱⁱ —Gd1—N4	144.26 (19)
C12—C13—H13	121.1	O11 ⁱⁱⁱ —Gd1—N4	74.83 (18)
C15—C14—C13	120.1 (5)	O1—Gd1—N4	70.35 (17)
C15—C14—C16	119.7 (5)	O8—Gd1—N4	112.27 (17)
C13—C14—C16	120.2 (5)	O7—Gd1—N4	77.29 (16)
C14—C15—C10	120.5 (6)	N3—Gd1—N4	63.6 (2)
C14—C15—H15	119.8	O4—N1—O3	124.4 (8)
C10—C15—H15	119.8	O4—N1—C4	117.7 (8)

N3—C17—C18	123.5 (10)	O3—N1—C4	117.9 (7)
N3—C17—H17	118.2	O10—N2—O9	122.6 (6)
C18—C17—H17	118.2	O10—N2—C12	119.1 (6)
C19—C18—C17	119.6 (10)	O9—N2—C12	118.3 (6)
C19—C18—H18	120.2	C17—N3—C28	117.6 (6)
C17—C18—H18	120.2	C17—N3—Gd1	121.2 (5)
C18—C19—C20	119.8 (8)	C28—N3—Gd1	119.7 (5)
C18—C19—H19	120.1	C26—N4—C27	117.6 (7)
C20—C19—H19	120.1	C26—N4—Gd1	124.1 (5)
C28—C20—C21	119.8 (10)	C27—N4—Gd1	116.9 (5)
C28—C20—C19	116.8 (9)	C1—O1—Gd1	130.5 (4)
C21—C20—C19	123.4 (9)	C1—O2—Gd1 ⁱⁱ	153.7 (4)
C22—C21—C20	121.3 (10)	C8—O5—H5A	109.5
C22—C21—H21	119.4	C9—O7—Gd1	90.7 (3)
C20—C21—H21	119.4	C9—O8—Gd1	94.0 (4)
C21—C22—C23	122.2 (10)	O12—C16—O11	124.9 (6)
C21—C22—H22	118.9	O12—C16—C14	117.3 (5)
C23—C22—H22	118.9	O11—C16—C14	117.8 (6)
C24—C23—C27	117.7 (10)	C16—O11—Gd1 ^{iv}	127.2 (4)
C24—C23—C22	124.9 (10)	C16—O12—Gd1 ⁱ	161.3 (4)
C27—C23—C22	117.4 (11)		

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+2, -y, -z+1$; (iii) $x+1, y, z$; (iv) $x-1, y, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H5A \cdots O7 ^v	0.82	2.03	2.737 (7)	145

Symmetry code: (v) $x, y, z+1$.