

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Dimethoxyphosphoryl)(furan-2-yl)-methyl 2-(2,4-dichlorophenoxy)acetate

Xiaosong Tan,* Hao Peng and Hongwu He

Key Laboratory of Pesticide and Chemical Biology, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China
Correspondence e-mail: txs3542@mail.ccnu.edu.cn


Received 15 October 2010; accepted 21 October 2010

Key indicators: single-crystal X-ray study; $T = 292$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;
 R factor = 0.049; wR factor = 0.133; data-to-parameter ratio = 18.1.

In the title compound, $\text{C}_{15}\text{H}_{15}\text{Cl}_2\text{O}_7\text{P}$, the benzene and furan rings form a dihedral angle of 73.54 (1)°. In the crystal, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into layers parallel to (100).

Related literature

For the synthesis and biological activity of 1-(substituted phenoxyacetoxy)alkylphosphonate derivatives, see: He *et al.* (2001, 2005); Chen *et al.* (2006). The synthesis and biological activity of the title compound have been discussed by Peng *et al.* (2007).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{15}\text{Cl}_2\text{O}_7\text{P}$
 $M_r = 409.14$
Monoclinic, $P2_1/c$
 $a = 8.5380$ (7) Å

$b = 17.3111$ (14) Å
 $c = 12.4003$ (10) Å
 $\beta = 98.475$ (1)°
 $V = 1812.8$ (3) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.48$ mm⁻¹

$T = 292$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
12335 measured reflections

4135 independent reflections
3158 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.133$
 $S = 1.05$
4135 reflections

228 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C6}-\text{H6}\cdots\text{O5}^i$	0.93	2.50	3.322 (3)	148
$\text{C9}-\text{H9}\cdots\text{O2}^{ii}$	0.98	2.35	3.270 (2)	157
$\text{C14}-\text{H14B}\cdots\text{O5}^i$	0.96	2.44	3.379 (3)	165

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINTE* (Bruker, 2001); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We gratefully acknowledge financial support of this work by the National Basic Research Program of China (grant No. 2010CB126100) and the National Natural Science Foundation of China (grant No. 20772042).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2778).

References

- Bruker (2001). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
Chen, T., Shen, P., Li, Y.-J. & He, H.-W. (2006). *J. Fluorine Chem.* **127**, 291–295.
He, H.-W. & Liu, Z.-J. (2001). *Chin. J. Org. Chem.* **21**, 878–883.
He, H.-W., Wang, T. & Yuan, J.-L. (2005). *J. Organomet. Chem.* **690**, 2608–2613.
Peng, H., Wang, T., Xie, P., Chen, T., He, H.-W. & Wan, J. (2007). *J. Agric. Food Chem.* **55**, 1871–1880.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2010). E66, o2962 [https://doi.org/10.1107/S1600536810042820]

(Dimethoxyphosphoryl)(furan-2-yl)methyl 2-(2,4-dichlorophenoxy)acetate**Xiaosong Tan, Hao Peng and Hongwu He****S1. Comment**

Phosphonate derivatives are particularly important in connection with their remarkable biological activities. They have been widely used as enzyme inhibitors, antibacterial agents, anti-HIV agents, and also used as pesticides. As a continuation of our search for novel herbicides, series of 1-(substituted phenoxyacetoxy)alkylphosphonate derivatives have been designed and synthesized. Some of them have shown good herbicidal activities acting as inhibitors of PDHc (He *et al.*, 2001, 2005; Chen *et al.*, 2006; Peng *et al.*, 2007). The title compound (I) can be used as herbicide to control broadleaf weeds and sedge weeds effectively and showed good selectivity between monocotyledonous crops and dicotyledonous weeds. Here, we report the crystal structure of (I).

The title compound (Fig. 1), has formed a racemate crystal with monoclinic ($P2_1/c$) symmetry. The benzene and furan rings in the molecule are nonplanar, and the dihedral angle between the two rings is $73.54(1)^\circ$.


In the crystal structure, weak intermolecular C—H \cdots O hydrogen bonds (Table 1) link the molecules into layers parallel to the (100) plane. Weak C—H $\cdots\pi$ interaction (C15 \cdots Cg1 = 3.690(3) Å, C15—H15 \cdots Cg1 = 137° , symmetry code: x, y, z , Cg1 is the centroid defined by atoms O4/C10—C13) was also observed in the crystal structure.

S2. Experimental


A solution of 2,4-dichlorophenoxyacetyl chloride (11 mmol) in trichloromethane (15 ml) was added to stirred mixture of 1-hydroxy (furan-2-yl)methylphosphonate (10 mmol) and triethyl amine (11 mmol) in trichloromethane (15 ml) at 273 K. The resultant mixture was stirred at ambient temperature for 3 h, then washed with dilute hydrochloric acid solution, saturated sodium hydrogen carbonate and brine separately, dried and evaporated. The residue was chromatographed on silica with acetone and petroleum ether as eluent to give the title compound as a white solid.

S3. Refinement

All hydrogen atoms were geometrically positioned [C—H 0.93–0.98 Å], and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5 U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of the title compound, showing the atom-labeling scheme for the non-H atoms and 50% probability displacement ellipsoids.

**Figure 2**

Part of the crystal packing, showing the C—H...O hydrogen bonds as dashed lines.

(Dimethoxyphosphoryl)(furan-2-yl)methyl 2-(2,4-dichlorophenoxy)acetate

Crystal data

C₁₅H₁₅Cl₂O₇P $M_r = 409.14$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 8.5380$ (7) Å $b = 17.3111$ (14) Å $c = 12.4003$ (10) Å $\beta = 98.475$ (1)° $V = 1812.8$ (3) Å³ $Z = 4$ $F(000) = 840$ $D_x = 1.499$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4066 reflections

 $\theta = 2.4$ – 28.2 ° $\mu = 0.48$ mm⁻¹ $T = 292$ K

Block, colourless

 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ϕ and ω scans

12335 measured reflections

4135 independent reflections

3158 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.048$ $\theta_{\text{max}} = 27.5$ °, $\theta_{\text{min}} = 2.0$ ° $h = -11 \rightarrow 11$ $k = -22 \rightarrow 19$ $l = -15 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$ $wR(F^2) = 0.133$ $S = 1.05$

4135 reflections

228 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0672P)^2 + 0.2806P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.001$ $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6128 (3)	0.35463 (14)	0.10119 (19)	0.0489 (5)
C2	0.4929 (3)	0.40648 (14)	0.10635 (19)	0.0545 (6)
H2	0.4350	0.4262	0.0429	0.065*
C3	0.4592 (3)	0.42900 (14)	0.20734 (19)	0.0474 (5)
C4	0.5448 (2)	0.40033 (12)	0.30291 (17)	0.0377 (4)

C5	0.6644 (2)	0.34767 (12)	0.29467 (18)	0.0411 (5)
H5	0.7220	0.3274	0.3578	0.049*
C6	0.7000 (3)	0.32463 (13)	0.1940 (2)	0.0459 (5)
H6	0.7812	0.2896	0.1892	0.055*
C7	0.5786 (2)	0.39588 (13)	0.49683 (17)	0.0412 (5)
H7A	0.5747	0.3400	0.4922	0.049*
H7B	0.5227	0.4116	0.5559	0.049*
C8	0.7491 (2)	0.42157 (12)	0.52171 (16)	0.0361 (4)
C9	0.9975 (2)	0.39109 (12)	0.62871 (16)	0.0361 (4)
H9	1.0269	0.4344	0.5849	0.043*
C10	1.0363 (2)	0.41187 (12)	0.74493 (17)	0.0390 (5)
C11	1.1050 (3)	0.47563 (13)	0.78872 (19)	0.0449 (5)
H11	1.1352	0.5186	0.7517	0.054*
C12	1.1229 (3)	0.46505 (18)	0.9034 (2)	0.0631 (7)
H12	1.1684	0.4998	0.9560	0.076*
C13	1.0633 (3)	0.39691 (18)	0.9214 (2)	0.0642 (7)
H13	1.0593	0.3757	0.9898	0.077*
C14	1.1220 (4)	0.35140 (18)	0.3883 (2)	0.0681 (8)
H14A	1.0862	0.4032	0.3976	0.102*
H14B	1.0867	0.3348	0.3149	0.102*
H14C	1.2355	0.3499	0.4026	0.102*
C15	1.3707 (3)	0.3126 (2)	0.7176 (2)	0.0762 (9)
H15A	1.3491	0.3517	0.7684	0.114*
H15B	1.4810	0.3138	0.7104	0.114*
H15C	1.3442	0.2628	0.7438	0.114*
Cl1	0.65651 (11)	0.32723 (5)	−0.02623 (6)	0.0777 (3)
Cl2	0.30927 (9)	0.49516 (5)	0.21544 (6)	0.0819 (3)
O1	0.50190 (16)	0.42765 (9)	0.39781 (12)	0.0444 (4)
O2	0.80530 (19)	0.47377 (9)	0.47743 (14)	0.0548 (4)
O3	0.82710 (15)	0.37669 (8)	0.59892 (11)	0.0394 (3)
O4	1.0077 (2)	0.36170 (10)	0.82338 (14)	0.0610 (5)
O5	1.0513 (2)	0.23366 (9)	0.64065 (14)	0.0552 (4)
O6	1.0582 (2)	0.30095 (10)	0.46306 (13)	0.0546 (4)
O7	1.27695 (17)	0.32711 (10)	0.61288 (14)	0.0510 (4)
P1	1.09650 (6)	0.30439 (3)	0.58986 (5)	0.04063 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0557 (13)	0.0517 (14)	0.0399 (12)	−0.0049 (11)	0.0090 (10)	−0.0065 (10)
C2	0.0591 (14)	0.0644 (16)	0.0368 (13)	0.0079 (12)	−0.0036 (11)	0.0024 (11)
C3	0.0411 (11)	0.0545 (13)	0.0442 (13)	0.0110 (10)	−0.0014 (10)	0.0008 (11)
C4	0.0301 (9)	0.0439 (11)	0.0377 (11)	−0.0011 (8)	0.0003 (8)	−0.0009 (9)
C5	0.0352 (10)	0.0446 (12)	0.0415 (12)	0.0010 (9)	−0.0015 (9)	0.0027 (9)
C6	0.0425 (11)	0.0432 (12)	0.0522 (14)	0.0026 (9)	0.0082 (10)	−0.0050 (10)
C7	0.0357 (10)	0.0517 (13)	0.0356 (11)	−0.0018 (9)	0.0031 (8)	0.0030 (9)
C8	0.0373 (10)	0.0402 (11)	0.0301 (10)	−0.0008 (8)	0.0023 (8)	−0.0016 (8)
C9	0.0308 (9)	0.0402 (11)	0.0362 (11)	−0.0047 (8)	0.0018 (8)	0.0062 (9)

C10	0.0356 (10)	0.0443 (12)	0.0369 (11)	0.0040 (9)	0.0046 (9)	0.0037 (9)
C11	0.0509 (13)	0.0404 (12)	0.0436 (13)	-0.0089 (10)	0.0070 (10)	-0.0034 (10)
C12	0.0587 (15)	0.0773 (19)	0.0502 (15)	0.0000 (14)	-0.0022 (12)	-0.0206 (14)
C13	0.0732 (17)	0.084 (2)	0.0352 (13)	0.0139 (16)	0.0064 (12)	0.0061 (13)
C14	0.0799 (19)	0.088 (2)	0.0378 (14)	-0.0214 (16)	0.0133 (13)	0.0000 (13)
C15	0.0442 (14)	0.116 (3)	0.0642 (19)	0.0139 (15)	-0.0057 (13)	0.0160 (17)
C11	0.1028 (6)	0.0855 (5)	0.0477 (4)	0.0091 (4)	0.0213 (4)	-0.0122 (3)
C12	0.0755 (5)	0.1100 (6)	0.0563 (4)	0.0552 (4)	-0.0029 (3)	0.0040 (4)
O1	0.0355 (7)	0.0607 (10)	0.0353 (8)	0.0096 (7)	-0.0004 (6)	-0.0011 (7)
O2	0.0500 (9)	0.0558 (10)	0.0541 (10)	-0.0142 (8)	-0.0072 (7)	0.0202 (8)
O3	0.0322 (7)	0.0458 (8)	0.0389 (8)	-0.0049 (6)	0.0004 (6)	0.0105 (6)
O4	0.0774 (12)	0.0568 (11)	0.0505 (10)	-0.0025 (9)	0.0147 (9)	0.0053 (8)
O5	0.0657 (10)	0.0453 (9)	0.0558 (10)	-0.0021 (8)	0.0123 (8)	0.0066 (8)
O6	0.0633 (10)	0.0619 (10)	0.0387 (9)	-0.0138 (8)	0.0074 (8)	-0.0046 (7)
O7	0.0370 (8)	0.0686 (11)	0.0468 (9)	0.0026 (7)	0.0043 (7)	0.0063 (8)
P1	0.0402 (3)	0.0448 (3)	0.0369 (3)	-0.0021 (2)	0.0060 (2)	0.0002 (2)

Geometric parameters (Å, °)

C1—C2	1.370 (3)	C9—H9	0.9800
C1—C6	1.377 (3)	C10—C11	1.328 (3)
C1—C11	1.742 (2)	C10—O4	1.353 (2)
C2—C3	1.382 (3)	C11—C12	1.419 (3)
C2—H2	0.9300	C11—H11	0.9300
C3—C4	1.389 (3)	C12—C13	1.317 (4)
C3—C12	1.732 (2)	C12—H12	0.9300
C4—O1	1.367 (2)	C13—O4	1.380 (3)
C4—C5	1.384 (3)	C13—H13	0.9300
C5—C6	1.386 (3)	C14—O6	1.438 (3)
C5—H5	0.9300	C14—H14A	0.9600
C6—H6	0.9300	C14—H14B	0.9600
C7—O1	1.415 (2)	C14—H14C	0.9600
C7—C8	1.509 (3)	C15—O7	1.444 (3)
C7—H7A	0.9700	C15—H15A	0.9600
C7—H7B	0.9700	C15—H15B	0.9600
C8—O2	1.194 (2)	C15—H15C	0.9600
C8—O3	1.333 (2)	O5—P1	1.4548 (16)
C9—O3	1.468 (2)	O6—P1	1.5591 (17)
C9—C10	1.475 (3)	O7—P1	1.5749 (16)
C9—P1	1.822 (2)		
C2—C1—C6	121.6 (2)	C11—C10—C9	128.6 (2)
C2—C1—C11	118.84 (19)	O4—C10—C9	120.50 (19)
C6—C1—C11	119.56 (19)	C10—C11—C12	106.2 (2)
C1—C2—C3	118.9 (2)	C10—C11—H11	126.9
C1—C2—H2	120.5	C12—C11—H11	126.9
C3—C2—H2	120.5	C13—C12—C11	107.3 (2)
C2—C3—C4	121.3 (2)	C13—C12—H12	126.3

C2—C3—C12	119.57 (18)	C11—C12—H12	126.3
C4—C3—C12	119.15 (18)	C12—C13—O4	109.7 (2)
O1—C4—C5	125.83 (19)	C12—C13—H13	125.1
O1—C4—C3	115.91 (19)	O4—C13—H13	125.1
C5—C4—C3	118.3 (2)	O6—C14—H14A	109.5
C4—C5—C6	121.2 (2)	O6—C14—H14B	109.5
C4—C5—H5	119.4	H14A—C14—H14B	109.5
C6—C5—H5	119.4	O6—C14—H14C	109.5
C1—C6—C5	118.7 (2)	H14A—C14—H14C	109.5
C1—C6—H6	120.6	H14B—C14—H14C	109.5
C5—C6—H6	120.6	O7—C15—H15A	109.5
O1—C7—C8	111.83 (16)	O7—C15—H15B	109.5
O1—C7—H7A	109.2	H15A—C15—H15B	109.5
C8—C7—H7A	109.2	O7—C15—H15C	109.5
O1—C7—H7B	109.2	H15A—C15—H15C	109.5
C8—C7—H7B	109.2	H15B—C15—H15C	109.5
H7A—C7—H7B	107.9	C4—O1—C7	117.64 (16)
O2—C8—O3	125.38 (18)	C8—O3—C9	116.97 (15)
O2—C8—C7	124.74 (19)	C10—O4—C13	105.9 (2)
O3—C8—C7	109.87 (17)	C14—O6—P1	125.73 (16)
O3—C9—C10	111.03 (15)	C15—O7—P1	121.16 (16)
O3—C9—P1	105.95 (13)	O5—P1—O6	112.02 (9)
C10—C9—P1	114.39 (14)	O5—P1—O7	116.52 (10)
O3—C9—H9	108.4	O6—P1—O7	104.26 (9)
C10—C9—H9	108.4	O5—P1—C9	114.57 (9)
P1—C9—H9	108.4	O6—P1—C9	105.17 (9)
C11—C10—O4	110.8 (2)	O7—P1—C9	103.06 (9)
C6—C1—C2—C3	0.0 (4)	C5—C4—O1—C7	-4.5 (3)
C11—C1—C2—C3	-179.22 (19)	C3—C4—O1—C7	175.69 (18)
C1—C2—C3—C4	0.2 (4)	C8—C7—O1—C4	71.0 (2)
C1—C2—C3—C12	179.42 (19)	O2—C8—O3—C9	-2.4 (3)
C2—C3—C4—O1	179.2 (2)	C7—C8—O3—C9	176.68 (16)
C12—C3—C4—O1	0.0 (3)	C10—C9—O3—C8	119.82 (19)
C2—C3—C4—C5	-0.6 (3)	P1—C9—O3—C8	-115.43 (16)
C12—C3—C4—C5	-179.82 (16)	C11—C10—O4—C13	-0.4 (3)
O1—C4—C5—C6	-178.97 (19)	C9—C10—O4—C13	177.36 (19)
C3—C4—C5—C6	0.8 (3)	C12—C13—O4—C10	0.0 (3)
C2—C1—C6—C5	0.2 (4)	C14—O6—P1—O5	-163.4 (2)
C11—C1—C6—C5	179.39 (16)	C14—O6—P1—O7	-36.6 (2)
C4—C5—C6—C1	-0.6 (3)	C14—O6—P1—C9	71.5 (2)
O1—C7—C8—O2	14.4 (3)	C15—O7—P1—O5	-36.0 (2)
O1—C7—C8—O3	-164.71 (17)	C15—O7—P1—O6	-160.0 (2)
O3—C9—C10—C11	-121.5 (2)	C15—O7—P1—C9	90.4 (2)
P1—C9—C10—C11	118.6 (2)	O3—C9—P1—O5	-57.93 (15)
O3—C9—C10—O4	61.2 (2)	C10—C9—P1—O5	64.71 (17)
P1—C9—C10—O4	-58.7 (2)	O3—C9—P1—O6	65.52 (14)
O4—C10—C11—C12	0.6 (3)	C10—C9—P1—O6	-171.84 (14)

C9—C10—C11—C12	-176.9 (2)	O3—C9—P1—O7	174.48 (12)
C10—C11—C12—C13	-0.6 (3)	C10—C9—P1—O7	-62.88 (16)
C11—C12—C13—O4	0.4 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C6—H6 \cdots O5 ⁱ	0.93	2.50	3.322 (3)	148
C9—H9 \cdots O2 ⁱⁱ	0.98	2.35	3.270 (2)	157
C14—H14 <i>B</i> \cdots O5 ⁱ	0.96	2.44	3.379 (3)	165

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x+2, -y+1, -z+1$.