

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[meso-Tetrakis(4-heptyloxyphenyl)-porphyrinato]nickel(II)

 Liang Chen,^a Hong-Bin Zhao,^{a,b*} Yu-Jia Xie,^a De-Liang Yang^a and Bang-Ying Wang^a

^aDepartment of Organic Chemistry, the College of Chemistry, Xiangtan University, Hunan 411105, People's Republic of China, and ^bEnvironmental Engineering, Dongguan University of Technology, Guangdong 523808, People's Republic of China

Correspondence e-mail: zhhbhanlf@163.com


Received 28 September 2010; accepted 16 October 2010

 Key indicators: single-crystal X-ray study; $T = 185$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.051; wR factor = 0.114; data-to-parameter ratio = 14.8.

In the title compound, $[\text{Ni}(\text{C}_{72}\text{H}_{84}\text{N}_4\text{O}_4)]$, the four-coordinate Ni^{II} ion in the middle of the planar 24-membered porphyrin ring is located on a crystallographic inversion center, with Ni–N distances of 1.946 (2)–1.951 (2) Å. The 4-heptyloxyphenyl groups are twisted with respect to the porphyrin mean plane, the dihedral angles being 88.5 (3) and 79.1 (2)°.

Related literature

For related structures, see: Scheidt (1977); Maclean *et al.* (1996); Jentzen *et al.* (1996). For background to porphyrins and metalloporphyrins, see: Kozaki *et al.* (2007); Kuciauskas *et al.* (1996); Suslick *et al.* (2005); Liu *et al.* (1985); Gross & Ini (1999); Wasielewski *et al.* (1993).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{72}\text{H}_{84}\text{N}_4\text{O}_4)]$
 $M_r = 1128.14$
 Monoclinic, $P2_1/c$
 $a = 15.8843$ (12) Å
 $b = 19.0602$ (15) Å
 $c = 10.2398$ (8) Å
 $\beta = 91.221$ (2)°

$V = 3099.5$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.37$ mm⁻¹
 $T = 185$ K
 $0.21 \times 0.16 \times 0.07$ mm

Data collection

Bruker APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)
 $T_{\text{min}} = 0.927$, $T_{\text{max}} = 0.975$

14663 measured reflections
 5469 independent reflections
 3578 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.062$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.114$
 $S = 0.96$
 5469 reflections

369 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This work was supported by National Analytical Research Center of Electrochemistry and Spectroscopy, Changchun Institute of Applied Chemistry, Changchun, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FK2026).

References

- Bruker (2002). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Gross, Z. & Ini, S. (1999). *Org. Lett.* **1**, 2077–2080.
 Jentzen, W., Turowska-Tyrk, I., Scheidt, W. R. & Shelnutt, J. A. (1996). *Inorg. Chem.* **35**, 3559–3567.
 Kozaki, M., Akita, K. & Okada, K. (2007). *Org. Lett.* **9**, 1509–1592.
 Kuciauskas, D., Lin, S., Seely, G. R., Moore, A. L., Moore, M. T. & Gust, D. (1996). *J. Phys. Chem.* **100**, 15926–15932.
 Liu, H. Y., Abdalmuhdi, I., Chang, C. K. & Anson, F. C. (1985). *J. Phys. Chem.* **89**, 665–670.
 Maclean, A. L., Foran, G. J., Kennedy, B. J., Turner, P. & Hambley, T. W. (1996). *Aust. J. Chem.* **49**, 1273–1278.
 Scheidt, W. R. (1977). *Acc. Chem. Res.* **10**, 339–345.
 Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Suslick, K. S., Bhyrappa, P., Chou, J. H., Kosal, M. E., Nakagaki, S., Smithenry, D. W. & Wilson, S. R. (2005). *Acc. Chem. Res.* **38**, 283–291.
 Wasielewski, M. R., Gaines, G. L., Wiederrecht, G. P., Svec, W. A. & Niemczyk, M. P. (1993). *J. Am. Chem. Soc.* **115**, 10442–10443.

supporting information

Acta Cryst. (2010). E66, m1455 [https://doi.org/10.1107/S1600536810041942]

[meso-Tetrakis(4-heptyloxyphenyl)porphyrinato]nickel(II)**Liang Chen, Hong-Bin Zhao, Yu-Jia Xie, De-Liang Yang and Bang-Ying Wang****S1. Comment**

Porphyrins and metalloporphyrins are researched in many aspects, such as electron and energy transfer (Kozaki *et al.*,2007; Kuciauskas *et al.*,1996), molecular recognition (Suslick *et al.*,2005), catalysts (Liu *et al.*,1985; Gross & Ini, 1999) or biomimetic models of photosynthetic systems (Wasielewski *et al.*,1993). In this paper, the structure of Nickel(II) meso-tetrakis[*p*-(heptyloxy)phenyl] porphyrinate is reported.

The 24-membered porphyrin moiety of the title compound is planar with a mean deviation of 0.045 (3) Å. The four-coordinate Ni^{II} ion is located at a crystallographic inversion center, with Ni—N distances of 1.946 (2) to 1.951 (2) Å, in agreement with those found in other nickel porphyrin compounds (Scheidt,1977; Maclean *et al.*1996; Jentzen *et al.*1996.).

The *p*-heptyloxyphenyl groups are rotated at angles of 88.5 (3)° and 79.1 (2)° with respect to the porphyrin mean plane, due to steric hindrance with the pyrrole-H atoms of the macrocycle.

S2. Experimental

Single crystals were recrystallised from a dichloromethane solution at room temperature.

S3. Refinement

H atoms were placed in calculated positions (C—H = 0.95, 0.98 or 0.99 Å) and refined in riding mode, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl and 1.2 for all other H atoms.




Figure 1

Molecular structure of (I), with the atom-labeling scheme and 50% probability displacement ellipsoids.

[meso-Tetrakis(4-heptyloxyphenyl)porphyrinato]nickel(II)

Crystal data

[Ni(C₇₂H₈₄N₄O₄)]

$M_r = 1128.14$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 15.8843 (12) \text{ \AA}$

$b = 19.0602 (15) \text{ \AA}$

$c = 10.2398 (8) \text{ \AA}$

$\beta = 91.221 (2)^\circ$

$V = 3099.5 (4) \text{ \AA}^3$

$Z = 2$

$F(000) = 1208$

$D_x = 1.209 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1476 reflections

$\theta = 2.3\text{--}22.6^\circ$

$\mu = 0.37 \text{ mm}^{-1}$

$T = 185 \text{ K}$

Block, purple

$0.21 \times 0.16 \times 0.07 \text{ mm}$

Data collection

Bruker APEX CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
phi and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2004)
 $T_{\min} = 0.927$, $T_{\max} = 0.975$

14663 measured reflections
5469 independent reflections
3578 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.062$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.7^\circ$
 $h = -18 \rightarrow 16$
 $k = -22 \rightarrow 22$
 $l = -8 \rightarrow 12$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.114$
 $S = 0.96$
5469 reflections
369 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0487P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	1.0000	0.0000	0.0000	0.02268 (15)
N1	0.93762 (14)	0.08453 (11)	0.0426 (2)	0.0234 (5)
N2	0.90680 (13)	-0.05759 (11)	0.0596 (2)	0.0234 (5)
O1	1.09497 (12)	0.46905 (9)	-0.07128 (19)	0.0323 (5)
O2	0.51784 (12)	0.09270 (11)	0.4285 (2)	0.0408 (6)
C1	1.03791 (18)	0.17651 (14)	-0.0219 (3)	0.0242 (7)
C2	0.96102 (17)	0.15389 (14)	0.0243 (3)	0.0244 (7)
C3	0.89751 (17)	0.20052 (15)	0.0693 (3)	0.0281 (7)
H3	0.8985	0.2503	0.0646	0.034*
C4	0.83648 (18)	0.16108 (15)	0.1191 (3)	0.0301 (7)
H4	0.7866	0.1777	0.1583	0.036*
C5	0.86001 (17)	0.08937 (14)	0.1025 (3)	0.0244 (7)
C6	0.81084 (17)	0.03361 (15)	0.1406 (3)	0.0259 (7)
C7	0.83248 (17)	-0.03534 (15)	0.1162 (3)	0.0267 (7)
C8	0.78014 (18)	-0.09352 (15)	0.1457 (3)	0.0327 (8)
H8	0.7260	-0.0914	0.1832	0.039*

C9	0.82105 (17)	-0.15206 (16)	0.1109 (3)	0.0325 (8)
H9	0.8013	-0.1989	0.1187	0.039*
C10	0.90037 (17)	-0.13037 (14)	0.0595 (3)	0.0243 (7)
C11	1.05300 (17)	0.25403 (14)	-0.0347 (3)	0.0252 (7)
C12	1.01380 (19)	0.29119 (15)	-0.1341 (3)	0.0330 (8)
H12	0.9778	0.2668	-0.1940	0.040*
C13	1.02490 (18)	0.36291 (15)	-0.1501 (3)	0.0333 (8)
H13	0.9960	0.3872	-0.2186	0.040*
C14	1.07832 (17)	0.39857 (14)	-0.0654 (3)	0.0276 (7)
C15	1.11814 (18)	0.36205 (15)	0.0355 (3)	0.0325 (7)
H15	1.1545	0.3863	0.0949	0.039*
C16	1.10554 (18)	0.29106 (15)	0.0505 (3)	0.0334 (8)
H16	1.1333	0.2670	0.1203	0.040*
C17	1.04975 (18)	0.50907 (14)	-0.1690 (3)	0.0321 (7)
H17A	0.9892	0.5107	-0.1485	0.039*
H17B	1.0557	0.4869	-0.2558	0.039*
C18	1.08534 (18)	0.58205 (14)	-0.1706 (3)	0.0326 (7)
H18A	1.0880	0.6005	-0.0802	0.039*
H18B	1.0472	0.6127	-0.2227	0.039*
C19	1.17292 (18)	0.58474 (14)	-0.2277 (3)	0.0339 (8)
H19A	1.2107	0.5542	-0.1746	0.041*
H19B	1.1700	0.5650	-0.3172	0.041*
C20	1.21164 (18)	0.65730 (15)	-0.2341 (3)	0.0360 (8)
H20A	1.2239	0.6742	-0.1442	0.043*
H20B	1.1706	0.6900	-0.2753	0.043*
C21	1.29262 (19)	0.65811 (16)	-0.3114 (3)	0.0384 (8)
H21A	1.3319	0.6231	-0.2729	0.046*
H21B	1.2792	0.6432	-0.4021	0.046*
C22	1.3369 (2)	0.72806 (17)	-0.3155 (4)	0.0528 (10)
H22A	1.3526	0.7424	-0.2253	0.063*
H22B	1.2975	0.7637	-0.3517	0.063*
C23	1.4159 (2)	0.7267 (2)	-0.3977 (4)	0.0689 (12)
H23A	1.4527	0.6883	-0.3680	0.103*
H23B	1.4458	0.7714	-0.3883	0.103*
H23C	1.3999	0.7194	-0.4897	0.103*
C24	0.73300 (17)	0.04826 (14)	0.2166 (3)	0.0266 (7)
C25	0.65543 (18)	0.06211 (16)	0.1569 (3)	0.0376 (8)
H25	0.6505	0.0617	0.0643	0.045*
C26	0.58512 (19)	0.07654 (17)	0.2299 (3)	0.0417 (8)
H26	0.5328	0.0862	0.1871	0.050*
C27	0.59084 (18)	0.07690 (15)	0.3649 (3)	0.0321 (8)
C28	0.66699 (18)	0.06206 (16)	0.4264 (3)	0.0380 (8)
H28	0.6715	0.0615	0.5191	0.046*
C29	0.73707 (19)	0.04797 (16)	0.3511 (3)	0.0363 (8)
H29	0.7893	0.0378	0.3940	0.044*
C30	0.5209 (2)	0.08959 (17)	0.5680 (3)	0.0431 (9)
H30A	0.5677	0.1190	0.6024	0.052*
H30B	0.5308	0.0407	0.5971	0.052*

C31	0.4376 (2)	0.11602 (17)	0.6195 (3)	0.0469 (9)
H31A	0.4344	0.1043	0.7134	0.056*
H31B	0.3910	0.0914	0.5731	0.056*
C32	0.42590 (19)	0.19494 (16)	0.6025 (3)	0.0435 (9)
H32A	0.4785	0.2189	0.6308	0.052*
H32B	0.4164	0.2052	0.5086	0.052*
C33	0.35324 (19)	0.22487 (16)	0.6787 (3)	0.0430 (9)
H33A	0.3588	0.2096	0.7709	0.052*
H33B	0.2997	0.2057	0.6425	0.052*
C34	0.3499 (2)	0.30462 (17)	0.6739 (4)	0.0500 (9)
H34A	0.4052	0.3234	0.7034	0.060*
H34B	0.3404	0.3195	0.5821	0.060*
C35	0.2822 (2)	0.33668 (17)	0.7564 (4)	0.0568 (11)
H35A	0.2896	0.3201	0.8476	0.068*
H35B	0.2264	0.3206	0.7237	0.068*
C36	0.2846 (3)	0.41618 (19)	0.7547 (5)	0.0884 (15)
H36A	0.3389	0.4324	0.7906	0.133*
H36B	0.2390	0.4346	0.8077	0.133*
H36C	0.2775	0.4329	0.6646	0.133*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0215 (3)	0.0189 (3)	0.0279 (3)	−0.0005 (2)	0.0057 (2)	0.0001 (3)
N1	0.0224 (13)	0.0219 (13)	0.0262 (14)	0.0005 (11)	0.0043 (10)	−0.0013 (11)
N2	0.0220 (13)	0.0215 (13)	0.0269 (14)	0.0000 (10)	0.0044 (11)	−0.0015 (11)
O1	0.0413 (13)	0.0198 (11)	0.0359 (13)	−0.0051 (9)	0.0003 (10)	0.0042 (9)
O2	0.0328 (12)	0.0430 (14)	0.0473 (15)	0.0041 (10)	0.0168 (11)	−0.0031 (11)
C1	0.0256 (16)	0.0231 (16)	0.0238 (17)	0.0005 (13)	0.0022 (13)	0.0022 (13)
C2	0.0277 (16)	0.0217 (16)	0.0237 (17)	0.0032 (13)	0.0014 (13)	0.0000 (13)
C3	0.0311 (17)	0.0199 (16)	0.0336 (18)	0.0022 (13)	0.0032 (14)	−0.0009 (13)
C4	0.0271 (17)	0.0261 (17)	0.038 (2)	0.0063 (14)	0.0090 (15)	−0.0013 (14)
C5	0.0225 (16)	0.0208 (16)	0.0302 (17)	0.0005 (13)	0.0040 (13)	−0.0022 (13)
C6	0.0218 (16)	0.0264 (17)	0.0295 (18)	−0.0008 (13)	0.0045 (13)	−0.0022 (14)
C7	0.0230 (16)	0.0245 (17)	0.0330 (18)	−0.0021 (13)	0.0067 (14)	0.0008 (13)
C8	0.0240 (16)	0.0315 (18)	0.043 (2)	−0.0019 (14)	0.0118 (15)	0.0010 (15)
C9	0.0307 (18)	0.0250 (17)	0.042 (2)	−0.0039 (14)	0.0105 (15)	0.0005 (15)
C10	0.0264 (16)	0.0224 (16)	0.0243 (16)	−0.0012 (13)	0.0040 (13)	−0.0002 (13)
C11	0.0245 (15)	0.0206 (15)	0.0308 (17)	−0.0011 (13)	0.0060 (13)	−0.0015 (13)
C12	0.0396 (19)	0.0265 (17)	0.0330 (18)	−0.0060 (14)	0.0016 (15)	−0.0026 (14)
C13	0.0415 (19)	0.0260 (17)	0.0321 (18)	−0.0024 (15)	−0.0036 (15)	0.0037 (14)
C14	0.0297 (17)	0.0214 (15)	0.0318 (18)	−0.0023 (13)	0.0058 (14)	0.0015 (14)
C15	0.0371 (18)	0.0258 (17)	0.0343 (19)	−0.0076 (14)	−0.0072 (15)	−0.0027 (14)
C16	0.0359 (18)	0.0281 (17)	0.0361 (19)	−0.0002 (14)	−0.0031 (15)	0.0068 (15)
C17	0.0352 (17)	0.0246 (17)	0.0366 (18)	0.0001 (14)	0.0020 (14)	0.0064 (15)
C18	0.0372 (18)	0.0242 (17)	0.0367 (19)	0.0024 (14)	0.0072 (15)	0.0033 (14)
C19	0.0361 (18)	0.0227 (17)	0.043 (2)	−0.0001 (14)	0.0035 (15)	0.0013 (15)
C20	0.0364 (19)	0.0295 (17)	0.042 (2)	−0.0007 (15)	0.0066 (16)	0.0028 (15)

C21	0.041 (2)	0.0336 (19)	0.041 (2)	-0.0012 (16)	0.0039 (16)	0.0004 (16)
C22	0.053 (2)	0.037 (2)	0.070 (3)	-0.0073 (18)	0.025 (2)	0.0005 (19)
C23	0.064 (3)	0.063 (3)	0.081 (3)	-0.017 (2)	0.034 (2)	0.001 (2)
C24	0.0243 (16)	0.0194 (15)	0.0365 (19)	0.0005 (12)	0.0073 (14)	-0.0004 (13)
C25	0.0309 (18)	0.045 (2)	0.038 (2)	0.0039 (16)	0.0093 (15)	0.0022 (16)
C26	0.0257 (18)	0.051 (2)	0.049 (2)	0.0065 (15)	0.0046 (16)	0.0006 (18)
C27	0.0255 (17)	0.0257 (17)	0.046 (2)	0.0032 (13)	0.0144 (15)	-0.0016 (15)
C28	0.0343 (19)	0.046 (2)	0.034 (2)	0.0044 (16)	0.0071 (16)	-0.0030 (16)
C29	0.0246 (17)	0.043 (2)	0.041 (2)	0.0055 (15)	0.0044 (15)	-0.0006 (16)
C30	0.045 (2)	0.037 (2)	0.049 (2)	0.0030 (16)	0.0177 (18)	-0.0038 (17)
C31	0.043 (2)	0.041 (2)	0.058 (2)	0.0002 (17)	0.0284 (18)	-0.0068 (18)
C32	0.038 (2)	0.037 (2)	0.056 (2)	-0.0019 (16)	0.0214 (17)	-0.0056 (17)
C33	0.0360 (19)	0.0348 (19)	0.059 (2)	-0.0013 (16)	0.0176 (17)	-0.0069 (17)
C34	0.046 (2)	0.038 (2)	0.066 (3)	0.0018 (17)	0.0165 (19)	-0.0047 (19)
C35	0.045 (2)	0.040 (2)	0.086 (3)	0.0044 (17)	0.015 (2)	-0.013 (2)
C36	0.091 (3)	0.043 (3)	0.132 (4)	0.012 (2)	0.034 (3)	-0.014 (3)

Geometric parameters (Å, °)

Ni1—N1	1.946 (2)	C19—H19A	0.9900
Ni1—N1 ⁱ	1.946 (2)	C19—H19B	0.9900
Ni1—N2 ⁱ	1.951 (2)	C20—C21	1.525 (4)
Ni1—N2	1.951 (2)	C20—H20A	0.9900
N1—C2	1.387 (3)	C20—H20B	0.9900
N1—C5	1.391 (3)	C21—C22	1.508 (4)
N2—C10	1.391 (3)	C21—H21A	0.9900
N2—C7	1.392 (3)	C21—H21B	0.9900
O1—C14	1.371 (3)	C22—C23	1.526 (4)
O1—C17	1.438 (3)	C22—H22A	0.9900
O2—C27	1.376 (3)	C22—H22B	0.9900
O2—C30	1.430 (4)	C23—H23A	0.9800
C1—C10 ⁱ	1.378 (4)	C23—H23B	0.9800
C1—C2	1.388 (4)	C23—H23C	0.9800
C1—C11	1.503 (4)	C24—C29	1.378 (4)
C2—C3	1.428 (4)	C24—C25	1.389 (4)
C3—C4	1.337 (4)	C25—C26	1.384 (4)
C3—H3	0.9500	C25—H25	0.9500
C4—C5	1.428 (4)	C26—C27	1.383 (4)
C4—H4	0.9500	C26—H26	0.9500
C5—C6	1.380 (4)	C27—C28	1.381 (4)
C6—C7	1.382 (4)	C28—C29	1.394 (4)
C6—C24	1.502 (4)	C28—H28	0.9500
C7—C8	1.422 (4)	C29—H29	0.9500
C8—C9	1.343 (4)	C30—C31	1.520 (4)
C8—H8	0.9500	C30—H30A	0.9900
C9—C10	1.437 (4)	C30—H30B	0.9900
C9—H9	0.9500	C31—C32	1.525 (4)
C10—C1 ⁱ	1.378 (4)	C31—H31A	0.9900

C11—C12	1.378 (4)	C31—H31B	0.9900
C11—C16	1.387 (4)	C32—C33	1.518 (4)
C12—C13	1.388 (4)	C32—H32A	0.9900
C12—H12	0.9500	C32—H32B	0.9900
C13—C14	1.380 (4)	C33—C34	1.522 (4)
C13—H13	0.9500	C33—H33A	0.9900
C14—C15	1.387 (4)	C33—H33B	0.9900
C15—C16	1.377 (4)	C34—C35	1.510 (4)
C15—H15	0.9500	C34—H34A	0.9900
C16—H16	0.9500	C34—H34B	0.9900
C17—C18	1.502 (4)	C35—C36	1.516 (5)
C17—H17A	0.9900	C35—H35A	0.9900
C17—H17B	0.9900	C35—H35B	0.9900
C18—C19	1.521 (4)	C36—H36A	0.9800
C18—H18A	0.9900	C36—H36B	0.9800
C18—H18B	0.9900	C36—H36C	0.9800
C19—C20	1.516 (4)		
N1—Ni1—N1 ⁱ	179.996 (1)	C21—C20—H20A	109.1
N1—Ni1—N2 ⁱ	89.88 (9)	C19—C20—H20B	109.1
N1 ⁱ —Ni1—N2 ⁱ	90.13 (9)	C21—C20—H20B	109.1
N1—Ni1—N2	90.13 (9)	H20A—C20—H20B	107.9
N1 ⁱ —Ni1—N2	89.87 (9)	C22—C21—C20	115.0 (3)
N2 ⁱ —Ni1—N2	180.0	C22—C21—H21A	108.5
C2—N1—C5	103.8 (2)	C20—C21—H21A	108.5
C2—N1—Ni1	128.28 (19)	C22—C21—H21B	108.5
C5—N1—Ni1	127.87 (18)	C20—C21—H21B	108.5
C10—N2—C7	104.0 (2)	H21A—C21—H21B	107.5
C10—N2—Ni1	128.07 (18)	C21—C22—C23	112.9 (3)
C7—N2—Ni1	127.92 (18)	C21—C22—H22A	109.0
C14—O1—C17	117.2 (2)	C23—C22—H22A	109.0
C27—O2—C30	116.9 (2)	C21—C22—H22B	109.0
C10 ⁱ —C1—C2	122.2 (2)	C23—C22—H22B	109.0
C10 ⁱ —C1—C11	119.2 (3)	H22A—C22—H22B	107.8
C2—C1—C11	118.6 (2)	C22—C23—H23A	109.5
C1—C2—N1	125.6 (2)	C22—C23—H23B	109.5
C1—C2—C3	123.3 (3)	H23A—C23—H23B	109.5
N1—C2—C3	110.9 (2)	C22—C23—H23C	109.5
C4—C3—C2	107.2 (3)	H23A—C23—H23C	109.5
C4—C3—H3	126.4	H23B—C23—H23C	109.5
C2—C3—H3	126.4	C29—C24—C25	117.6 (3)
C3—C4—C5	107.4 (3)	C29—C24—C6	119.8 (3)
C3—C4—H4	126.3	C25—C24—C6	122.6 (3)
C5—C4—H4	126.3	C24—C25—C26	121.2 (3)
C6—C5—N1	125.8 (2)	C24—C25—H25	119.4
C6—C5—C4	123.5 (3)	C26—C25—H25	119.4
N1—C5—C4	110.6 (2)	C27—C26—C25	120.2 (3)
C5—C6—C7	122.5 (3)	C27—C26—H26	119.9

C5—C6—C24	118.7 (3)	C25—C26—H26	119.9
C7—C6—C24	118.7 (3)	O2—C27—C28	124.6 (3)
C6—C7—N2	125.5 (3)	O2—C27—C26	115.9 (3)
C6—C7—C8	123.7 (3)	C28—C27—C26	119.5 (3)
N2—C7—C8	110.8 (2)	C27—C28—C29	119.3 (3)
C9—C8—C7	107.6 (3)	C27—C28—H28	120.4
C9—C8—H8	126.2	C29—C28—H28	120.4
C7—C8—H8	126.2	C24—C29—C28	122.1 (3)
C8—C9—C10	106.9 (3)	C24—C29—H29	119.0
C8—C9—H9	126.5	C28—C29—H29	119.0
C10—C9—H9	126.5	O2—C30—C31	108.7 (3)
C1 ⁱ —C10—N2	125.8 (3)	O2—C30—H30A	109.9
C1 ⁱ —C10—C9	123.6 (3)	C31—C30—H30A	109.9
N2—C10—C9	110.6 (2)	O2—C30—H30B	109.9
C12—C11—C16	117.3 (3)	C31—C30—H30B	109.9
C12—C11—C1	119.9 (3)	H30A—C30—H30B	108.3
C16—C11—C1	122.7 (3)	C30—C31—C32	113.1 (3)
C11—C12—C13	122.5 (3)	C30—C31—H31A	109.0
C11—C12—H12	118.8	C32—C31—H31A	109.0
C13—C12—H12	118.8	C30—C31—H31B	109.0
C14—C13—C12	119.3 (3)	C32—C31—H31B	109.0
C14—C13—H13	120.4	H31A—C31—H31B	107.8
C12—C13—H13	120.4	C33—C32—C31	113.9 (3)
O1—C14—C13	124.9 (3)	C33—C32—H32A	108.8
O1—C14—C15	116.1 (3)	C31—C32—H32A	108.8
C13—C14—C15	119.1 (3)	C33—C32—H32B	108.8
C16—C15—C14	120.7 (3)	C31—C32—H32B	108.8
C16—C15—H15	119.6	H32A—C32—H32B	107.7
C14—C15—H15	119.6	C32—C33—C34	112.7 (3)
C15—C16—C11	121.1 (3)	C32—C33—H33A	109.1
C15—C16—H16	119.4	C34—C33—H33A	109.1
C11—C16—H16	119.4	C32—C33—H33B	109.1
O1—C17—C18	108.4 (2)	C34—C33—H33B	109.1
O1—C17—H17A	110.0	H33A—C33—H33B	107.8
C18—C17—H17A	110.0	C35—C34—C33	114.3 (3)
O1—C17—H17B	110.0	C35—C34—H34A	108.7
C18—C17—H17B	110.0	C33—C34—H34A	108.7
H17A—C17—H17B	108.4	C35—C34—H34B	108.7
C17—C18—C19	112.5 (2)	C33—C34—H34B	108.7
C17—C18—H18A	109.1	H34A—C34—H34B	107.6
C19—C18—H18A	109.1	C34—C35—C36	112.3 (3)
C17—C18—H18B	109.1	C34—C35—H35A	109.1
C19—C18—H18B	109.1	C36—C35—H35A	109.1
H18A—C18—H18B	107.8	C34—C35—H35B	109.1
C20—C19—C18	115.0 (2)	C36—C35—H35B	109.1
C20—C19—H19A	108.5	H35A—C35—H35B	107.9
C18—C19—H19A	108.5	C35—C36—H36A	109.5
C20—C19—H19B	108.5	C35—C36—H36B	109.5

C18—C19—H19B	108.5	H36A—C36—H36B	109.5
H19A—C19—H19B	107.5	C35—C36—H36C	109.5
C19—C20—C21	112.3 (2)	H36A—C36—H36C	109.5
C19—C20—H20A	109.1	H36B—C36—H36C	109.5

Symmetry code: (i) $-x+2, -y, -z$.