

# Poly[[[1-ethyl-6,8-difluoro-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylato]cadmium]- $\mu$ -benzene-1,4-dicarboxylato] trihydrate]

Xin-Ping Kang,<sup>a,b</sup> Zhe An<sup>b</sup> and Rena Kasimu<sup>a\*</sup>

<sup>a</sup>School of Pharmacological Sciences, Xinjiang Medical University, Urumqi 830054, People's Republic of China, and <sup>b</sup>School of Chemistry and Life Science, Guangdong University of Petrochemical Technology, Maoming 525000, People's Republic of China

Correspondence e-mail: kasimu\_xmu@163.com


Received 21 October 2010; accepted 23 October 2010

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.009$  Å;  $R$  factor = 0.044;  $wR$  factor = 0.135; data-to-parameter ratio = 12.4.

In the title layered coordination polymer,  $\{[\text{Cd}(\text{C}_{17}\text{H}_{18}\text{F}_2\text{N}_3\text{O}_3)(\text{C}_8\text{H}_4\text{O}_4)] \cdot 3\text{H}_2\text{O}\}_n$ , the  $\text{Cd}^{\text{II}}$  atom exhibits a very distorted  $\text{CdO}_6$  octahedral geometry defined by one  $O^3, O^4$ -bidentate 1-ethyl-6,8-difluoro-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate (lome) ligand, one  $O, O'$ -bidentate benzene-1,4-dicarboxylate (bdc) dianion and two  $O$ -monodentate bdc dianions. Both the bdc species in the asymmetric unit are completed by crystallographic inversion symmetry. The bridging bdc dianions link the cadmium nodes into a rectangular grid lying parallel to  $(01\bar{1})$ . A network of  $\text{N}-\text{H} \cdots \text{O}$  and  $\text{O}-\text{H} \cdots \text{O}$  hydrogen bonds helps to establish the packing.

## Related literature

For background on the medicinal uses of lomefloxacin, see: Mizuki *et al.* (1996).



## Experimental

### Crystal data

$[\text{Cd}(\text{C}_{17}\text{H}_{18}\text{F}_2\text{N}_3\text{O}_3)(\text{C}_8\text{H}_4\text{O}_4)] \cdot 3\text{H}_2\text{O}$   
 $M_r = 680.90$   
 Triclinic,  $P\bar{1}$   
 $a = 9.7924$  (7) Å  
 $b = 11.9788$  (8) Å  
 $c = 13.3981$  (9) Å  
 $\alpha = 114.138$  (1)°  
 $\beta = 103.430$  (1)°  
 $\gamma = 100.295$  (1)°  
 $V = 1327.00$  (16) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.90$  mm<sup>-1</sup>  
 $T = 295$  K  
 $0.32 \times 0.24 \times 0.18$  mm

### Data collection

Bruker SMART CCD diffractometer  
 Absorption correction: multi-scan (*SADABS*; Bruker, 2001)  
 $T_{\text{min}} = 0.762$ ,  $T_{\text{max}} = 0.855$   
 6581 measured reflections  
 4624 independent reflections  
 4108 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.014$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$   
 $wR(F^2) = 0.135$   
 $S = 1.11$   
 4624 reflections  
 372 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 1.76$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.81$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cd1—O6	2.213 (4)	Cd1—O1	2.299 (4)
Cd1—O3	2.238 (3)	Cd1—O5	2.322 (4)
Cd1—O7 <sup>i</sup>	2.295 (4)	Cd1—O4	2.510 (4)

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3A} \cdots \text{O8}$	0.86	2.33	2.749 (8)	110
$\text{O8}-\text{H8A} \cdots \text{O2}^{\text{ii}}$	0.85	2.02	2.854 (7)	166
$\text{O8}-\text{H8B} \cdots \text{O4}^{\text{iii}}$	0.85	2.01	2.837 (7)	166
$\text{O9}-\text{H9B} \cdots \text{O10}^{\text{iv}}$	0.85	1.84	2.687 (14)	178
$\text{O9}-\text{H9A} \cdots \text{O5}^{\text{v}}$	0.85	1.93	2.784 (9)	178
$\text{O10}-\text{H5O} \cdots \text{O2}$	0.85	2.13	2.971 (10)	171
$\text{O10}-\text{H6O} \cdots \text{O9}$	0.85	2.42	2.957 (14)	121

Symmetry codes: (ii)  $-x + 2, -y, -z + 1$ ; (iii)  $x, y - 1, z - 1$ ; (iv)  $-x + 1, -y, -z + 2$ ; (v)  $x, y - 1, z$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors acknowledge financial support from Xinjiang Medical University and Guangdong University of Petrochemical Technology.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5699).

## References

- Bruker (2001). *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mizuki, Y., Fujiwara, I. & Yamaguchi, T. (1996). *J. Antimicrob. Chemother.* **37** (Suppl. A), 41–45.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2010). E66, m1490–m1491 [https://doi.org/10.1107/S160053681004328X]

**Poly[[[1-ethyl-6,8-difluoro-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydro-quinoline-3-carboxylato]cadmium]- $\mu$ -benzene-1,4-dicarboxylato] trihydrate]**

**Xin-Ping Kang, Zhe An and Rena Kasimu**

### S1. Comment

Lomefloxacin (*H*-Lome, 1,4-dihydro-6,8-difluoro-1-ethyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinoline carboxylic acid) is a new member of the class of quinolones that is used to treat infections (Mizuki *et al.*, 1996). The metal complexes of lomefloxacin have not been reported.

The title cadmium(II)-containing complex of lomefloxacin, (I), is reported here (Fig. 1).

The structure of (I) is built up from Cd<sup>2+</sup> cations, lome ligands, 1,4-benzenedicarboxylate anions, three uncoordinated water molecules (Fig. 1). The Cd atom exhibits a distorted CdO<sub>6</sub> octahedral geometry, two O atom come from one bidentate O,O-bonded 1,4-dihydro-6,8-difluoro-1-ethyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinoline carboxylic (lome) and four O atom come from three 1,4-benzenedicarboxylic acid molecules.

In title compound (I) form a square grid propagating in (Fig. 2) which 1,4-benzenedicarboxylic acid is bridged ligands.

The components of (I) are linked by O—H $\cdots$ O and O—H $\cdots$ N hydrogen bonds involving all the potential donors, generating a three-dimensional supramolecular network.

### S2. Experimental

A mixture of Cd(NO<sub>3</sub>)<sub>2</sub>·2H<sub>2</sub>O (0.5 mmol), lomefloxacin (0.6 mmol), 1,4-benzenedicarboxylic acid (0.25 mmol), and water (12 ml) was stirred for 30 min in air. The mixture was then transferred to a 25 ml Teflon reactor and kept at 433 K for 72 h under autogenous pressure. Colourless single crystals of (I) suitable for X-ray analysis were obtained from the reaction mixture.

### S3. Refinement

The H on the C atoms and N atoms were positioned geometrically (C—H = 0.93–0.97 Å, N—H = 0.86 Å) and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C or N})$ . The H atoms on water molecules were placed at chemically sensible positions on the basis of hydrogen bonds, but these were not refined, O—H = 0.85 Å and  $U(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ .




Figure 1

The asymmetric unit of (I), expanded to show the Cd coordination, showing 50% displacement ellipsoids. Symmetry code: (i)  $x, y, z$ .




Figure 2

A view of part of a two-dimensional polymeric sheet in (I) showing the square-grid connectivity (lomefloxacin molecule, H atoms and water molecule have been omitted for clarity).

Poly[[[1-ethyl-6,8-difluoro-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylato]cadmium(II)]- $\mu$ -benzene-1,4-dicarboxylato] trihydrate]

Crystal data

[Cd(C<sub>17</sub>H<sub>18</sub>F<sub>2</sub>N<sub>3</sub>O<sub>3</sub>)(C<sub>8</sub>H<sub>4</sub>O<sub>4</sub>)]·3H<sub>2</sub>O

$M_r$  = 680.90

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a$  = 9.7924 (7) Å

$b$  = 11.9788 (8) Å

$c$  = 13.3981 (9) Å

$\alpha$  = 114.138 (1)°

$\beta$  = 103.430 (1)°

$\gamma$  = 100.295 (1)°

$V$  = 1327.00 (16) Å<sup>3</sup>

$Z$  = 2

$F(000)$  = 690

$D_x$  = 1.704 Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 6498 reflections

$\theta$  = 1.8–25.0°

$\mu$  = 0.90 mm<sup>-1</sup>

$T$  = 295 K

Prism, colourless

0.32 × 0.24 × 0.18 mm

Data collection

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan

(SADABS; Bruker, 2001)

$T_{\min}$  = 0.762,  $T_{\max}$  = 0.855

6581 measured reflections

4624 independent reflections

4108 reflections with  $I > 2\sigma(I)$

$R_{\text{int}}$  = 0.014

$\theta_{\max}$  = 25.0°,  $\theta_{\min}$  = 1.8°

$h$  = -11→11

$k$  = -14→13

$l$  = -15→15

Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)]$  = 0.044

$wR(F^2)$  = 0.135

$S$  = 1.11

4624 reflections

372 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.077P)^2 + 3.3034P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max}$  = 0.001

$\Delta\rho_{\max}$  = 1.76 e Å<sup>-3</sup>

$\Delta\rho_{\min}$  = -0.81 e Å<sup>-3</sup>

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.60296 (4)	0.51272 (3)	0.66873 (3)	0.02772 (15)
F1	0.7760 (5)	0.0511 (4)	0.2452 (3)	0.0572 (10)

F2	0.7085 (5)	-0.1943 (3)	0.4378 (3)	0.0528 (9)
N1	0.6840 (5)	0.0195 (4)	0.6257 (4)	0.0319 (9)
N2	0.7709 (6)	-0.1778 (5)	0.2569 (4)	0.0469 (12)
N3	0.7833 (7)	-0.4099 (5)	0.0884 (5)	0.0608 (16)
H3A	0.8174	-0.4574	0.0387	0.073*
O1	0.5949 (5)	0.4247 (4)	0.7917 (3)	0.0419 (9)
O2	0.6845 (5)	0.3373 (4)	0.8987 (3)	0.0499 (11)
O3	0.6497 (5)	0.3294 (3)	0.5729 (3)	0.0384 (9)
O4	0.8541 (4)	0.6499 (4)	0.8163 (4)	0.0457 (10)
O5	0.6668 (4)	0.7265 (4)	0.8070 (4)	0.0501 (11)
O6	0.3731 (4)	0.5173 (4)	0.6448 (4)	0.0480 (10)
O7	0.3361 (4)	0.4186 (4)	0.4565 (3)	0.0381 (9)
O8	1.0201 (6)	-0.3309 (6)	0.0299 (5)	0.0805 (17)
H8A	1.1082	-0.3335	0.0399	0.121*
H8B	0.9839	-0.3370	-0.0370	0.121*
O9	0.4490 (9)	-0.1577 (9)	0.8478 (8)	0.147 (4)
H9A	0.5155	-0.1934	0.8367	0.220*
H9B	0.4402	-0.1481	0.9123	0.220*
O10	0.5836 (11)	0.1231 (9)	0.9489 (9)	0.142 (3)
H5O	0.6181	0.1897	0.9428	0.213*
H6O	0.5537	0.0588	0.8811	0.213*
C1	0.6442 (6)	0.3397 (5)	0.8035 (4)	0.0346 (12)
C2	0.6576 (6)	0.2330 (5)	0.6991 (4)	0.0303 (11)
C3	0.6622 (6)	0.2386 (5)	0.5956 (4)	0.0302 (11)
C4	0.6902 (6)	0.1292 (5)	0.5103 (4)	0.0294 (10)
C5	0.7112 (6)	0.1352 (5)	0.4121 (5)	0.0360 (12)
H5	0.7033	0.2059	0.4011	0.043*
C6	0.7428 (7)	0.0384 (6)	0.3335 (5)	0.0396 (13)
C7	0.7459 (6)	-0.0757 (5)	0.3398 (4)	0.0363 (12)
C8	0.7193 (6)	-0.0807 (5)	0.4357 (4)	0.0344 (12)
C9	0.6984 (6)	0.0206 (5)	0.5248 (4)	0.0298 (10)
C10	0.6692 (6)	0.1241 (5)	0.7075 (4)	0.0324 (11)
H10	0.6666	0.1227	0.7760	0.039*
C11	0.7087 (7)	-0.0836 (5)	0.6569 (5)	0.0432 (14)
H11A	0.6522	-0.1672	0.5909	0.052*
H11B	0.6741	-0.0756	0.7210	0.052*
C12	0.8695 (7)	-0.0743 (6)	0.6911 (6)	0.0514 (16)
H12A	0.9038	-0.0823	0.6276	0.077*
H12B	0.8822	-0.1420	0.7096	0.077*
H12C	0.9252	0.0075	0.7578	0.077*
C13	0.7198 (10)	-0.2110 (8)	0.1326 (5)	0.075 (3)
H13A	0.6421	-0.1739	0.1176	0.090*
H13B	0.8009	-0.1747	0.1139	0.090*
C14	0.6651 (10)	-0.3490 (8)	0.0593 (6)	0.077 (2)
H14A	0.6393	-0.3686	-0.0218	0.092*
H14B	0.5769	-0.3844	0.0712	0.092*
C15	0.8317 (11)	-0.3763 (7)	0.2137 (7)	0.072 (2)
H15	0.7487	-0.4155	0.2302	0.086*

C16	0.8786 (9)	-0.2384 (7)	0.2861 (6)	0.065 (2)
H16A	0.9699	-0.2010	0.2789	0.078*
H16B	0.8990	-0.2196	0.3666	0.078*
C17	0.9593 (11)	-0.4326 (8)	0.2388 (8)	0.076 (3)
H17A	1.0309	-0.4109	0.2061	0.113*
H17B	0.9204	-0.5244	0.2044	0.113*
H17C	1.0053	-0.3972	0.3213	0.113*
C18	0.8017 (6)	0.7408 (5)	0.8452 (4)	0.0354 (12)
C19	0.9057 (6)	0.8760 (5)	0.9286 (4)	0.0316 (11)
C20	0.8504 (6)	0.9742 (5)	0.9815 (5)	0.0378 (12)
H20	0.7497	0.9570	0.9688	0.045*
C21	0.9451 (6)	1.0990 (5)	1.0537 (5)	0.0368 (12)
H21	0.9081	1.1652	1.0898	0.044*
C22	0.2939 (5)	0.4741 (5)	0.5409 (5)	0.0325 (11)
C23	0.1424 (5)	0.4880 (5)	0.5192 (4)	0.0282 (10)
C24	0.1114 (6)	0.5848 (5)	0.6051 (5)	0.0343 (11)
H24	0.1860	0.6422	0.6753	0.041*
C25	-0.0300 (5)	0.5958 (5)	0.5861 (5)	0.0332 (11)
H25	-0.0498	0.6596	0.6442	0.040*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cd1	0.0276 (2)	0.0200 (2)	0.0276 (2)	0.00944 (14)	0.00617 (15)	0.00477 (15)
F1	0.098 (3)	0.053 (2)	0.045 (2)	0.041 (2)	0.045 (2)	0.0264 (18)
F2	0.089 (3)	0.0242 (16)	0.0424 (19)	0.0259 (17)	0.0203 (18)	0.0111 (15)
N1	0.041 (2)	0.025 (2)	0.029 (2)	0.0112 (18)	0.0114 (19)	0.0126 (18)
N2	0.069 (3)	0.038 (3)	0.029 (2)	0.032 (3)	0.015 (2)	0.006 (2)
N3	0.093 (5)	0.036 (3)	0.052 (3)	0.032 (3)	0.040 (3)	0.005 (3)
O1	0.062 (3)	0.035 (2)	0.035 (2)	0.0243 (19)	0.0226 (19)	0.0142 (17)
O2	0.077 (3)	0.047 (2)	0.025 (2)	0.030 (2)	0.017 (2)	0.0125 (18)
O3	0.063 (3)	0.0288 (19)	0.035 (2)	0.0267 (18)	0.0252 (19)	0.0163 (17)
O4	0.047 (2)	0.0251 (19)	0.044 (2)	0.0065 (17)	0.0060 (19)	0.0040 (17)
O5	0.035 (2)	0.030 (2)	0.052 (3)	0.0015 (17)	-0.0012 (19)	0.0012 (19)
O6	0.031 (2)	0.063 (3)	0.044 (2)	0.0183 (19)	0.0057 (18)	0.022 (2)
O7	0.036 (2)	0.038 (2)	0.052 (2)	0.0202 (17)	0.0229 (18)	0.0239 (19)
O8	0.065 (3)	0.116 (5)	0.087 (4)	0.031 (3)	0.023 (3)	0.072 (4)
O9	0.108 (6)	0.150 (8)	0.122 (7)	0.065 (6)	0.011 (5)	0.014 (6)
O10	0.150 (8)	0.133 (8)	0.190 (10)	0.070 (6)	0.079 (7)	0.095 (8)
C1	0.041 (3)	0.027 (3)	0.030 (3)	0.011 (2)	0.013 (2)	0.007 (2)
C2	0.035 (3)	0.027 (3)	0.024 (2)	0.011 (2)	0.009 (2)	0.006 (2)
C3	0.035 (3)	0.023 (2)	0.028 (3)	0.012 (2)	0.010 (2)	0.007 (2)
C4	0.036 (3)	0.022 (2)	0.026 (2)	0.012 (2)	0.009 (2)	0.006 (2)
C5	0.052 (3)	0.029 (3)	0.036 (3)	0.022 (2)	0.021 (3)	0.017 (2)
C6	0.055 (3)	0.041 (3)	0.028 (3)	0.023 (3)	0.020 (2)	0.014 (2)
C7	0.046 (3)	0.028 (3)	0.024 (3)	0.018 (2)	0.006 (2)	0.002 (2)
C8	0.046 (3)	0.020 (2)	0.030 (3)	0.015 (2)	0.008 (2)	0.007 (2)
C9	0.036 (3)	0.023 (2)	0.026 (2)	0.011 (2)	0.007 (2)	0.009 (2)

C10	0.044 (3)	0.025 (3)	0.023 (2)	0.011 (2)	0.010 (2)	0.008 (2)
C11	0.072 (4)	0.026 (3)	0.039 (3)	0.019 (3)	0.020 (3)	0.019 (2)
C12	0.064 (4)	0.047 (4)	0.050 (4)	0.029 (3)	0.014 (3)	0.028 (3)
C13	0.100 (6)	0.076 (5)	0.027 (3)	0.057 (5)	0.005 (3)	0.002 (3)
C14	0.101 (6)	0.071 (5)	0.045 (4)	0.048 (5)	0.023 (4)	0.008 (4)
C15	0.109 (7)	0.055 (4)	0.081 (5)	0.047 (5)	0.057 (5)	0.037 (4)
C16	0.084 (5)	0.058 (4)	0.040 (3)	0.050 (4)	0.008 (3)	0.005 (3)
C17	0.127 (7)	0.062 (5)	0.089 (6)	0.069 (5)	0.069 (6)	0.049 (5)
C18	0.043 (3)	0.028 (3)	0.025 (3)	0.004 (2)	0.008 (2)	0.008 (2)
C19	0.036 (3)	0.026 (3)	0.022 (2)	0.003 (2)	0.003 (2)	0.007 (2)
C20	0.032 (3)	0.032 (3)	0.035 (3)	0.003 (2)	0.005 (2)	0.009 (2)
C21	0.041 (3)	0.026 (3)	0.032 (3)	0.010 (2)	0.009 (2)	0.006 (2)
C22	0.027 (3)	0.029 (3)	0.043 (3)	0.007 (2)	0.008 (2)	0.021 (2)
C23	0.025 (2)	0.029 (2)	0.034 (3)	0.010 (2)	0.011 (2)	0.016 (2)
C24	0.027 (3)	0.034 (3)	0.031 (3)	0.005 (2)	0.006 (2)	0.009 (2)
C25	0.029 (3)	0.032 (3)	0.033 (3)	0.013 (2)	0.011 (2)	0.010 (2)

*Geometric parameters (Å, °)*

Cd1—O6	2.213 (4)	C5—H5	0.9300
Cd1—O3	2.238 (3)	C6—C7	1.408 (8)
Cd1—O7 <sup>i</sup>	2.295 (4)	C7—C8	1.391 (8)
Cd1—O1	2.299 (4)	C8—C9	1.398 (7)
Cd1—O5	2.322 (4)	C10—H10	0.9300
Cd1—O4	2.510 (4)	C11—C12	1.504 (9)
F1—C6	1.356 (6)	C11—H11A	0.9700
F2—C8	1.358 (6)	C11—H11B	0.9700
N1—C10	1.345 (6)	C12—H12A	0.9600
N1—C9	1.397 (7)	C12—H12B	0.9600
N1—C11	1.496 (6)	C12—H12C	0.9600
N2—C7	1.383 (7)	C13—C14	1.447 (11)
N2—C16	1.451 (8)	C13—H13A	0.9700
N2—C13	1.477 (8)	C13—H13B	0.9700
N3—C15	1.489 (10)	C14—H14A	0.9700
N3—C14	1.534 (10)	C14—H14B	0.9700
N3—H3A	0.8600	C15—C16	1.447 (10)
O1—C1	1.251 (6)	C15—C17	1.555 (11)
O2—C1	1.258 (7)	C15—H15	0.9800
O3—C3	1.261 (6)	C16—H16A	0.9700
O4—C18	1.243 (7)	C16—H16B	0.9700
O5—C18	1.253 (7)	C17—H17A	0.9600
O6—C22	1.255 (7)	C17—H17B	0.9600
O7—C22	1.261 (6)	C17—H17C	0.9600
O7—Cd1 <sup>i</sup>	2.295 (4)	C18—C19	1.520 (7)
O8—H8A	0.8500	C19—C20	1.380 (8)
O8—H8B	0.8501	C19—C21 <sup>ii</sup>	1.380 (8)
O9—H9A	0.8501	C20—C21	1.391 (8)
O9—H9B	0.8500	C20—H20	0.9300



O10—H5O	0.8523	C21—C19 <sup>ii</sup>	1.380 (8)
O10—H6O	0.8530	C21—H21	0.9300
C1—C2	1.513 (7)	C22—C23	1.497 (7)
C2—C10	1.377 (7)	C23—C25 <sup>iii</sup>	1.391 (7)
C2—C3	1.427 (7)	C23—C24	1.399 (7)
C3—C4	1.466 (7)	C24—C25	1.389 (7)
C4—C9	1.405 (7)	C24—H24	0.9300
C4—C5	1.406 (7)	C25—C23 <sup>iii</sup>	1.391 (7)
C5—C6	1.353 (7)	C25—H25	0.9300
O6—Cd1—O3	120.64 (16)	N1—C11—H11B	109.4
O6—Cd1—O7 <sup>i</sup>	102.79 (15)	C12—C11—H11B	109.4
O3—Cd1—O7 <sup>i</sup>	89.23 (13)	H11A—C11—H11B	108.0
O6—Cd1—O1	91.71 (16)	C11—C12—H12A	109.5
O3—Cd1—O1	79.38 (13)	C11—C12—H12B	109.5
O7 <sup>i</sup> —Cd1—O1	164.82 (13)	H12A—C12—H12B	109.5
O6—Cd1—O5	84.71 (16)	C11—C12—H12C	109.5
O3—Cd1—O5	154.35 (15)	H12A—C12—H12C	109.5
O7 <sup>i</sup> —Cd1—O5	88.65 (16)	H12B—C12—H12C	109.5
O1—Cd1—O5	97.27 (15)	C14—C13—N2	111.1 (7)
O6—Cd1—O4	136.90 (15)	C14—C13—H13A	109.4
O3—Cd1—O4	100.57 (14)	N2—C13—H13A	109.4
O7 <sup>i</sup> —Cd1—O4	88.88 (14)	C14—C13—H13B	109.4
O1—Cd1—O4	83.45 (15)	N2—C13—H13B	109.4
O5—Cd1—O4	53.83 (14)	H13A—C13—H13B	108.0
C10—N1—C9	119.0 (4)	C13—C14—N3	109.9 (7)
C10—N1—C11	117.2 (4)	C13—C14—H14A	109.7
C9—N1—C11	123.0 (4)	N3—C14—H14A	109.7
C7—N2—C16	122.9 (5)	C13—C14—H14B	109.7
C7—N2—C13	121.7 (5)	N3—C14—H14B	109.7
C16—N2—C13	113.6 (5)	H14A—C14—H14B	108.2
C15—N3—C14	111.3 (5)	C16—C15—N3	110.6 (6)
C15—N3—H3A	124.3	C16—C15—C17	110.9 (8)
C14—N3—H3A	124.3	N3—C15—C17	108.5 (6)
C1—O1—Cd1	132.5 (3)	C16—C15—H15	108.9
C3—O3—Cd1	132.3 (3)	N3—C15—H15	108.9
C18—O4—Cd1	87.0 (3)	C17—C15—H15	108.9
C18—O5—Cd1	95.5 (3)	C15—C16—N2	113.7 (6)
C22—O6—Cd1	114.2 (4)	C15—C16—H16A	108.8
C22—O7—Cd1 <sup>i</sup>	129.7 (3)	N2—C16—H16A	108.8
H8A—O8—H8B	108.7	C15—C16—H16B	108.8
H9A—O9—H9B	108.4	N2—C16—H16B	108.8
H5O—O10—H6O	107.2	H16A—C16—H16B	107.7
O1—C1—O2	123.6 (5)	C15—C17—H17A	109.5
O1—C1—C2	119.5 (5)	C15—C17—H17B	109.5
O2—C1—C2	117.0 (5)	H17A—C17—H17B	109.5
C10—C2—C3	118.5 (4)	C15—C17—H17C	109.5
C10—C2—C1	116.5 (4)	H17A—C17—H17C	109.5

C3—C2—C1	125.0 (5)	H17B—C17—H17C	109.5
O3—C3—C2	126.5 (4)	O4—C18—O5	123.0 (5)
O3—C3—C4	117.8 (4)	O4—C18—C19	118.8 (5)
C2—C3—C4	115.6 (4)	O5—C18—C19	118.2 (5)
C9—C4—C5	119.6 (4)	O4—C18—Cd1	66.0 (3)
C9—C4—C3	122.0 (5)	O5—C18—Cd1	57.4 (3)
C5—C4—C3	118.3 (4)	C19—C18—Cd1	170.3 (4)
C6—C5—C4	120.1 (5)	C20—C19—C21 <sup>ii</sup>	120.3 (5)
C6—C5—H5	119.9	C20—C19—C18	120.2 (5)
C4—C5—H5	119.9	C21 <sup>ii</sup> —C19—C18	119.5 (5)
C5—C6—F1	119.3 (5)	C19—C20—C21	120.1 (5)
C5—C6—C7	123.1 (5)	C19—C20—H20	120.0
F1—C6—C7	117.6 (5)	C21—C20—H20	120.0
N2—C7—C8	121.2 (5)	C19 <sup>ii</sup> —C21—C20	119.6 (5)
N2—C7—C6	123.6 (5)	C19 <sup>ii</sup> —C21—H21	120.2
C8—C7—C6	115.2 (5)	C20—C21—H21	120.2
F2—C8—C7	116.0 (4)	O6—C22—O7	122.8 (5)
F2—C8—C9	119.7 (5)	O6—C22—C23	117.1 (5)
C7—C8—C9	124.2 (5)	O7—C22—C23	120.1 (5)
N1—C9—C8	124.1 (5)	C25 <sup>iii</sup> —C23—C24	119.0 (5)
N1—C9—C4	118.6 (4)	C25 <sup>iii</sup> —C23—C22	120.5 (5)
C8—C9—C4	117.3 (5)	C24—C23—C22	120.5 (5)
N1—C10—C2	125.9 (5)	C25—C24—C23	120.5 (5)
N1—C10—H10	117.0	C25—C24—H24	119.8
C2—C10—H10	117.0	C23—C24—H24	119.8
N1—C11—C12	111.1 (5)	C24—C25—C23 <sup>iii</sup>	120.5 (5)
N1—C11—H11A	109.4	C24—C25—H25	119.7
C12—C11—H11A	109.4	C23 <sup>iii</sup> —C25—H25	119.7
O6—Cd1—O1—C1	140.3 (5)	F2—C8—C9—N1	8.1 (8)
O3—Cd1—O1—C1	19.4 (5)	C7—C8—C9—N1	-174.4 (5)
O7 <sup>i</sup> —Cd1—O1—C1	-22.5 (9)	F2—C8—C9—C4	-171.3 (5)
O5—Cd1—O1—C1	-134.8 (5)	C7—C8—C9—C4	6.2 (8)
O4—Cd1—O1—C1	-82.7 (5)	C5—C4—C9—N1	177.0 (5)
C18—Cd1—O1—C1	-108.1 (5)	C3—C4—C9—N1	-2.1 (7)
O6—Cd1—O3—C3	-86.7 (5)	C5—C4—C9—C8	-3.5 (8)
O7 <sup>i</sup> —Cd1—O3—C3	168.9 (5)	C3—C4—C9—C8	177.3 (5)
O1—Cd1—O3—C3	-1.0 (5)	C9—N1—C10—C2	4.2 (8)
O5—Cd1—O3—C3	83.6 (6)	C11—N1—C10—C2	174.6 (5)
O4—Cd1—O3—C3	80.2 (5)	C3—C2—C10—N1	-0.4 (8)
C18—Cd1—O3—C3	83.7 (5)	C1—C2—C10—N1	-179.1 (5)
O6—Cd1—O4—C18	-22.9 (4)	C10—N1—C11—C12	-99.4 (6)
O3—Cd1—O4—C18	173.7 (3)	C9—N1—C11—C12	70.5 (7)
O7 <sup>i</sup> —Cd1—O4—C18	84.7 (3)	C7—N2—C13—C14	-140.9 (7)
O1—Cd1—O4—C18	-108.4 (3)	C16—N2—C13—C14	54.0 (10)
O5—Cd1—O4—C18	-4.4 (3)	N2—C13—C14—N3	-55.0 (9)
O6—Cd1—O5—C18	171.8 (4)	C15—N3—C14—C13	56.4 (9)
O3—Cd1—O5—C18	0.2 (6)	C14—N3—C15—C16	-54.0 (9)

O7 <sup>i</sup> —Cd1—O5—C18	-85.2 (4)	C14—N3—C15—C17	-175.8 (7)
O1—Cd1—O5—C18	80.8 (4)	N3—C15—C16—N2	52.3 (9)
O4—Cd1—O5—C18	4.4 (3)	C17—C15—C16—N2	172.7 (6)
O3—Cd1—O6—C22	-54.1 (4)	C7—N2—C16—C15	142.6 (7)
O7 <sup>i</sup> —Cd1—O6—C22	42.7 (4)	C13—N2—C16—C15	-52.6 (10)
O1—Cd1—O6—C22	-132.8 (4)	Cd1—O4—C18—O5	7.9 (6)
O5—Cd1—O6—C22	130.1 (4)	Cd1—O4—C18—C19	-170.6 (4)
O4—Cd1—O6—C22	145.0 (3)	Cd1—O5—C18—O4	-8.6 (6)
C18—Cd1—O6—C22	134.0 (4)	Cd1—O5—C18—C19	170.0 (4)
Cd1—O1—C1—O2	149.8 (5)	O6—Cd1—C18—O4	163.4 (3)
Cd1—O1—C1—C2	-30.0 (8)	O3—Cd1—C18—O4	-7.8 (4)
O1—C1—C2—C10	-161.2 (5)	O7 <sup>i</sup> —Cd1—C18—O4	-94.2 (3)
O2—C1—C2—C10	18.9 (7)	O1—Cd1—C18—O4	70.6 (3)
O1—C1—C2—C3	20.2 (8)	O5—Cd1—C18—O4	172.1 (6)
O2—C1—C2—C3	-159.6 (5)	O6—Cd1—C18—O5	-8.7 (4)
Cd1—O3—C3—C2	-4.1 (8)	O3—Cd1—C18—O5	-179.9 (3)
Cd1—O3—C3—C4	178.5 (3)	O7 <sup>i</sup> —Cd1—C18—O5	93.7 (4)
C10—C2—C3—O3	178.2 (5)	O1—Cd1—C18—O5	-101.5 (4)
C1—C2—C3—O3	-3.2 (9)	O4—Cd1—C18—O5	-172.1 (6)
C10—C2—C3—C4	-4.3 (7)	O6—Cd1—C18—C19	-74 (2)
C1—C2—C3—C4	174.2 (5)	O3—Cd1—C18—C19	114 (2)
O3—C3—C4—C9	-176.7 (5)	O7 <sup>i</sup> —Cd1—C18—C19	28 (2)
C2—C3—C4—C9	5.6 (7)	O1—Cd1—C18—C19	-167 (2)
O3—C3—C4—C5	4.1 (7)	O5—Cd1—C18—C19	-66 (2)
C2—C3—C4—C5	-173.5 (5)	O4—Cd1—C18—C19	122 (2)
C9—C4—C5—C6	-1.7 (8)	O4—C18—C19—C20	-166.3 (5)
C3—C4—C5—C6	177.5 (5)	O5—C18—C19—C20	15.0 (8)
C4—C5—C6—F1	-174.0 (5)	Cd1—C18—C19—C20	76 (2)
C4—C5—C6—C7	4.7 (9)	O4—C18—C19—C21 <sup>ii</sup>	17.0 (8)
C16—N2—C7—C8	-50.8 (9)	O5—C18—C19—C21 <sup>ii</sup>	-161.7 (5)
C13—N2—C7—C8	145.6 (7)	Cd1—C18—C19—C21 <sup>ii</sup>	-101 (2)
C16—N2—C7—C6	130.3 (7)	C21 <sup>ii</sup> —C19—C20—C21	-0.5 (9)
C13—N2—C7—C6	-33.4 (10)	C18—C19—C20—C21	-177.1 (5)
C5—C6—C7—N2	176.7 (6)	C19—C20—C21—C19 <sup>ii</sup>	0.5 (9)
F1—C6—C7—N2	-4.5 (9)	Cd1—O6—C22—O7	8.1 (7)
C5—C6—C7—C8	-2.3 (9)	Cd1—O6—C22—C23	-173.2 (3)
F1—C6—C7—C8	176.5 (5)	Cd1 <sup>i</sup> —O7—C22—O6	-109.6 (5)
N2—C7—C8—F2	-4.8 (8)	Cd1 <sup>i</sup> —O7—C22—C23	71.8 (6)
C6—C7—C8—F2	174.2 (5)	O6—C22—C23—C25 <sup>iii</sup>	-155.0 (5)
N2—C7—C8—C9	177.6 (5)	O7—C22—C23—C25 <sup>iii</sup>	23.7 (7)
C6—C7—C8—C9	-3.4 (8)	O6—C22—C23—C24	24.4 (7)
C10—N1—C9—C8	177.8 (5)	O7—C22—C23—C24	-156.9 (5)
C11—N1—C9—C8	8.1 (8)	C25 <sup>iii</sup> —C23—C24—C25	1.0 (9)
C10—N1—C9—C4	-2.8 (7)	C22—C23—C24—C25	-178.4 (5)
C11—N1—C9—C4	-172.5 (5)	C23—C24—C25—C23 <sup>iii</sup>	-1.0 (9)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+2, -y+2, -z+2$ ; (iii)  $-x, -y+1, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N3—H3A $\cdots$ O8	0.86	2.33	2.749 (8)	110
O8—H8A $\cdots$ O2 <sup>iv</sup>	0.85	2.02	2.854 (7)	166
O8—H8B $\cdots$ O4 <sup>v</sup>	0.85	2.01	2.837 (7)	166
O9—H9B $\cdots$ O10 <sup>vi</sup>	0.85	1.84	2.687 (14)	178
O9—H9A $\cdots$ O5 <sup>vii</sup>	0.85	1.93	2.784 (9)	178
O10—H5O $\cdots$ O2	0.85	2.13	2.971 (10)	171
O10—H6O $\cdots$ O9	0.85	2.42	2.957 (14)	121

Symmetry codes: (iv)  $-x+2, -y, -z+1$ ; (v)  $x, y-1, z-1$ ; (vi)  $-x+1, -y, -z+2$ ; (vii)  $x, y-1, z$ .