

Poly[di- μ_2 -aqua- μ_4 -chlorido- μ_4 -(2-mercaptopyrimidine-4,6-diolato- κ^4 O:O:O':O')-disodium(I)]

Bao Li, Wen Li, Ling Ye, Guang-Feng Hou and Li-Xin Wu*

State Key Laboratory of Supramolecular Structure and Materials, College of Chemistry, Jilin University, Changchun 130012, People's Republic of China
Correspondence e-mail: wulx@jlu.edu.cn


Received 5 November 2010; accepted 8 November 2010

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.024; wR factor = 0.062; data-to-parameter ratio = 13.8.

In the title coordination polymer, $[\text{Na}_2(\text{C}_4\text{H}_3\text{N}_2\text{O}_2\text{S})\text{Cl}(\text{H}_2\text{O})_2]_n$, the Na^{I} ion lies on a twofold rotation axis and the chloride anion on an inversion center. The Na^{I} ion is six-coordinated by two O atoms from two zwitterionic 2-mercaptopyrimidine-4,6-diolate ligands ($mm2$ symmetry), two water O atoms (m symmetry) and two Cl atoms in a distorted octahedral geometry. Adjacent Na^{I} ions are bridged by an olate group, a water molecule and a chloride anion into a three-dimensional network.

Related literature

For organic-inorganic hybrid compounds with 2-mercaptopyrimidine-4,6-diol derivatives, see: Carballo *et al.* (1996).



Experimental

Crystal data

$[\text{Na}_2(\text{C}_4\text{H}_3\text{N}_2\text{O}_2\text{S})\text{Cl}(\text{H}_2\text{O})_2]$	$V = 982.2(3) \text{ \AA}^3$
$M_r = 260.61$	$Z = 4$
Orthorhombic, <i>Imma</i>	Mo $K\alpha$ radiation
$a = 16.815(3) \text{ \AA}$	$\mu = 0.68 \text{ mm}^{-1}$
$b = 6.5938(13) \text{ \AA}$	$T = 290 \text{ K}$
$c = 8.8587(18) \text{ \AA}$	$0.12 \times 0.11 \times 0.09 \text{ mm}$

Data collection

Rigaku R-Axis RAPID diffractometer	4752 measured reflections
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	633 independent reflections
$T_{\text{min}} = 0.923$, $T_{\text{max}} = 0.942$	581 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.024$	46 parameters
$wR(F^2) = 0.062$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
633 reflections	$\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{Cl1}$	0.86	2.41	3.2457 (15)	165
$\text{O2}-\text{H2A}\cdots\text{O1}^{\text{i}}$	0.89	1.94	2.8164 (19)	169
$\text{O2}-\text{H2B}\cdots\text{S1}^{\text{ii}}$	0.87	2.49	3.3545 (15)	176

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + \frac{5}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

This work was supported financially by the National Basic Research Program of China (grant No. 2007CB808003) and the National Natural Science Foundation of China (grant Nos. 20973082, 20921003, 20703019).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2376).

References

- Carballo, R., Casas, J. S., Garcia-Tasende, M. S., Sanchez, A., Sordo, J. & Vazquez-Lopez, E. M. (1996). *J. Organomet. Chem.* **525**, 49–55.
Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS Inc., The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2010). E66, m1546 [https://doi.org/10.1107/S1600536810045836]

Poly[di- μ_2 -aqua- μ_4 -chlorido- μ_4 -(2-mercaptopyrimidine-4,6-diolato- κ^4 O:O:O':O')-disodium(I)]

Bao Li, Wen Li, Ling Ye, Guang-Feng Hou and Li-Xin Wu

S1. Comment

2-Mercaptopyrimidine-4,6-diol and its derivatives have been used widely to build organic-inorganic hybrids due to their diverse properties (Carballo *et al.* 1996). During the course of on-going crystal engineering studies on nickel salts, we obtained two types of crystals different in color, green and colorless. We herein report the crystal structure of the colorless one, the title compound.

The coordination environment around Na^I ion is shown in Fig. 1, with atom numbering scheme. The Na^I ion is six-coordinated in a distorted octahedral geometry by two O atoms from two 2-mercaptopyrimidine-4,6-diolate ligands, two water O atoms and two Cl atoms. The 2-mercaptopyrimidine-4,6-diolate anion serving as a bridging ligand coordinates to four Na^I ions. The crystal structure is stabilized by N—H \cdots Cl, O—H \cdots O and O—H \cdots S hydrogen bonds (Table 1).

S2. Experimental

2-Mercaptopyrimidine-4,6-diol (1.44 g, 10 mmol) and NiCl₂·6H₂O (3.28 g, 10 mmol) were dissolved in hot water (20 ml) and the pH value was adjusted to about 5 by using dilute NaOH solution with stirring. The mixture was heated for one hour and then cooled to room temperature. The precipitate was washed by dilute HCl and the filtrate was allowed to evaporate at room temperature for two weeks, generating two types of block crystals, one was colorless and the other was green.

S3. Refinement

C- and N-bound H atoms were positioned geometrically (C—H = 0.93, N—H = 0.86 Å) and refined as riding atom, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$. Water H atoms were located in a difference Fourier map and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.




Figure 1

The asymmetric unit of the title compound, with symmetry-related atoms to complete the ligand and Na coordination.

Displacement ellipsoids are drawn at the 30% probability level. [Symmetry codes: (A): 1-x, 1/2-y, z; (B): x, 1-y, 2-z; (C): x, -1/2+y, 2-z; (D): 1/2-x, 1-y, 1/2+z; (E): 1/2-x, -y, 1/2+z; (F) x, -y, 2-z.]

Poly[di- μ_2 -aqua- μ_4 -chlorido- μ_4 -(2-mercaptopyrimidine-4,6-diolato- κ^4 O:O:O':O')-disodium(I)]

Crystal data

[Na₂(C₄H₃N₂O₂S)Cl(H₂O)₂]

$M_r = 260.61$

Orthorhombic, *Imma*

Hall symbol: -I 2b 2

$a = 16.815$ (3) Å

$b = 6.5938$ (13) Å

$c = 8.8587$ (18) Å

$V = 982.2$ (3) Å³

$Z = 4$

$F(000) = 528$

$D_x = 1.762$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4272 reflections

$\theta = 3.3$ – 27.4°

$\mu = 0.68$ mm⁻¹

$T = 290$ K

Block, colorless

$0.12 \times 0.11 \times 0.09$ mm

Data collection

Rigaku R-AXIS RAPID diffractometer	4752 measured reflections
Radiation source: rotation anode	633 independent reflections
Graphite monochromator	581 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.017$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$\theta_{\text{max}} = 27.4^\circ$, $\theta_{\text{min}} = 3.9^\circ$
$T_{\text{min}} = 0.923$, $T_{\text{max}} = 0.942$	$h = -21 \rightarrow 19$
	$k = -8 \rightarrow 8$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.024$	H-atom parameters constrained
$wR(F^2) = 0.062$	$w = 1/[\sigma^2(F_o^2) + (0.0282P)^2 + 1.0461P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
633 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
46 parameters	$\Delta\rho_{\text{max}} = 0.26 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.30 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5000	0.2500	0.7898 (3)	0.0222 (5)
C2	0.42785 (10)	0.2500	1.02887 (18)	0.0174 (3)
C3	0.5000	0.2500	1.1064 (3)	0.0199 (5)
H3	0.5000	0.2500	1.2113	0.024*
Cl1	0.2500	0.2500	0.7500	0.02138 (17)
N1	0.43218 (8)	0.2500	0.87105 (16)	0.0204 (3)
H1	0.3881	0.2500	0.8220	0.025*
Na1	0.25570 (4)	0.5000	1.0000	0.0256 (2)
O1	0.35874 (7)	0.2500	1.08459 (14)	0.0242 (3)
O2	0.16839 (8)	0.2500	1.10168 (15)	0.0294 (3)
H2A	0.1533	0.2500	1.1978	0.044*
H2B	0.1232	0.2500	1.0538	0.044*
S1	0.5000	0.2500	0.60136 (8)	0.0531 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0155 (11)	0.0313 (13)	0.0197 (11)	0.000	0.000	0.000
C2	0.0158 (7)	0.0182 (7)	0.0182 (8)	0.000	0.0019 (6)	0.000
C3	0.0178 (11)	0.0262 (12)	0.0158 (10)	0.000	0.000	0.000
Cl1	0.0201 (3)	0.0286 (3)	0.0154 (3)	0.000	-0.0028 (2)	0.000
N1	0.0127 (7)	0.0314 (8)	0.0172 (7)	0.000	-0.0013 (5)	0.000
Na1	0.0288 (4)	0.0236 (4)	0.0245 (4)	0.000	0.000	-0.0031 (3)
O1	0.0141 (6)	0.0382 (7)	0.0204 (6)	0.000	0.0030 (5)	0.000
O2	0.0206 (6)	0.0449 (8)	0.0225 (6)	0.000	0.0013 (5)	0.000

S1	0.0199 (3)	0.1243 (10)	0.0150 (3)	0.000	0.000	0.000
----	------------	-------------	------------	-------	-------	-------

Geometric parameters (Å, °)

C1—N1	1.3484 (19)	N1—H1	0.8600
C1—S1	1.670 (3)	Na1—O2	2.3842 (11)
C2—O1	1.263 (2)	Na1—O1	2.5062 (11)
C2—C3	1.394 (2)	Na1—Cl1 ⁱ	2.7625 (4)
C2—N1	1.400 (2)	Na1—Na1 ⁱⁱ	3.2969 (6)
C3—H3	0.9300	O2—H2A	0.8885
Cl1—Na1	2.7625 (4)	O2—H2B	0.8701
N1 ⁱⁱⁱ —C1—N1	115.5 (2)	O2—Na1—Cl1 ⁱ	95.07 (3)
N1 ⁱⁱⁱ —C1—S1	122.25 (11)	O2 ⁱⁱ —Na1—Cl1 ⁱ	82.47 (3)
N1—C1—S1	122.25 (11)	O1 ⁱⁱ —Na1—Cl1 ⁱ	82.60 (3)
O1—C2—C3	127.48 (16)	O1—Na1—Cl1 ⁱ	100.18 (3)
O1—C2—N1	115.99 (15)	Cl1—Na1—Cl1 ⁱ	176.03 (3)
C3—C2—N1	116.53 (15)	O2—Na1—Na1 ⁱⁱ	133.74 (3)
C2 ⁱⁱⁱ —C3—C2	121.0 (2)	O2 ⁱⁱ —Na1—Na1 ⁱⁱ	46.26 (3)
C2 ⁱⁱⁱ —C3—H3	119.5	O1 ⁱⁱ —Na1—Na1 ⁱⁱ	48.87 (2)
C2—C3—H3	119.5	O1—Na1—Na1 ⁱⁱ	131.13 (2)
Na1—Cl1—Na1 ^{iv}	180.0	Cl1—Na1—Na1 ⁱⁱ	126.636 (8)
Na1—Cl1—Na1 ^v	73.272 (15)	Cl1 ⁱ —Na1—Na1 ⁱⁱ	53.364 (8)
Na1 ^{iv} —Cl1—Na1 ^v	106.728 (15)	O2—Na1—Na1 ^v	46.26 (3)
Na1—Cl1—Na1 ^{vi}	106.728 (15)	O2 ⁱⁱ —Na1—Na1 ^v	133.74 (3)
Na1 ^{iv} —Cl1—Na1 ^{vi}	73.272 (16)	O1 ⁱⁱ —Na1—Na1 ^v	131.13 (2)
Na1 ^v —Cl1—Na1 ^{vi}	180.0	O1—Na1—Na1 ^v	48.87 (2)
C1—N1—C2	125.23 (16)	Cl1—Na1—Na1 ^v	53.364 (8)
C1—N1—H1	117.4	Cl1 ⁱ —Na1—Na1 ^v	126.636 (8)
C2—N1—H1	117.4	Na1 ⁱⁱ —Na1—Na1 ^v	180.00 (5)
O2—Na1—O2 ⁱⁱ	103.98 (6)	C2—O1—Na1 ^v	121.29 (7)
O2—Na1—O1 ⁱⁱ	173.45 (5)	C2—O1—Na1	121.29 (7)
O2 ⁱⁱ —Na1—O1 ⁱⁱ	81.84 (4)	Na1 ^v —O1—Na1	82.26 (4)
O2—Na1—O1	81.84 (4)	Na1—O2—Na1 ^v	87.48 (5)
O2 ⁱⁱ —Na1—O1	173.45 (5)	Na1—O2—H2A	122.5
O1 ⁱⁱ —Na1—O1	92.53 (5)	Na1 ^v —O2—H2A	122.5
O2—Na1—Cl1	82.47 (3)	Na1—O2—H2B	110.7
O2 ⁱⁱ —Na1—Cl1	95.07 (3)	Na1 ^v —O2—H2B	110.7
O1 ⁱⁱ —Na1—Cl1	100.18 (3)	H2A—O2—H2B	102.6
O1—Na1—Cl1	82.60 (3)		

Symmetry codes: (i) $-x+1/2, -y+1, z+1/2$; (ii) $x, y+1/2, -z+2$; (iii) $-x+1, -y+1/2, z$; (iv) $-x+1/2, -y+1/2, -z+3/2$; (v) $x, y-1/2, -z+2$; (vi) $-x+1/2, -y+1, z-1/2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots Cl1	0.86	2.41	3.2457 (15)	165

O2—H2A···O1 ^{vii}	0.89	1.94	2.8164 (19)	169
O2—H2B···S1 ^{viii}	0.87	2.49	3.3545 (15)	176

Symmetry codes: (vii) $-x+1/2, -y+1/2, -z+5/2$; (viii) $x-1/2, -y+1/2, -z+3/2$.