

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-Allyl-3-methyl-3',5'-diphenylspiro-[quinoxaline-2(1*H*),2'(3'*H*)-[1,3,4]-thiadiazole]

 Caleb Ahoya Anothane,^a Rachid Bouhfid,^a Hafid Zouihri,^a El Mokhtar Essassi^a and Seik Weng Ng^{b*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 9 November 2010; accepted 12 November 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; disorder in main residue; R factor = 0.050; wR factor = 0.164; data-to-parameter ratio = 29.3.

In the title spiro compound, $\text{C}_{25}\text{H}_{22}\text{N}_4\text{S}$, the planar quinoxaline (r.m.s. deviation = 0.070 Å) and planar thiadiazole (r.m.s. deviation = 0.060 Å) ring systems share a common C atom; their mean planes are aligned at 89.7 (1)°. The thiazole ring possesses two aromatic ring substituents and is nearly coplanar with these rings [the dihedral angles between the thiadiazole and phenyl rings are 5.7 (1) and 10.7 (1)°]. The allyl unit is disordered over two positions in a 0.65 (1):0.35 (1) ratio.

Related literature

For pharmacologically active compounds derived from the 1,3-dipolar addition of diphenylnitrilimine to double bonds, see: Ahabchane & Essassi (2000); Canara *et al.* (2004); Ghomsi *et al.* (2004); Mustaphil *et al.* (2005).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{22}\text{N}_4\text{S}$	$\gamma = 89.826$ (1)°
$M_r = 410.53$	$V = 1054.50$ (2) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.9201$ (1) Å	Mo $K\alpha$ radiation
$b = 10.0713$ (1) Å	$\mu = 0.17$ mm ⁻¹
$c = 13.6642$ (2) Å	$T = 293$ K
$\alpha = 78.296$ (1)°	$0.40 \times 0.10 \times 0.10$ mm
$\beta = 81.277$ (1)°	

Data collection

Bruker X8 APEXII diffractometer	38334 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	8514 independent reflections
$T_{\min} = 0.818$, $T_{\max} = 0.862$	5771 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$	28 restraints
$wR(F^2) = 0.164$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.32$ e Å ⁻³
8514 reflections	$\Delta\rho_{\text{min}} = -0.20$ e Å ⁻³
291 parameters	

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2073).

References

- Ahabchane, N. E. & Essassi, E. M. (2000). *J. Soc. Chem. Tunis.* **4**, 753–760.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Canara, H. D., Attar, K., Benchidmi, M., Essassi, E. M. & Garrigues, B. (2004). *Indian J. Chem. Sect. B*, **43**, 660–666.
 Ghomsi, J. N. T., Ahabchane, N. H. & Essassi, E. M. (2004). *Phosphorus Sulfur Silicon Relat. Elem.* **179**, 353–364.
 Mustaphil, N. E. H., Ferfra, S., Essassi, E. M. & Garrigues, B. (2005). *Phosphorus Sulfur Silicon Relat. Elem.* **180**, 2193–2200.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2010). E66, o3227 [https://doi.org/10.1107/S1600536810046994]

1-Allyl-3-methyl-3',5'-diphenylspiro[quinoxaline-2(1*H*),2'(3'*H*)-[1,3,4]thiadiazole]

Caleb Ahoya Anothane, Rachid Bouhfid, Hafid Zouihri, El Mokhtar Essassi and Seik Weng Ng

S1. Comment

In our studies on pharmacologically active compounds, we have used diphenylnitrilimine to undergo 1,3-dipolar additions to double bonds (Ahabchane & Essassi, 2000; Canara *et al.*, 2004; Ghomsi *et al.*, 2004; Mustaphil *et al.*, 2005). In the present study, this compound reacts with an aromatic thione to yield the title spiro compound (Scheme I, Fig. 1). The quinoxaline and the thiadiazole ring systems share a common C atom; their mean planes are aligned at 89.7 (1)°.

S2. Experimental

To a solution of 1-allyl-3-methylquinoxaline-2-thione (1.00 g, 4.62 mmol) and diphenylnitrilimine (1.28 g, 5.55 mmol) in THF (20 ml) was added triethylamine (0.78 ml, 5.55 mmol). The mixture was heated for 24 h. The precipitate was recovered by filtration and was separated by chromatography on silica gel (hexane/ethyl acetate: 9/1). The title compound was obtained as yellow crystals upon evaporation of the solvent.

S3. Refinement

C-bound H-atoms were placed in calculated positions (C—H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U_{\text{eq}}(\text{C})$. The allyl unit is disordered over two positions in a 65 (1):35 (1) ratio.




Figure 1

Displacement ellipsoid plot of $C_{25}H_{22}N_4S$ at the 50% probability level; hydrogen atoms are drawn as arbitrary radius. The minor disorder component is not shown.

1-Allyl-3-methyl-3',5'-diphenylspiro[quinoxaline- 2(1H),2'(3'H)-[1,3,4]thiadiazole]

Crystal data

$C_{25}H_{22}N_4S$

$M_r = 410.53$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.9201(1) \text{ \AA}$

$b = 10.0713(1) \text{ \AA}$

$c = 13.6642(2) \text{ \AA}$

$\alpha = 78.296(1)^\circ$

$\beta = 81.277(1)^\circ$

$\gamma = 89.826(1)^\circ$

$V = 1054.50(2) \text{ \AA}^3$

$Z = 2$

$F(000) = 432$

$D_x = 1.293 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9958 reflections

$\theta = 2.3\text{--}33.0^\circ$

$\mu = 0.17 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Prism, yellow

$0.40 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Bruker X8 APEXII
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

$T_{\min} = 0.818$, $T_{\max} = 0.862$

38334 measured reflections

8514 independent reflections

5771 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 33.9^\circ$, $\theta_{\min} = 2.1^\circ$

$h = -12 \rightarrow 12$

$k = -15 \rightarrow 15$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.050$

$wR(F^2) = 0.164$

$S = 1.03$

8514 reflections

291 parameters

28 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0836P)^2 + 0.1531P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.55728 (4)	0.85155 (3)	0.59838 (2)	0.04746 (11)	
N1	0.50074 (13)	0.71267 (12)	0.79755 (8)	0.0449 (2)	
N2	0.15933 (13)	0.76834 (10)	0.77459 (8)	0.0433 (2)	
N3	0.50993 (16)	0.59101 (10)	0.66315 (9)	0.0507 (3)	
N4	0.64611 (15)	0.60663 (10)	0.58661 (9)	0.0470 (2)	
C1	0.8280 (5)	0.5283 (5)	0.9283 (4)	0.0709 (12)	0.654 (11)
H1A	0.9040	0.5982	0.9290	0.085*	0.654 (11)
H1B	0.8382	0.4425	0.9674	0.085*	0.654 (11)
C2	0.7059 (6)	0.5510 (3)	0.8718 (4)	0.0527 (9)	0.654 (11)
H2	0.6314	0.4796	0.8723	0.063*	0.654 (11)
C1'	0.8401 (10)	0.5022 (8)	0.8777 (11)	0.085 (3)	0.346 (11)
H1'1	0.9090	0.5189	0.8148	0.102*	0.346 (11)
H1'2	0.8650	0.4320	0.9287	0.102*	0.346 (11)
C2'	0.7060 (13)	0.5787 (8)	0.8948 (4)	0.0579 (19)	0.346 (11)
H2'	0.6338	0.5656	0.9565	0.070*	0.346 (11)
C3	0.68125 (17)	0.68768 (17)	0.80623 (11)	0.0584 (4)	
H3A	0.7262	0.7583	0.8348	0.070*	
H3B	0.7456	0.6924	0.7393	0.070*	
C4	0.39793 (16)	0.76799 (11)	0.86969 (9)	0.0406 (2)	
C5	0.4548 (2)	0.79361 (17)	0.95621 (12)	0.0577 (3)	
H5	0.5670	0.7761	0.9662	0.069*	
C6	0.3444 (3)	0.84499 (19)	1.02685 (13)	0.0684 (4)	
H6	0.3840	0.8623	1.0839	0.082*	
C7	0.1772 (3)	0.87124 (17)	1.01491 (12)	0.0649 (4)	
H7	0.1043	0.9053	1.0634	0.078*	
C8	0.11982 (19)	0.84633 (14)	0.93028 (11)	0.0527 (3)	
H8	0.0071	0.8637	0.9216	0.063*	
C9	0.22789 (15)	0.79542 (11)	0.85726 (9)	0.0400 (2)	
C10	0.25755 (16)	0.72698 (12)	0.70458 (10)	0.0432 (2)	
C11	0.1835 (2)	0.69551 (19)	0.61697 (13)	0.0644 (4)	
H11A	0.0655	0.7193	0.6226	0.097*	
H11B	0.1924	0.6004	0.6172	0.097*	
H11C	0.2452	0.7467	0.5549	0.097*	
C12	0.44934 (15)	0.71136 (11)	0.70212 (9)	0.0396 (2)	
C13	0.44801 (17)	0.45774 (11)	0.70791 (10)	0.0446 (3)	
C14	0.3295 (2)	0.43079 (15)	0.79643 (12)	0.0581 (4)	
H14	0.2914	0.5007	0.8287	0.070*	
C15	0.2686 (2)	0.29836 (16)	0.83610 (14)	0.0659 (4)	
H15	0.1879	0.2807	0.8945	0.079*	

C16	0.3252 (2)	0.19377 (15)	0.79093 (16)	0.0700 (5)
H16	0.2842	0.1056	0.8185	0.084*
C17	0.4432 (2)	0.22055 (14)	0.70430 (15)	0.0648 (4)
H17	0.4825	0.1496	0.6736	0.078*
C18	0.50517 (19)	0.35175 (13)	0.66164 (12)	0.0513 (3)
H18	0.5843	0.3684	0.6025	0.062*
C19	0.68575 (16)	0.73197 (11)	0.54723 (9)	0.0412 (2)
C20	0.82935 (16)	0.77223 (12)	0.46545 (9)	0.0411 (2)
C21	0.92074 (19)	0.67363 (16)	0.42316 (11)	0.0535 (3)
H21	0.8911	0.5823	0.4474	0.064*
C22	1.0549 (2)	0.7115 (2)	0.34546 (13)	0.0680 (4)
H22	1.1148	0.6456	0.3169	0.082*
C23	1.1010 (2)	0.8468 (2)	0.30971 (13)	0.0704 (5)
H23	1.1924	0.8716	0.2576	0.085*
C24	1.0122 (2)	0.94469 (19)	0.35095 (13)	0.0668 (4)
H24	1.0432	1.0358	0.3267	0.080*
C25	0.87667 (19)	0.90801 (14)	0.42849 (11)	0.0538 (3)
H25	0.8167	0.9747	0.4561	0.065*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0582 (2)	0.03341 (13)	0.04398 (17)	0.00401 (11)	0.00390 (13)	-0.00090 (10)
N1	0.0345 (5)	0.0579 (6)	0.0390 (5)	0.0047 (4)	-0.0036 (4)	-0.0039 (4)
N2	0.0388 (5)	0.0427 (5)	0.0474 (5)	0.0021 (4)	-0.0070 (4)	-0.0069 (4)
N3	0.0578 (6)	0.0322 (4)	0.0531 (6)	0.0008 (4)	0.0132 (5)	-0.0036 (4)
N4	0.0507 (6)	0.0375 (4)	0.0478 (6)	0.0013 (4)	0.0057 (5)	-0.0070 (4)
C1	0.0616 (17)	0.0673 (18)	0.077 (2)	-0.0016 (14)	-0.0272 (18)	0.0131 (17)
C2	0.0502 (14)	0.0530 (13)	0.0565 (19)	0.0043 (11)	-0.0175 (15)	-0.0084 (13)
C1'	0.071 (4)	0.072 (4)	0.103 (6)	0.016 (3)	-0.015 (4)	0.002 (4)
C2'	0.056 (3)	0.074 (4)	0.047 (3)	0.008 (3)	-0.008 (2)	-0.018 (2)
C3	0.0351 (6)	0.0824 (10)	0.0485 (7)	0.0064 (6)	-0.0035 (5)	0.0055 (7)
C4	0.0419 (6)	0.0393 (5)	0.0373 (5)	-0.0035 (4)	-0.0028 (4)	-0.0028 (4)
C5	0.0560 (8)	0.0696 (9)	0.0488 (7)	-0.0062 (7)	-0.0127 (6)	-0.0113 (6)
C6	0.0862 (12)	0.0760 (10)	0.0476 (8)	-0.0061 (9)	-0.0109 (8)	-0.0226 (7)
C7	0.0810 (11)	0.0617 (8)	0.0523 (8)	0.0085 (8)	0.0023 (7)	-0.0215 (7)
C8	0.0535 (7)	0.0476 (6)	0.0539 (7)	0.0087 (5)	0.0015 (6)	-0.0106 (5)
C9	0.0413 (6)	0.0346 (5)	0.0412 (6)	-0.0003 (4)	-0.0016 (4)	-0.0043 (4)
C10	0.0432 (6)	0.0420 (5)	0.0447 (6)	0.0019 (4)	-0.0094 (5)	-0.0079 (4)
C11	0.0673 (10)	0.0751 (10)	0.0600 (9)	0.0064 (8)	-0.0238 (8)	-0.0249 (8)
C12	0.0429 (6)	0.0347 (4)	0.0386 (5)	0.0029 (4)	-0.0020 (4)	-0.0046 (4)
C13	0.0450 (6)	0.0344 (5)	0.0506 (7)	-0.0022 (4)	-0.0066 (5)	-0.0004 (4)
C14	0.0583 (8)	0.0463 (6)	0.0601 (8)	-0.0068 (6)	0.0047 (7)	0.0009 (6)
C15	0.0617 (9)	0.0574 (8)	0.0671 (9)	-0.0179 (7)	-0.0073 (8)	0.0125 (7)
C16	0.0749 (11)	0.0423 (6)	0.0889 (12)	-0.0179 (7)	-0.0306 (10)	0.0090 (7)
C17	0.0720 (10)	0.0383 (6)	0.0876 (12)	-0.0024 (6)	-0.0275 (9)	-0.0097 (7)
C18	0.0541 (7)	0.0392 (5)	0.0615 (8)	-0.0002 (5)	-0.0122 (6)	-0.0099 (5)
C19	0.0466 (6)	0.0377 (5)	0.0369 (5)	0.0017 (4)	-0.0022 (4)	-0.0051 (4)

C20	0.0414 (6)	0.0463 (6)	0.0342 (5)	-0.0002 (4)	-0.0053 (4)	-0.0053 (4)
C21	0.0511 (7)	0.0577 (7)	0.0533 (7)	0.0037 (6)	-0.0035 (6)	-0.0185 (6)
C22	0.0553 (9)	0.0885 (12)	0.0608 (9)	0.0084 (8)	0.0044 (7)	-0.0265 (9)
C23	0.0495 (8)	0.1031 (14)	0.0499 (8)	-0.0017 (8)	0.0073 (6)	-0.0061 (8)
C24	0.0571 (9)	0.0690 (9)	0.0609 (9)	-0.0085 (7)	0.0044 (7)	0.0082 (7)
C25	0.0539 (8)	0.0477 (6)	0.0519 (7)	-0.0015 (5)	0.0028 (6)	0.0000 (5)

Geometric parameters (Å, °)

S1—C19	1.7585 (12)	C8—C9	1.3917 (17)
S1—C12	1.8872 (12)	C8—H8	0.9300
N1—C4	1.3863 (15)	C10—C11	1.4986 (19)
N1—C12	1.4263 (16)	C10—C12	1.5227 (17)
N1—C3	1.4692 (16)	C11—H11A	0.9600
N2—C10	1.2757 (16)	C11—H11B	0.9600
N2—C9	1.3991 (17)	C11—H11C	0.9600
N3—N4	1.3689 (15)	C13—C18	1.3887 (18)
N3—C13	1.4128 (15)	C13—C14	1.394 (2)
N3—C12	1.4709 (14)	C14—C15	1.391 (2)
N4—C19	1.2874 (15)	C14—H14	0.9300
C1—C2	1.318 (3)	C15—C16	1.367 (3)
C1—H1A	0.9300	C15—H15	0.9300
C1—H1B	0.9300	C16—C17	1.373 (3)
C2—C3	1.512 (3)	C16—H16	0.9300
C2—H2	0.9300	C17—C18	1.390 (2)
C1'—C2'	1.330 (5)	C17—H17	0.9300
C1'—H1'1	0.9300	C18—H18	0.9300
C1'—H1'2	0.9300	C19—C20	1.4629 (17)
C2'—C3	1.496 (4)	C20—C25	1.3896 (18)
C2'—H2'	0.9300	C20—C21	1.3948 (18)
C3—H3A	0.9700	C21—C22	1.378 (2)
C3—H3B	0.9700	C21—H21	0.9300
C4—C5	1.3975 (19)	C22—C23	1.381 (3)
C4—C9	1.4025 (17)	C22—H22	0.9300
C5—C6	1.381 (2)	C23—C24	1.374 (3)
C5—H5	0.9300	C23—H23	0.9300
C6—C7	1.377 (3)	C24—C25	1.382 (2)
C6—H6	0.9300	C24—H24	0.9300
C7—C8	1.373 (2)	C25—H25	0.9300
C7—H7	0.9300		
C19—S1—C12	89.97 (5)	C10—C11—H11A	109.5
C4—N1—C12	120.78 (10)	C10—C11—H11B	109.5
C4—N1—C3	120.12 (11)	H11A—C11—H11B	109.5
C12—N1—C3	117.17 (10)	C10—C11—H11C	109.5
C10—N2—C9	119.02 (11)	H11A—C11—H11C	109.5
N4—N3—C13	117.81 (10)	H11B—C11—H11C	109.5
N4—N3—C12	118.38 (9)	N1—C12—N3	111.72 (10)

C13—N3—C12	123.41 (10)	N1—C12—C10	112.44 (10)
C19—N4—N3	112.75 (10)	N3—C12—C10	111.80 (10)
C2—C1—H1A	120.0	N1—C12—S1	112.47 (8)
C2—C1—H1B	120.0	N3—C12—S1	100.87 (7)
H1A—C1—H1B	120.0	C10—C12—S1	106.89 (8)
C1—C2—C3	123.1 (3)	C18—C13—C14	119.45 (12)
C1—C2—H2	118.5	C18—C13—N3	119.04 (12)
C3—C2—H2	118.5	C14—C13—N3	121.50 (12)
C2'—C1'—H1'1	120.0	C15—C14—C13	119.39 (15)
C2'—C1'—H1'2	120.0	C15—C14—H14	120.3
H1'1—C1'—H1'2	120.0	C13—C14—H14	120.3
C1'—C2'—C3	114.3 (7)	C16—C15—C14	121.27 (17)
C1'—C2'—H2'	122.9	C16—C15—H15	119.4
C3—C2'—H2'	122.9	C14—C15—H15	119.4
N1—C3—C2'	113.4 (4)	C15—C16—C17	119.15 (14)
N1—C3—C2	112.4 (2)	C15—C16—H16	120.4
N1—C3—H3A	109.1	C17—C16—H16	120.4
C2'—C3—H3A	92.9	C16—C17—C18	121.24 (16)
C2—C3—H3A	109.1	C16—C17—H17	119.4
N1—C3—H3B	109.1	C18—C17—H17	119.4
C2'—C3—H3B	122.5	C13—C18—C17	119.49 (15)
C2—C3—H3B	109.1	C13—C18—H18	120.3
H3A—C3—H3B	107.9	C17—C18—H18	120.3
N1—C4—C5	122.78 (12)	N4—C19—C20	122.02 (11)
N1—C4—C9	118.77 (11)	N4—C19—S1	115.81 (9)
C5—C4—C9	118.41 (12)	C20—C19—S1	122.17 (9)
C6—C5—C4	119.98 (15)	C25—C20—C21	119.00 (12)
C6—C5—H5	120.0	C25—C20—C19	121.04 (11)
C4—C5—H5	120.0	C21—C20—C19	119.96 (11)
C7—C6—C5	121.61 (15)	C22—C21—C20	120.01 (15)
C7—C6—H6	119.2	C22—C21—H21	120.0
C5—C6—H6	119.2	C20—C21—H21	120.0
C8—C7—C6	118.93 (14)	C21—C22—C23	120.40 (16)
C8—C7—H7	120.5	C21—C22—H22	119.8
C6—C7—H7	120.5	C23—C22—H22	119.8
C7—C8—C9	120.97 (14)	C24—C23—C22	120.08 (15)
C7—C8—H8	119.5	C24—C23—H23	120.0
C9—C8—H8	119.5	C22—C23—H23	120.0
C8—C9—N2	117.88 (12)	C23—C24—C25	120.06 (16)
C8—C9—C4	120.09 (12)	C23—C24—H24	120.0
N2—C9—C4	121.98 (11)	C25—C24—H24	120.0
N2—C10—C11	119.02 (12)	C24—C25—C20	120.45 (14)
N2—C10—C12	123.99 (11)	C24—C25—H25	119.8
C11—C10—C12	116.96 (12)	C20—C25—H25	119.8
C13—N3—N4—C19	176.13 (12)	C13—N3—C12—S1	-171.95 (12)
C12—N3—N4—C19	-10.90 (18)	N2—C10—C12—N1	15.62 (17)
C4—N1—C3—C2'	67.3 (4)	C11—C10—C12—N1	-166.40 (12)

C12—N1—C3—C2'	-128.4 (4)	N2—C10—C12—N3	142.24 (12)
C4—N1—C3—C2	86.4 (3)	C11—C10—C12—N3	-39.79 (15)
C12—N1—C3—C2	-109.3 (3)	N2—C10—C12—S1	-108.26 (12)
C1'—C2'—C3—N1	144.0 (11)	C11—C10—C12—S1	69.71 (13)
C1'—C2'—C3—C2	53.6 (12)	C19—S1—C12—N1	106.99 (9)
C1—C2—C3—N1	-147.7 (7)	C19—S1—C12—N3	-12.17 (9)
C12—N1—C4—C5	-168.30 (12)	C19—S1—C12—C10	-129.14 (9)
C3—N1—C4—C5	-4.53 (19)	N4—N3—C13—C18	-11.6 (2)
C12—N1—C4—C9	13.84 (17)	C12—N3—C13—C18	175.83 (12)
C3—N1—C4—C9	177.61 (11)	N4—N3—C13—C14	169.30 (14)
N1—C4—C5—C6	-177.99 (14)	C12—N3—C13—C14	-3.3 (2)
C9—C4—C5—C6	-0.1 (2)	C18—C13—C14—C15	-0.9 (2)
C4—C5—C6—C7	0.5 (3)	N3—C13—C14—C15	178.20 (15)
C5—C6—C7—C8	-0.4 (3)	C13—C14—C15—C16	1.2 (3)
C6—C7—C8—C9	0.0 (2)	C14—C15—C16—C17	-0.5 (3)
C7—C8—C9—N2	177.98 (13)	C15—C16—C17—C18	-0.4 (3)
C7—C8—C9—C4	0.4 (2)	C14—C13—C18—C17	0.0 (2)
C10—N2—C9—C8	177.11 (12)	N3—C13—C18—C17	-179.11 (14)
C10—N2—C9—C4	-5.33 (17)	C16—C17—C18—C13	0.6 (2)
N1—C4—C9—C8	177.66 (11)	N3—N4—C19—C20	178.86 (11)
C5—C4—C9—C8	-0.29 (18)	N3—N4—C19—S1	-0.79 (16)
N1—C4—C9—N2	0.16 (17)	C12—S1—C19—N4	8.47 (11)
C5—C4—C9—N2	-177.80 (12)	C12—S1—C19—C20	-171.19 (11)
C9—N2—C10—C11	178.88 (12)	N4—C19—C20—C25	-175.82 (13)
C9—N2—C10—C12	-3.19 (18)	S1—C19—C20—C25	3.81 (18)
C4—N1—C12—N3	-147.06 (11)	N4—C19—C20—C21	4.4 (2)
C3—N1—C12—N3	48.70 (15)	S1—C19—C20—C21	-175.98 (10)
C4—N1—C12—C10	-20.40 (15)	C25—C20—C21—C22	-0.5 (2)
C3—N1—C12—C10	175.36 (11)	C19—C20—C21—C22	179.35 (14)
C4—N1—C12—S1	100.32 (11)	C20—C21—C22—C23	0.7 (3)
C3—N1—C12—S1	-63.91 (13)	C21—C22—C23—C24	-0.6 (3)
N4—N3—C12—N1	-104.20 (13)	C22—C23—C24—C25	0.2 (3)
C13—N3—C12—N1	68.34 (16)	C23—C24—C25—C20	0.1 (3)
N4—N3—C12—C10	128.80 (13)	C21—C20—C25—C24	0.0 (2)
C13—N3—C12—C10	-58.66 (16)	C19—C20—C25—C24	-179.75 (14)
N4—N3—C12—S1	15.50 (14)		
